arXiv:1412.1709v2 [math.AT] 10 Dec 2014

THE HIT PROBLEM FOR THE POLYNOMIAL ALGEBRA OF FOUR VARIABLES

NGUYỄN SUM

Abstract. We study the problem of determining a minimal set of generators for the polynomial algebra $\mathbb{F}_2[x_1, x_2, \ldots, x_k]$ as a module over the mod-2 Steenrod algebra $\mathcal{A}$. In this paper, we give an explicit answer in terms of the monomials for $k = 4$.

1. Introduction

Let E^k be an elementary abelian 2-group of rank k. Denote by BE^k the classifying space of E^k. It may be thought of as the product of k copies of real projective space $\mathbb{R}P^\infty$. We have

$$P_k := H^*(BE^k) = \mathbb{F}_2[x_1, x_2, \ldots, x_k],$$

a polynomial algebra on k generators $x_1, x_2, \ldots, x_k$, each of degree 1. Here the cohomology is taken with coefficients in the prime field $\mathbb{F}_2$ of two elements.

It is well known that P_k is a module over the mod-2 Steenrod algebra $\mathcal{A}$. The action of $\mathcal{A}$ on P_k is determined by the formula

$$Sq^i(x_j) = \begin{cases} x_j, & i = 0, \\ x_j^2, & i = 1, \\ 0, & \text{otherwise}, \end{cases}$$

and the Cartan formula

$$Sq^n(fg) = \sum_{i=0}^{n} Sq^i(f)Sq^{n-i}(g),$$

for $f, g \in P_k$ (see Steenrod [15]).

Many authors study the *hit problem* of determination of the minimal set of generators for P_k as a module over the Steenrod algebra, or equivalently, a basis of $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k$. This problem has first been studied by Peterson [10], Wood [16], Singer [13], Priddy [11], Carlisle-Wood [3], who show its relationship to several classical problems in homotopy theory.

Peterson conjectured in [10] that as a module over the Steenrod algebra $\mathcal{A}$, P_k is generated by monomials in degrees n that satisfy $\alpha(n + k) \leqslant k$,

[1] 2000 *Mathematics Subject Classification.* Primary 55S10; 55S05, 55T15.

[2] *Keywords and phrases:* Steenrod squares, polynomial algebra.

[3] This version is a revision of a preprint of Quy Nhơn University, Việt Nam, 2007.

where $\alpha(n)$ denotes the number of ones in dyadic expansion of n, and proved it for $k \leqslant 2$. The conjecture was established in general by Wood [16]. This is a useful tool for determining $\mathcal{A}$-generators for P_k.

The tensor product $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k$ has explicitly been calculated by Peterson [10] for $k = 2$ and Kameko for $k = 3$ in his thesis [8] and in "generic degrees" for all k by Nam [9]. Singer [14], Silverman [12], Hưng-Nam [6, 7] have showed that many polynomials in P_k are in $\mathcal{A}^+.P_k$, where $\mathcal{A}^+$ denotes the augmentation ideal in $\mathcal{A}$.

Recently, many authors showed their interest in the study of $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k$ in conjunction with the transfer, which was defined by Singer [13]. The transfer is a homomorphism

$$Tr_k : \mathrm{Tor}_k^{\mathcal{A}}(\mathbb{F}_2, \mathbb{F}_2) \longrightarrow (\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k)^{GL_k(\mathbb{F}_2)}.$$

Here $\mathrm{Tor}_k^{\mathcal{A}}(\mathbb{F}_2, \mathbb{F}_2)$ is isomorphic to $\mathrm{Ext}_{\mathcal{A}}^{k}(\mathbb{F}_2, \mathbb{F}_2)$, the E_2 term of the Adams spectral sequence of spheres and $GL_k(\mathbb{F}_2)$ is the general linear group which acts on $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k$ in the usual manner. It was shown that the transfer is an isomorphism for $k = 1, 2$ by Singer in [13] and for $k = 3$ by Boardman in [1]. Bruner-Hà-Hưng [2], Hưng [5] and Ha [4] have studied the transfer for $k = 4, 5$.

One of important tools in Kameko's computation of $\mathcal{A}$-generators for P_3 is the squaring operation

$$Sq_*^0 : (\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k)_n \longrightarrow (\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k)_{\frac{n-k}{2}},$$

which is determined for all $n \geqslant k$ such that $n-k$ is even. Kameko showed in [8] that if $\beta(n) = k$ then Sq_*^0 is an isomorphism of $\mathbb{F}_2$-vector spaces, where

$$\beta(n) = \min\{m \in \mathbb{Z} \,;\ \alpha(n+m) \leqslant m\}.$$

From this and Wood's theorem, the hit problem is reduced to the cases $\beta(n) < k$. Hence, for $k = 4$, it suffices to consider six cases:

$$\begin{aligned}
&1)\ n = 2^{s+1} - 3,\\
&2)\ n = 2^{s+1} - 2,\\
&3)\ n = 2^{s+1} - 1,\\
&4)\ n = 2^{s+t+1} + 2^{s+1} - 3,\\
&5)\ n = 2^{s+t} + 2^{s} - 2,\\
&6)\ n = 2^{s+t+u} + 2^{s+t} + 2^{s} - 3,
\end{aligned}$$

where s, t, u are the positive integers.

In this paper, we explicitly determine $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k$ for $k = 4$. We have

Theorem 1.1. *Let n be an arbitrary positive integer with $\beta(n) < 4$. The dimension of the $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_n$ is given by the following table*

n	$s=1$	$s=2$	$s=3$	$s=4$	$s\geqslant 5$
$2^{s+1}-3$	4	15	35	45	45
$2^{s+1}-2$	6	24	50	70	80
$2^{s+1}-1$	14	35	75	89	85
$2^{s+2}+2^{s+1}-3$	46	94	105	105	105
$2^{s+3}+2^{s+1}-3$	87	135	150	150	150
$2^{s+4}+2^{s+1}-3$	136	180	195	195	195
$2^{s+t+1}+2^{s+1}-3, t\geqslant 4$	150	195	210	210	210
$2^{s+1}+2^s-2$	21	70	116	164	175
$2^{s+2}+2^s-2$	55	126	192	240	255
$2^{s+3}+2^s-2$	73	165	241	285	300
$2^{s+4}+2^s-2$	95	179	255	300	315
$2^{s+5}+2^s-2$	115	175	255	300	315
$2^{s+t}+2^s-2, t\geqslant 6$	125	175	255	300	315
$2^{s+2}+2^{s+1}+2^s-3$	64	120	120	120	120
$2^{s+3}+2^{s+2}+2^s-3$	155	210	210	210	210
$2^{s+t+1}+2^{s+t}+2^s-3, t\geqslant 3$	140	210	210	210	210
$2^{s+3}+2^{s+1}+2^s-3$	140	225	225	225	225
$2^{s+u+1}+2^{s+1}+2^s-3, u\geqslant 3$	120	210	210	210	210
$2^{s+u+2}+2^{s+2}+2^s-3, u\geqslant 2$	225	315	315	315	315
$2^{s+t+u}+2^{s+t}+2^s-3, u\geqslant 2, t\geqslant 3$	210	315	315	315	315.

From this theorem, we see that Kameko's conjecture

$$\sup_n \dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k)_n = \prod_{1\leqslant i\leqslant k} (2^i-1)$$

is true for $k=4$. Furthermore, we have the following conjecture.

Conjecture 1.2. *Let $n = 2^{s_1+s_2+\ldots+s_{k-1}} + 2^{s_1+s_2+\ldots+s_{k-2}} + \ldots + 2^{s_1} - k + 1$ with s_j the positive integer, $j = 1, 2, \ldots, k-1$. If $s_j \geqslant 2$ for all j then*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k)_n = \prod_{1\leqslant i\leqslant k} (2^i-1).$$

This paper is organized as follows.

1. Introduction.
2. Preliminaries.
3. The indecomposables of P_4 in degree $2^{s+1}-3$.
4. The indecomposables of P_4 in degree $2^{s+1}-2$.
5. The indecomposables of P_4 in degree $2^{s+1}-1$.
6. The indecomposables of P_4 in degree $2^{s+t+1}+2^{s+1}-3$.
7. The indecomposables of P_4 in degree $2^{s+t}+2^s-2$.
8. The indecomposables of P_4 in degree $2^{s+t+u}+2^{s+t}+2^s-3$.
9. Final remark.

In Section 2, we recall some results on the admissible monomials in P_k and Kameko's squaring operation. We compute $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4$ in the next sections by describing explicitly a basis of it in terms of the monomials. In each case, we order the monomials in a basis by the product of powers of variables and then by the lexicographical order on the set of σ-sequences of monomials (see Section 2).

Finally, we give in Section 9 some matrices from which we can describe the matrices associated with admissible monomials in P_4.

Acknowledgements. It is a pleasure for me to thank Prof. Nguyễn H. V. Hưng for his valuable suggestions and encouragement. My thanks also go to all colleagues at Department of Mathematics, University of Quy Nhơn for many helpful conversations.

2. Preliminaries

In this section, we recall some results in Kameko [8], Wood [16], Singer [14] on the admissible monomials and the hit monomials in P_k. Recall that an element in P_k is called hit if it belongs to $\mathcal{A}^+.P_k$.

Let $x = x_1^{a_1}x_2^{a_2}\dots x_k^{a_k}$ in P_k. Following Kameko [8], we assign a matrix $(\varepsilon_{ij}(x))$ to monomial x, where $\varepsilon_{ij}(x) = \alpha_{i-1}(a_j)$, the $(i-1)$-th coefficient in dyadic expansion of a_j for $i = 1, 2, \dots$ and $j = 1, 2, \dots, k$.

From now on, if we say M is a matrix then we assume that its entries belong to $\{0, 1\}$ and the number of non-zero entries is finite. If $M = (\varepsilon_{ij})$ is a matrix then the monomial $x = x_1^{a_1}x_2^{a_2}\dots x_k^{a_k}$ corresponding to M is determined by

$$a_j = \sum_{i\geqslant 1} 2^{i-1}\varepsilon_{ij},\ j = 1, 2, \dots, k.$$

Define two sequences associated with monomial x by

$$\begin{aligned}\tau(x) &= (\tau_1(x); \tau_2(x); \dots; \tau_i(x); \dots),\\ \sigma(x) &= (a_1; a_2; \dots; a_k),\end{aligned}$$

where

$$\tau_i(x) = \sum_{j=1}^{k} \varepsilon_{ij}(x),\ i = 1, 2, \dots.$$

We call $\tau(x)$ the τ-sequence and $\sigma(x)$ the σ-sequence associated with monomial x.

We identify a finite sequence $(\xi_1; \xi_2; \dots; \xi_m)$ with $(\xi_1; \xi_2; \dots; \xi_m; 0; \dots)$.

Definition 2.1. Let x, y be the monomials in P_k. We say that $x < y$ if and only if one of the following holds

1. $\tau(x) < \tau(y)$;
2. $\tau(x) = \tau(y)$ and $\sigma(x) < \sigma(y)$.

Here the order on the set of sequences of non-negative integers is the lexicographical one.

Definition 2.2. A monomial x is said to be inadmissible if there exists the monomials $y_1, y_2, \ldots, y_r$ such that

$$x = y_1 + y_2 + \ldots + y_r \bmod \mathcal{A}^+.P_k \text{ and } y_j < x,\ j = 1, 2, \ldots, r.$$

A monomial x is said to be admissible if it is not inadmissible.

Obviously, the set of all admissible monomials in P_k is a minimal set of $\mathcal{A}$-generators of P_k.

Definition 2.3. Let M be an $s \times k$-matrix and x the monomial corresponding to M. The matrix M is said to be strictly inadmissible if and only if there exists the monomials $y_1, y_2, \ldots, y_r$ such that

$$x = y_1 + y_2 + \ldots + y_r + \sum_{0<i<2^s} \gamma_i Sq^i(z_i),\ \text{and } y_j < x,\ j = 1, 2, \ldots, r,$$

where $\gamma_i \in \mathbb{F}_2$, and $z_i \in P_k$ with $0 < i < 2^s$.

For a sequence of non-negative integers $\tau = (\tau_1; \tau_2; \ldots; \tau_m)$, we set

$$\mathcal{L}_k(\tau) = \text{Span}\Big\{x = x_1^{a_1}x_2^{a_2}\ldots x_k^{a_k} \in P_k\ ; \deg x = \sum_{i=1}^{m} 2^{i-1}\tau_i,\ \tau(x) < \tau\Big\}.$$

If x is a monomial in P_k and y is a monomial in $\mathcal{L}_k(\tau(x))$ then $y < x$. The space $\mathcal{L}_k(\tau)$ is an Σ_k-submodule of P_k, where Σ_k denote the symmetric group on k letters.

Suppose M is an $s \times k$-matrix and x is the monomial corresponding to M. Observe that if there exists the monomials $y_1, y_2, \ldots, y_r$ such that

$$x = y_1 + y_2 + \ldots + y_r + \sum_{0<i<2^s} \gamma_i Sq^i(z_i) \mod \mathcal{L}_k(\tau(x)),$$

where $\gamma_i \in \mathbb{F}_2, z_i \in P_k$ with $0 < i < 2^s$ and $y_j < x,\ j = 1, 2, \ldots, r$, then the matrix M is strictly inadmissible.

Notation 2.4. Let x be a monomial and $\Delta = (\delta_{ij})$ be an $s \times k$-matrix. If there is some non-negative integer r such that $\delta_{ij} = \varepsilon_{i+r,j}(x)$ for $1 \leqslant i \leqslant s$ and $1 \leqslant j \leqslant k$ then we write $\Delta \triangleright x$.

The following theorem is one of our main tools.

Theorem 2.5 (Kameko [8])**.** *Let x be a monomial and $\Delta = (\delta_{ij})$ be an $s \times k$-matrix. If the matrix Δ is strictly inadmissible and $\Delta \triangleright x$ then x is inadmissible.*

We need the following.

Lemma 2.6. *Let M be an $2 \times k$-matrix and x the monomial corresponding to M. If either $\tau_1(x) = 0, \tau_2(x) > 0$ or $\tau_1(x) < k, \tau_2(x) = k$ then M is strictly inadmissible.*

Proof. If $\tau_1(x) = 0$ and $\tau_2(x) > 0$ then $x = Sq^1(y)$ for some monomial y in P_k. Hence, the lemma is true. If $0 < \tau_1(x) = m < k$ and $\tau_2(x) = k$ then there is a ring homomorphism $f : P_k \to P_k$ induced by a permutation of $\{x_1, x_2, \ldots, x_k\}$ such that $x = f(x_1^3 \ldots x_m^3 x_{m+1}^2 x_{m+2}^2 \ldots x_k^2)$. It is easy to see that f is an $\mathcal{A}$-homomorphism. It sends monomials to monomials and preserves the associated τ-sequences. Obviously, we have

$$x_1^3 \ldots x_m^3 x_{m+1}^2 \ldots x_k^2 = Sq^1(x_1^3 \ldots x_m^3 x_{m+1} x_{m+2}^2 \ldots x_k^2) + \sum_{i=1}^{m} x_1^3 \ldots x_i^4 \ldots x_m^3 x_{m+1} x_{m+2}^2 \ldots x_k^2.$$

Since

$$\tau(x_1^3 \ldots x_i^4 \ldots x_m^3 x_{m+1} x_{m+2}^2 \ldots x_k^2) = (m; k-2; 1) < (m; k) = \tau(x_1^3 \ldots x_m^3 x_{m+1}^2 \ldots x_k^2),$$

for $i = 1, 2, \ldots, m$, the lemma holds. □

From Theorem 2.5 and Lemma 2.6, we easily obtain

Proposition 2.7. *Let x be an admissible monomial in P_k. Then we have*

1. If there is an index i_0 such that $\tau_{i_0}(x) = 0$ then $\tau_i(x) = 0$ for all $i > i_0$.

2. If there is an index i_0 such that $\tau_{i_0}(x) < k$ then $\tau_i(x) < k$ for all $i > i_0$.

For latter use, we set

$$Q_k = \mathrm{Span}\{x = x_1^{a_1} x_2^{a_2} \ldots x_k^{a_k} \ ; \ a_1 a_2 \ldots a_k = 0\},$$
$$R_k = \mathrm{Span}\{x = x_1^{a_1} x_2^{a_2} \ldots x_k^{a_k} \ ; \ a_1 a_2 \ldots a_k > 0\}.$$

It is easy to see that Q_k and R_k are the $\mathcal{A}$-submodules of P_k. Furethermore, we have the following.

Proposition 2.8. *We have a direct summand decomposition of the $\mathbb{F}_2$-vector spaces*

$$\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k = (\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_k) \oplus (\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_k).$$

From this proposition, if we know the set of all admissible monomials of P_{k-1} then we can easily determine a basis of $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_k$. So, to determine $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k$, we need only to determine a basis of $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_k$.

Now, we recall a result in Kameko [8] on the squaring operation.

Definition 2.9. Define the homomorphisms $\phi, \overline{Sq^0_*} : P_k \to P_k$ by

$$\phi(x) = x_1 x_2 \ldots x_k x^2,$$

$$\overline{Sq^0_*}(x) = \begin{cases} y, & \text{if } x = \phi(y), \\ 0, & \text{otherwise,} \end{cases}$$

for any monomial x in P_k.

Note that $\overline{Sq^0_*}$ commutes with the doubling map on $\mathcal{A}$. That is $Sq^t\overline{Sq^0_*} = \overline{Sq^0_*}Sq^{2t}$. Hence, $\overline{Sq^0_*}$ induces a homomorphism of $\mathbb{F}_2$-vector spaces

$$Sq^0_* : (\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k)_n \longrightarrow (\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k)_{\frac{n-k}{2}},$$

for all $n \geqslant k$ such that $n-k$ is even.

In general, $Sq^{2t}\phi \neq \phi Sq^t$. However, in one particular situation, we have the following.

Theorem 2.10 (Kameko [8]). *If $\beta(n) = k$ then Sq^0_* is an isomorphism of vector spaces and ϕ induces an inverse of it.*

Next, we recall some results of Wood [16] and Singer [14] on the hit monomials in P_k.

Theorem 2.11 (Wood [16]). *Let x be a monomial of degree n. If $\beta(n) > \tau_1(x)$ then x is hit.*

Definition 2.12. A monomial $z = x_1^{b_1}x_2^{b_2}\ldots x_k^{b_k}$ is called the spike if $b_j = 2^{s_j} - 1$ for s_j a non-negative integer and $j = 1, 2, \ldots, k$. If z is the spike with $s_1 > s_2 > \ldots > s_{r-1} \geqslant s_r > 0$ and $s_j = 0$ for $j > r$ then it is called the minimal spike.

The following is a main tool in our computation for P_4.

Theorem 2.13 (Singer [14]). *Suppose $x \in P_k$ is a monomial of degree n, where $\alpha(n+k) \leqslant k$. Let z be the minimal spike of degree n. If $\tau(x) < \tau(z)$ then x is hit.*

From this theorem, we see that if z is a minimal spike then $\mathcal{L}_k(\tau(z)) \subset \mathcal{A}^+.P_k$.

We end this section by defining some homomorphisms of $\mathcal{A}$-modules from P_4 to P_3 and some endomorphisms of P_4 which will be used in the next sections.

Let V_k denote the $\mathbb{F}_2$-vector subspace of P_k generated by $x_1, x_2, \ldots, x_k$. If $\widetilde{\varphi} : V_k \to V_{k'}$ is a homomorphism of $\mathbb{F}_2$-vector spaces then there exists uniquely ring homomorphism $\overline{\varphi} : P_k \to P_{k'}$ such that $\overline{\varphi}(x_i) = \widetilde{\varphi}(x_i)$ for $i = 1, 2, \ldots, k$. This homomorphism is also a homomorphism of $\mathcal{A}$-modules. So, it induces a homomorphism of $\mathbb{F}_2$-vector spaces $\varphi : \mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k \to \mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_{k'}$.

Note that if $\overline{\varphi} : P_k \to P_k$ is induced by a permutation of $\{x_1, x_2, \ldots, x_k\}$ then it sends monomials to monomials and preserves the associated τ-sequences.

In particular, consider the $\mathcal{A}$-homomorphisms $\overline{f}_i, \overline{g}_j, 1 \leqslant i \leqslant 6, 1 \leqslant j \leqslant 4$ and $\overline{h}$ from P_4 to P_3 defined by the following table

x	x_1	x_2	x_3	x_4
$\overline{f}_1(x)$	x_1	x_1	x_2	x_3
$\overline{f}_2(x)$	x_1	x_2	x_1	x_3
$\overline{f}_3(x)$	x_1	x_2	x_3	x_1
$\overline{f}_4(x)$	x_1	x_2	x_2	x_3
$\overline{f}_5(x)$	x_1	x_2	x_3	x_2
$\overline{f}_6(x)$	x_1	x_2	x_3	x_3
$\overline{g}_1(x)$	x_1+x_2	x_1	x_2	x_3
$\overline{g}_2(x)$	x_1+x_2	x_1	x_3	x_2
$\overline{g}_3(x)$	x_1+x_2	x_3	x_1	x_2
$\overline{g}_4(x)$	x_3	x_1+x_2	x_1	x_2
$\overline{h}(x)$	$x_1+x_2+x_3$	x_1	x_2	x_3.

These homomorphisms induce the homomorphisms f_i, g_j and h of $\mathbb{F}_2$-vector spaces from $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4$ to $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3$.

We use the above homomorphisms and the results in Kameko [8] to prove a certain subset of $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4$ is linearly independent. More precisely, to prove a subset of $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4$ is linearly independent, we consider a linear relation of the elements in this subset with coefficients in $\mathbb{F}_2$. By using Theorem 2.13, we compute the images of this linear relation under the action of the homomorphisms f_i, g_j, h. From the relations in $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3$ and the results in [8], we can show that all coefficients in the linear relation are zero.

Next, consider the endomorphisms $\overline{\varphi}_i, i = 1, 2, 3, 4$, of P_4 defined by the following table

x	x_1	x_2	x_3	x_4
$\overline{\varphi}_1(x)$	x_2	x_1	x_3	x_4
$\overline{\varphi}_2(x)$	x_1	x_3	x_2	x_4
$\overline{\varphi}_3(x)$	x_1	x_2	x_4	x_3
$\overline{\varphi}_4(x)$	x_1+x_2	x_2	x_3	x_4.

These endomorphisms induce the isomorphisms $\varphi_i, i = 1, 2, 3, 4$, from $\mathbb{F}_2$-vector space $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4$ to itself.

Note that the general linear group $GL_4(\mathbb{F}_2)$ is generated by $\overline{\varphi}_i, i = 1, 2, 3, 4$.

For simplicity, from now on, we denote the monomial $x = x_1^{a_1}x_2^{a_2}\dots x_k^{a_k}$ in P_k by $(a_1, a_2, \dots, a_k)$ and denote by $[x] = [a_1, a_2, \dots, a_k]$ the class in $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_k$ represented by the monomial $(a_1, a_2, \dots, a_k)$.

Suppose I is a finite subset of non-negative integers and $\gamma_i \in \mathbb{F}_2$ for $i \in I$. Denote by

$$\gamma_I = \sum_{i\in I} \gamma_i.$$

3. The indecomposables of P_4 in degree $2^{s+1}-3$

The purpose of this section is to prove Theorem 1.1 for the case $n = 2^{s+1}-3$.

According to Kameko [8], $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+1}-3} = 3$ with a basis given by the classes

$$v_{s,1} = [2^{s-1}-1, 2^{s-1}-1, 2^s-1], \quad v_{s,2} = [2^{s-1}-1, 2^s-1, 2^{s-1}-1],$$
$$v_{s,3} = [2^s-1, 2^{s-1}-1, 2^{s-1}-1].$$

So, we easily obtain

Proposition 3.1.

1. $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_1$ *is an* $\mathbb{F}_2$*-vector space of dimension 4 with a basis consisting of all the classes represented by the following monomials:*

$$a_{1,1} = (0,0,0,1),\ a_{1,2} = (0,0,1,0),\ a_{1,3} = (0,1,0,0),\ a_{1,4} = (1,0,0,0).$$

2. For $s \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+1}-3}$ *is an* $\mathbb{F}_2$*-vector space of dimension 12 with a basis consisting of all the classes represented by the following monomials:*

$$a_{s,1} = (0, 2^{s-1}-1, 2^{s-1}-1, 2^s-1), \quad a_{s,2} = (0, 2^{s-1}-1, 2^s-1, 2^{s-1}-1),$$
$$a_{s,3} = (0, 2^s-1, 2^{s-1}-1, 2^{s-1}-1), \quad a_{s,4} = (2^{s-1}-1, 0, 2^{s-1}-1, 2^s-1),$$
$$a_{s,5} = (2^{s-1}-1, 0, 2^s-1, 2^{s-1}-1), \quad a_{s,6} = (2^{s-1}-1, 2^{s-1}-1, 0, 2^s-1),$$
$$a_{s,7} = (2^{s-1}-1, 2^{s-1}-1, 2^s-1, 0), \quad a_{s,8} = (2^{s-1}-1, 2^s-1, 0, 2^{s-1}-1),$$
$$a_{s,9} = (2^{s-1}-1, 2^s-1, 2^{s-1}-1, 0), \quad a_{s,10} = (2^s-1, 0, 2^{s-1}-1, 2^{s-1}-1),$$
$$a_{s,11} = (2^s-1, 2^{s-1}-1, 0, 2^{s-1}-1), \quad a_{s,12} = (2^s-1, 2^{s-1}-1, 2^{s-1}-1, 0).$$

By Proposition 2.8, we need only to determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-3}$.

Set $\mu_1(1) = 4, \mu_1(2) = 15, \mu_1(3) = 35$ and $\mu_1(s) = 45$ for $s \geqslant 4$. We have

Theorem 3.2. *For any integer* $s \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-3}$ *is an* $\mathbb{F}_2$*-vector space of dimension* $\mu_1(s) - 12$ *with a basis consisting of all the classes represented by the following monomials:*

For $s = 2$,

$$a_{2,13} = (1,1,1,2), \quad a_{2,14} = (1,1,2,1), \quad a_{2,15} = (1,2,1,1).$$

For $s \geqslant 3$,

$$a_{s,13} = (1, 2^{s-1}-2, 2^{s-1}-1, 2^s-1), \quad a_{s,14} = (1, 2^{s-1}-2, 2^s-1, 2^{s-1}-1),$$
$$a_{s,15} = (1, 2^{s-1}-1, 2^{s-1}-2, 2^s-1), \quad a_{s,16} = (1, 2^{s-1}-1, 2^s-1, 2^{s-1}-2),$$
$$a_{s,17} = (1, 2^s-1, 2^{s-1}-2, 2^{s-1}-1), \quad a_{s,18} = (1, 2^s-1, 2^{s-1}-1, 2^{s-1}-2),$$
$$a_{s,19} = (2^{s-1}-1, 1, 2^{s-1}-2, 2^s-1), \quad a_{s,20} = (2^{s-1}-1, 1, 2^s-1, 2^{s-1}-2),$$
$$a_{s,21} = (2^{s-1}-1, 2^s-1, 1, 2^{s-1}-2), \quad a_{s,22} = (2^s-1, 1, 2^{s-1}-2, 2^{s-1}-1),$$
$$a_{s,23} = (2^s-1, 1, 2^{s-1}-1, 2^{s-1}-2), \quad a_{s,24} = (2^s-1, 2^{s-1}-1, 1, 2^{s-1}-2),$$
$$a_{s,25} = (1, 2^{s-1}-1, 2^{s-1}-1, 2^s-2), \quad a_{s,26} = (1, 2^{s-1}-1, 2^s-2, 2^{s-1}-1),$$
$$a_{s,27} = (1, 2^s-2, 2^{s-1}-1, 2^{s-1}-1), \quad a_{s,28} = (2^{s-1}-1, 1, 2^{s-1}-1, 2^s-2),$$
$$a_{s,29} = (2^{s-1}-1, 1, 2^s-2, 2^{s-1}-1), \quad a_{s,30} = (2^{s-1}-1, 2^{s-1}-1, 1, 2^s-2).$$

For $s = 3$,

$$a_{3,31} = (3,3,5,2), \quad a_{3,32} = (3,5,2,3), \quad a_{3,33} = (3,5,3,2),$$
$$a_{3,34} = (3,3,3,4), \quad a_{3,35} = (3,3,4,3).$$

For $s \geqslant 4$,

$a_{s,31} = (3, 2^{s-1}-3, 2^{s-1}-2, 2^s-1)$, $a_{s,32} = (3, 2^{s-1}-3, 2^s-1, 2^{s-1}-2)$,
$a_{s,33} = (3, 2^s-1, 2^{s-1}-3, 2^{s-1}-2)$, $a_{s,34} = (2^s-1, 3, 2^{s-1}-3, 2^{s-1}-2)$,
$a_{s,35} = (3, 2^{s-1}-3, 2^{s-1}-1, 2^s-2)$, $a_{s,36} = (3, 2^{s-1}-3, 2^s-2, 2^{s-1}-1)$,
$a_{s,37} = (3, 2^{s-1}-1, 2^{s-1}-3, 2^s-2)$, $a_{s,38} = (2^{s-1}-1, 3, 2^{s-1}-3, 2^s-2)$,
$a_{s,39} = (3, 2^{s-1}-1, 2^s-3, 2^{s-1}-2)$, $a_{s,40} = (3, 2^s-3, 2^{s-1}-2, 2^{s-1}-1)$,
$a_{s,41} = (3, 2^s-3, 2^{s-1}-1, 2^{s-1}-2)$, $a_{s,42} = (2^{s-1}-1, 3, 2^s-3, 2^{s-1}-2)$,
$a_{s,43} = (7, 2^s-5, 2^{s-1}-3, 2^{s-1}-2)$.

For $s = 4$,

$$a_{4,44} = (7,7,9,6), \quad a_{4,45} = (7,7,7,8).$$

For $s \geqslant 5$,

$$a_{s,44} = (7, 2^{s-1}-5, 2^{s-1}-3, 2^s-2), \quad a_{s,45} = (7, 2^{s-1}-5, 2^s-3, 2^{s-1}-2).$$

We prove this theorem by proving the following propositions.

Proposition 3.3. *For $s \geqslant 2$, the $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-3}$ is generated by the $\mu_1(s) - 12$ elements listed in Theorem 3.2.*

The proposition is proved by combining Theorem 2.5 and the following lemmas.

Lemma 3.4. *If x is an admissible monomial of degree $2^{s+1}-3$ in P_4 then*

$$\tau(x) = (\underbrace{3;3;\ldots;3}_{s-1 \text{ times}};1).$$

Proof. It is easy to see that the lemma holds for $s = 1$. Suppose $s \geqslant 2$. Obviously, $z = (2^s-1, 2^{s-1}-1, 2^{s-1}-1, 0)$ is the minimal spike of degree $2^{s+1}-3$ in P_4 and $\tau(z) = (\underbrace{3;3;\ldots;3}_{s-1 \text{ times}};1)$. Since $2^{s+1}-3$ is odd, we get either $\tau_1(x) = 1$ or $\tau_1(x) = 3$. If $\tau_1(x) = 1$ then $\tau(x) < \tau(z)$. By Theorem 2.13, x is hit. This contradicts the fact that x is admissible. Hence, we have $\tau_1(x) = 3$. Using Proposition 2.7 and Theorem 2.13, we obtain $\tau_i(x) = 3$, $i = 1, 2, \ldots, s-1$. From this, it implies

$$2^{s+1}-3 = \deg x = \sum_{i \geqslant 1} 2^{i-1}\tau_i(x) = 3(2^{s-1}-1) + \sum_{i \geqslant s} 2^{i-1}\tau_i(x).$$

The last equality implies $\tau_s(x) = 1$ and $\tau_i(x) = 0$ for $i > s$. The lemma is proved. □

By this lemma, it suffices to consider monomials whose associated τ-sequences are $(\underbrace{3;3;\ldots;3}_{s-1 \text{ times}};1)$.

Lemma 3.5. *The following matrices are strictly inadmissible*

$$\Delta_1 = \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 1 & 0 \end{pmatrix},\ \Delta_2 = \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 1 & 1 & 0 \end{pmatrix},\ \Delta_3 = \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 1 & 0 & 1 \end{pmatrix},$$

$$\Delta_4 = \begin{pmatrix} 0 & 1 & 1 & 1\\ 1 & 1 & 1 & 0 \end{pmatrix},\ \Delta_5 = \begin{pmatrix} 0 & 1 & 1 & 1\\ 1 & 1 & 0 & 1 \end{pmatrix},\ \Delta_6 = \begin{pmatrix} 0 & 1 & 1 & 1\\ 1 & 0 & 1 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to six matrices in the lemma respectively are

$$(3,3,2,1), (3,2,3,1), (3,2,1,3), (2,3,3,1), (2,3,1,3), (2,1,3,3).$$

If x is one of these monomials then $\tau(x) = (3;3)$ and there is an $\mathcal{A}$-homomorphism $f : P_4 \to P_4$ induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = f(3,3,2,1)$. Furethermore, f sends monomials to monomials and preserves the associated τ-sequences. By a direct computation, we have

$$(3,3,2,1) = Sq^1(3,3,1,1) + (4,3,1,1) + (3,4,1,1) + (3,3,1,2).$$

It is easy to see that

$$\begin{aligned} &\tau(4,3,1,1) = \tau(3,4,1,1) = (3;1;1) < (3;3),\\ &\tau(3,3,1,2) = \tau(3,3,2,1) \text{ and } \sigma(3,3,1,2) < \sigma(3,3,2,1). \end{aligned}$$

Hence, the lemma is proved. □

Lemma 3.6. *The following matrices are strictly inadmissible*

$$\Delta_7 = \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 1 \end{pmatrix},\ \Delta_8 = \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 1 \end{pmatrix},\ \Delta_9 = \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 0 & 1 & 1 \end{pmatrix},$$

$$\Delta_{10} = \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 0 & 1 & 1 & 1 \end{pmatrix},\ \Delta_{11} = \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 0 & 1 & 1 & 1 \end{pmatrix},\ \Delta_{12} = \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 1\\ 0 & 1 & 1 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to six matrices in this lemma respectively are

$$(7,7,3,4), (7,3,7,4), (7,3,4,7), (3,7,7,4), (3,7,4,7), (3,4,7,7).$$

If x is one of these monomials then $\tau(x) = (3;3;3)$ and there is an $\mathcal{A}$-homomorphism $g : P_4 \to P_4$ induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = g(7,7,3,4)$. By a direct computation, we get

$$\begin{aligned} (7,7,3,4) = Sq^1\big((7,7,3,3) + (8,7,2,3)\big) + Sq^2(7,7,2,3)\\ + (8,7,3,3) + (7,8,3,3) + (9,7,2,3)\\ + (7,9,2,3) + (7,8,2,4) + (7,7,2,5). \end{aligned}$$

So, the lemma is proved. □

Lemma 3.7. *The following matrices are strictly inadmissible*

$$\Delta_{13} = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad \Delta_{14} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix},$$

$$\Delta_{15} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad \Delta_{16} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices are

$$(2,1,1,1), (3,4,3,3), (7,7,8,7), (15,15,15,16).$$

By a direct computation, we have

$$(2,1,1,1) = Sq^1(1,1,1,1) + (1,2,1,1) + (1,1,2,1) + (1,1,1,2).$$

Since $\tau(1,2,1,1) = \tau(1,1,2,1) = \tau(1,1,1,2) = \tau(2,1,1,1) = (3;1)$ and $\sigma(1,2,1,1), \sigma(1,1,2,1), \sigma(1,1,1,2) < \sigma(2,1,1,1)$, the lemma is true.

$$\begin{aligned}(3,4,3,3) = Sq^1\big((3,3,3,3) + (2,4,3,3)\big) + Sq^2(2,3,3,3)\\ + (3,3,4,3) + (3,3,3,4) + (2,5,3,3)\\ + (2,3,5,3) + (2,3,3,5) + (2,3,4,4).\end{aligned}$$

Let y be a monomial in $(P_4)_{13}$. If y is a term in the right hand side of the above equality and $y \neq (2,3,4,4)$ then $\tau(y) = \tau(3,4,3,3) = (3;3;1)$ and $\sigma(y) < \sigma(3,4,3,3)$. If $y = (2,3,4,4)$ then $\tau(y) = (1;2;2) < (3;3;1)$. Hence, the lemma holds.

$$\begin{aligned}(7,7,8,7) = Sq^1\big((7,7,7,7) + (6,8,7,7) + (8,6,7,7) + (6,6,9,7)\\ + (6,6,7,9)\big) + Sq^2\big((6,7,7,7) + (7,6,7,7) + (5,8,7,7)\big)\\ + Sq^4(5,6,7,7) + (7,7,7,8) + (6,9,7,7) + (6,7,9,7) + (6,7,7,9)\\ + (7,6,9,7) + (7,6,7,9) + (5,10,7,7) + (5,6,11,7) + (5,6,7,11)\\ + (6,7,8,8) + (7,6,8,8) + (5,6,9,9) + (5,6,10,8) + (5,6,8,10).\end{aligned}$$

Using the above equality and by an argument analogous to the previous one, we see that the lemma is true.

By a direct computation, we get

$$\begin{aligned}(15,15,15,16) = Sq^1(15,15,15,15) + Sq^2\big((14,15,15,15) + (15,14,15,15)\\ + (15,15,14,15)\big) + Sq^4\big((13,14,15,15) + (15,13,14,15)\big)\\ + Sq^8(11,13,14,15) + (14,17,15,15) + (14,15,17,15) + (14,15,15,17)\\ + (15,14,17,15) + (15,14,15,17) + (15,15,14,17) + (13,18,15,15)\\ + (13,14,19,15) + (13,14,15,19) + (15,13,18,15) + (15,13,14,19)\\ + (11,21,14,15) + (11,13,22,15) + (11,13,14,23) \bmod \mathcal{L}_4(3;3;3;3;1).\end{aligned}$$

Let y be a monomial in $(P_4)_{61}$. If y is a term of the right hand side of the above equality then $\tau(y) = \tau(15,15,15,16) = (3;3;3;3;1)$ and $\sigma(y) < \sigma(15,15,15,16)$. If $y \in \mathcal{L}_4(3;3;3;3;1)$ then $\tau(y) < (3;3;3;3;1)$. So $y < (15,15,15,16)$ and the lemma holds. □

Now we prove Proposition 3.3 by using Theorem 2.5, Lemmas 3.4, 3.5, 3.6, 3.7.

Proof of Proposition 3.3. Let x be a monomial of degree $2^{s+1}-3$ in P_4 with $\tau_i(x) = 3, i = 1,2,\ldots,s-1, \tau_s(x) = 1, \tau_i(x) = 0, i > s$. Then we can write $x = z_i y^2$ for $1 \leqslant i \leqslant 4$, where

$$z_1 = (0,1,1,1),\ z_2 = (1,0,1,1),\ z_3 = (1,1,0,1),\ z_4 = (1,1,1,0),$$

and y is a monomial of degree $2^s - 3$ with $\tau_i(y) = 3, i = 1,2,\ldots,s-2, \tau_{s-1}(y) = 1, \tau_i(y) = 0, i \geqslant s$.

We prove the proposition by showing that if $x \neq a_{s,j}, 1 \leqslant j \leqslant \mu_1(s)$ then there is a strictly inadmissible matrix Δ such that $\Delta \triangleright x$.

The proof proceeds by induction on s. The case $s = 2$ is trivial. By the inductive hypothesis and Theorem 2.5, it suffices to consider monomials $x = z_i y^2$ with $y = a_{s,j}, 1 \leqslant j \leqslant \mu_1(s)$.

For $s = 3$, a direct computation shows

$$\begin{array}{llll}
a_{3,1} = z_1 a_{2,1}^2, & a_{3,2} = z_1 a_{2,2}^2, & a_{3,3} = z_1 a_{2,3}^2, & a_{3,4} = z_2 a_{2,4}^2,\\
a_{3,5} = z_2 a_{2,5}^2, & a_{3,6} = z_3 a_{2,6}^2, & a_{3,7} = z_4 a_{2,7}^2, & a_{3,8} = z_3 a_{2,8}^2,\\
a_{3,9} = z_4 a_{2,9}^2, & a_{3,10} = z_2 a_{2,10}^2, & a_{3,11} = z_3 a_{2,11}^2, & a_{3,12} = z_4 a_{2,12}^2,\\
a_{3,13} = z_2 a_{2,1}^2, & a_{3,14} = z_2 a_{2,2}^2, & a_{3,15} = z_3 a_{2,1}^2, & a_{3,16} = z_4 a_{2,2}^2,\\
a_{3,17} = z_3 a_{2,3}^2, & a_{3,18} = z_4 a_{2,3}^2, & a_{3,19} = z_3 a_{2,4}^2, & a_{3,20} = z_4 a_{2,5}^2,\\
a_{3,21} = z_4 a_{2,8}^2, & a_{3,22} = z_3 a_{2,10}^2, & a_{3,23} = z_4 a_{2,10}^2, & a_{3,24} = z_4 a_{2,11}^2,\\
a_{3,25} = z_4 a_{2,1}^2, & a_{3,26} = z_3 a_{2,2}^2, & a_{3,27} = z_2 a_{2,3}^2, & a_{3,28} = z_4 a_{2,4}^2,\\
a_{3,29} = z_3 a_{2,5}^2, & a_{3,30} = z_4 a_{2,6}^2, & a_{3,31} = z_4 a_{2,14}^2, & a_{3,32} = z_3 a_{2,15}^2,\\
a_{3,33} = z_4 a_{2,15}^2, & a_{3,34} = z_4 a_{2,13}^2, & a_{3,35} = z_3 a_{2,14}^2. &
\end{array}$$

We have $\Delta_1 \triangleright z_3 a_{2,j}^2$, $\Delta_2 \triangleright z_2 a_{2,j}^2$, $\Delta_4 \triangleright z_1 a_{2,j}^2$ for $j = 7, 9, 12, 13$; $\Delta_3 \triangleright z_2 a_{2,j}^2$, $\Delta_5 \triangleright z_1 a_{2,j}^2$ for $j = 6, 8, 11, 14$; $\Delta_6 \triangleright z_1 a_{2,j}^2$ for $j = 4, 5, 10, 15$; $\Delta_{14} \triangleright z_2 a_{2,15}^2$.

From the above equalities, Lemmas 3.5, 3.6, 3.7 and Theorem 2.5, we see that if $x \neq a_{3,j}, 1 \leqslant j \leqslant 35$ then x is inadmissible. So, the case $s = 3$ is true.

Let $s = 4$. By a direct computation we obtain

$$\begin{array}{llll}
a_{4,1} = z_1 a_{3,1}^2, & a_{4,2} = z_1 a_{3,2}^2, & a_{4,3} = z_1 a_{3,3}^2, & a_{4,4} = z_2 a_{3,4}^2,\\
a_{4,5} = z_2 a_{3,5}^2, & a_{4,6} = z_3 a_{3,6}^2, & a_{4,7} = z_4 a_{3,7}^2, & a_{4,8} = z_3 a_{3,8}^2,\\
a_{4,9} = z_4 a_{3,9}^2, & a_{4,10} = z_2 a_{3,10}^2, & a_{4,11} = z_3 a_{3,11}^2, & a_{4,12} = z_4 a_{3,12}^2,\\
a_{4,13} = z_2 a_{3,1}^2, & a_{4,14} = z_2 a_{3,2}^2, & a_{4,15} = z_3 a_{3,1}^2, & a_{4,16} = z_4 a_{3,2}^2,\\
a_{4,17} = z_3 a_{3,3}^2, & a_{4,18} = z_4 a_{3,3}^2, & a_{4,19} = z_3 a_{3,4}^2, & a_{4,20} = z_4 a_{3,5}^2,\\
a_{4,21} = z_4 a_{3,8}^2, & a_{4,22} = z_3 a_{3,10}^2, & a_{4,23} = z_4 a_{3,10}^2, & a_{4,24} = z_4 a_{3,11}^2,\\
a_{4,25} = z_4 a_{3,1}^2, & a_{4,26} = z_3 a_{3,2}^2, & a_{4,27} = z_2 a_{3,3}^2, & a_{4,28} = z_4 a_{3,4}^2,\\
a_{4,29} = z_3 a_{3,5}^2, & a_{4,30} = z_4 a_{3,6}^2, & a_{4,31} = z_3 a_{3,13}^2, & a_{4,32} = z_4 a_{3,14}^2,\\
a_{4,33} = z_4 a_{3,17}^2, & a_{4,34} = z_4 a_{3,22}^2, & a_{4,35} = z_4 a_{3,13}^2, & a_{4,36} = z_3 a_{3,14}^2,
\end{array}$$

$a_{4,37} = z_4a_{3,15}^2$, $a_{4,38} = z_4a_{3,19}^2$, $a_{4,39} = z_4a_{3,26}^2$, $a_{4,40} = z_3a_{3,27}^2$, $a_{4,41} = z_4a_{3,27}^2$, $a_{4,42} = z_4a_{3,29}^2$, $a_{4,43} = z_4a_{3,32}^2$, $a_{4,44} = z_4a_{3,35}^2$, $a_{4,45} = z_4a_{3,34}^2$.

We have also $\Delta_1 \triangleright z_3a_{3,j}^2$, $\Delta_2 \triangleright z_2a_{3,j}^2$, $\Delta_4 \triangleright z_1a_{3,j}^2$ for $j = 7, 9, 12, 16, 18, 20, 21, 23, 24, 25, 28, 30, 31, 33, 34$; $\Delta_3 \triangleright z_2a_{3,j}^2$, $\Delta_5 \triangleright z_1a_{3,j}^2$ for $j = 6, 8, 11, 15, 17, 19, 22, 26, 29, 32, 35$; $\Delta_6 \triangleright z_1a_{3,j}^2$ for $j = 4, 5, 10, 13, 14, 27$; $\Delta_7 \triangleright z_4a_{3,j}^2$ for $j = 21, 24, 30, 31$; $\Delta_8 \triangleright z_4a_{3,j}^2$ for $j = 20, 23, 28, 33$; $\Delta_9 \triangleright z_3a_{3,j}^2$ for $j = 19, 22, 29, 32$; $\Delta_{10} \triangleright z_4a_{3,j}^2$ for $j = 16, 18, 25$; $\Delta_{11} \triangleright z_3a_{3,j}^2$ for $j = 15, 17, 26$; $\Delta_{12} \triangleright z_2a_{3,j}^2$ for $j = 13, 14, 27$; $\Delta_{15} \triangleright z_3a_{3,25}^2$. These equalities, Lemmas 3.5, 3.6, 3.7 and Theorem 2.5 imply that if $x \neq a_{4,j}, 1 \leqslant j \leqslant 45$ then x is inadmissible. Hence, the case $s = 4$ is true.

Suppose $s \geqslant 4$ and the proposition holds for s. By a direct computation, we have $\Delta_1 \triangleright z_3a_{s,j}^2$, $\Delta_2 \triangleright z_2a_{s,j}^2$, $\Delta_4 \triangleright z_1a_{s,j}^2$ for $j = 7, 9, 12, 16, 18, 20, 21, 23, 24, 25, 28, 30, 32, 33, 34, 35, 37, 38, 39, 41, 42, 43, 44, 45$; $\Delta_3 \triangleright z_2a_{s,j}^2$, $\Delta_5 \triangleright z_1a_{s,j}^2$ for $j = 6, 8, 11, 15, 17, 19, 22, 26, 29, 31, 36, 40$; $\Delta_6 \triangleright z_1a_{s,j}^2$ for $j = 4, 5, 10, 13, 14, 27$; $\Delta_7 \triangleright z_4a_{s,j}^2$ for $j = 21, 24, 30, 33, 34, 37, 38, 39, 42, 43, 44$; $\Delta_{16} \triangleright z_4a_{s,45}^2$ for $s = 4$; $\Delta_7 \triangleright z_4a_{s,45}^2$ for $s \geqslant 5$; $\Delta_8 \triangleright z_4a_{s,j}^2$ for $j = 20, 23, 28, 32, 35, 41$; $\Delta_9 \triangleright z_3a_{s,j}^2$ for $j = 19, 22, 29, 31, 36, 40$; $\Delta_{10} \triangleright z_4a_{s,j}^2$ for $j = 16, 18, 25$; $\Delta_{11} \triangleright z_3a_{s,j}^2$ for $j = 15, 17, 26$; $\Delta_{12} \triangleright z_2a_{s,j}^2$ for $j = 13, 14, 27$ and

$a_{s+1,1} = z_1a_{s,1}^2$, $a_{s+1,2} = z_1a_{s,2}^2$, $a_{s+1,3} = z_1a_{s,3}^2$, $a_{s+1,4} = z_2a_{s,4}^2$, $a_{s+1,5} = z_2a_{s,5}^2$, $a_{s+1,6} = z_3a_{s,6}^2$, $a_{s+1,7} = z_4a_{s,7}^2$, $a_{s+1,8} = z_3a_{s,8}^2$, $a_{s+1,9} = z_4a_{s,9}^2$, $a_{s+1,10} = z_2a_{s,10}^2$, $a_{s+1,11} = z_3a_{s,11}^2$, $a_{s+1,12} = z_4a_{s,12}^2$, $a_{s+1,13} = z_2a_{s,1}^2$, $a_{s+1,14} = z_2a_{s,2}^2$, $a_{s+1,15} = z_3a_{s,1}^2$, $a_{s+1,16} = z_4a_{s,2}^2$, $a_{s+1,17} = z_3a_{s,3}^2$, $a_{s+1,18} = z_4a_{s,3}^2$, $a_{s+1,19} = z_3a_{s,4}^2$, $a_{s+1,20} = z_4a_{s,5}^2$, $a_{s+1,21} = z_4a_{s,8}^2$, $a_{s+1,22} = z_3a_{s,10}^2$, $a_{s+1,23} = z_4a_{s,10}^2$, $a_{s+1,24} = z_4a_{s,11}^2$, $a_{s+1,25} = z_4a_{s,1}^2$, $a_{s+1,26} = z_3a_{s,2}^2$, $a_{s+1,27} = z_2a_{s,3}^2$, $a_{s+1,28} = z_4a_{s,4}^2$, $a_{s+1,29} = z_3a_{s,5}^2$, $a_{s+1,30} = z_4a_{s,6}^2$, $a_{s+1,31} = z_3a_{s,13}^2$, $a_{s+1,32} = z_4a_{s,14}^2$, $a_{s+1,33} = z_4a_{s,17}^2$, $a_{s+1,34} = z_4a_{s,22}^2$, $a_{s+1,35} = z_4a_{s,13}^2$, $a_{s+1,36} = z_3a_{s,14}^2$, $a_{s+1,37} = z_4a_{s,15}^2$, $a_{s+1,38} = z_4a_{s,19}^2$, $a_{s+1,39} = z_4a_{s,26}^2$, $a_{s+1,40} = z_3a_{s,27}^2$, $a_{s+1,41} = z_4a_{s,27}^2$, $a_{s+1,42} = z_4a_{s,29}^2$, $a_{s+1,43} = z_4a_{s,40}^2$, $a_{s+1,44} = z_4a_{s,36}^2$, $a_{s+1,45} = z_4a_{s,31}^2$.

Using these equalities, Lemmas 3.5, 3.6, 3.7 and Theorem 2.5, we obtain the proposition for $s + 1$. The proof is completed. □

Now, we show that $[a_{s,j}]; 13 \leqslant j \leqslant \mu_1(s)$, are linearly independent.

Proposition 3.8. *The elements* $[a_{2,13}], [a_{2,14}], [a_{2,15}]$ *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_5$.

Proof. Suppose there is a linear relation $\gamma_1[a_{2,13}] + \gamma_2[a_{2,14}] + \gamma_3[a_{2,15}] = 0$, where $\gamma_1, \gamma_2, \gamma_3 \in \mathbb{F}_2$. Consider the homomorphisms f_1, f_2, f_3. Under these homomorphisms, the images of the above linear relation respectively are

$$\gamma_3[3, 1, 1] = 0,\ \gamma_2[3, 1, 1] = 0,\ \gamma_1[3, 1, 1] = 0.$$

Hence, $\gamma_1 = \gamma_2 = \gamma_3 = 0$. The proposition is proved. □

Proposition 3.9. *The elements* $[a_{3,j}], 13 \leqslant j \leqslant 35$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{13}$.

Proof. Suppose there is a linear relation

$$\sum_{j=13}^{35} \gamma_j [a_{3,j}] = 0, \tag{3.9.1}$$

where $\gamma_j \in \mathbb{F}_2, 13 \leqslant j \leqslant 35$. By a direct calculation, we see that the homomorphisms f_i, for $i \leqslant i \leqslant 6$, send the relation (3.9.1) respectively to

$$\begin{aligned}
&\gamma_{13}[3,3,7] + \gamma_{14}[3,7,3] + \gamma_{27}[7,3,3] = 0,\\
&\gamma_{15}[3,3,7] + \gamma_{17}[3,7,3] + \gamma_{\{26,35\}}[7,3,3] = 0,\\
&\gamma_{16}[3,3,7] + \gamma_{18}[3,7,3] + \gamma_{\{25,34\}}[7,3,3] = 0,\\
&\gamma_{19}[3,3,7] + \gamma_{\{29,32,35\}}[3,7,3] + \gamma_{22}[7,3,3] = 0,\\
&\gamma_{20}[3,3,7] + \gamma_{\{28,33,34\}}[3,7,3] + \gamma_{23}[7,3,3] = 0,\\
&\gamma_{\{30,31,34,35\}}[3,3,7] + \gamma_{21}[3,7,3] + \gamma_{24}[7,3,3] = 0.
\end{aligned}$$

From the above equalities, we obtain

$$\begin{cases}
\gamma_j = 0,\ j = 13,14,15,16,17,18,19,20,21,22,23,24,27,\\
\gamma_{\{26,35\}} = \gamma_{\{25,34\}} = \gamma_{\{29,32,35\}} = \gamma_{\{28,33,34\}} = \gamma_{\{30,31,34,35\}} = 0.
\end{cases} \tag{3.9.2}$$

Substituting (3.9.2) into the relation (3.9.1), we have

$$\sum_{25 \leqslant j \leqslant 35, j \neq 27} \gamma_j [a_{3,j}] = 0. \tag{3.9.3}$$

The homomorphisms $g_i, i = 1,2,3$, send (3.9.3) respectively to

$$\begin{aligned}
&\gamma_{\{26,29,35\}}[3,7,3] + \gamma_{32}[7,3,3] = 0,\\
&\gamma_{\{25,28,34\}}[3,7,3] + \gamma_{33}[7,3,3] = 0,\\
&\gamma_{\{25,30,34\}}[3,7,3] + \gamma_{\{26,31,35\}}[7,3,3] = 0.
\end{aligned}$$

Hence, we get

$$\begin{cases}
\gamma_{32} = \gamma_{33} = 0,\\
\gamma_{\{26,29,35\}} = \gamma_{\{25,28,34\}} = 0,\\
\gamma_{\{25,30,34\}} = \gamma_{\{26,31,35\}} = 0.
\end{cases} \tag{3.9.4}$$

Combining (3.9.2) and (3.9.4), we obtain $\gamma_j = 0,\ j = 13,14,\ldots,35$. The proposition follows. □

Proposition 3.10. *The elements* $[a_{4,j}], 13 \leqslant j \leqslant 45$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{29}$.

Proof. Suppose there is a linear relation

$$\sum_{j=13}^{45} \gamma_j[a_{4,j}] = 0, \tag{3.10.1}$$

where $\gamma_j \in \mathbb{F}_2, 13 \leqslant j \leqslant 45$.

Applying the homomorphisms f_i, for $i = 1, 2, \ldots, 6$, to the relation (3.10.1), we get

$$\begin{aligned}
&\gamma_{13}[7,7,15] + \gamma_{14}[7,15,7] + \gamma_{27}[15,7,7] = 0,\\
&\gamma_{15}[7,7,15] + \gamma_{17}[7,15,7] + \gamma_{26}[15,7,7] = 0,\\
&\gamma_{16}[7,7,15] + \gamma_{18}[7,15,7] + \gamma_{\{25,45\}}[15,7,7] = 0,\\
&\gamma_{19}[7,7,15] + \gamma_{29}[7,15,7] + \gamma_{22}[15,7,7] = 0,\\
&\gamma_{20}[7,7,15] + \gamma_{\{28,45\}}[7,15,7] + \gamma_{23}[15,7,7] = 0,\\
&\gamma_{\{30,44,45\}}[7,7,15] + \gamma_{21}[7,15,7] + \gamma_{24}[15,7,7] = 0.
\end{aligned}$$

From these equalities, we obtain

$$\begin{cases}
\gamma_j = 0, j = 13, 14, \ldots, 24, 26, 27, 29\\
\gamma_{\{25,45\}} = \gamma_{\{28,45\}} = \gamma_{\{30,44,45\}} = 0.
\end{cases} \tag{3.10.2}$$

Substituting (3.10.2) into the relation (3.10.1), we get

$$\sum_{25\leqslant j\leqslant 45, j\neq 26,27,29} \gamma_j[a_{4,j}] = 0. \tag{3.10.3}$$

Apply the homomorphisms g_i, for $i = 1, 2, 3, 4$, to the relation (3.10.3), we obtain

$$\begin{aligned}
&\gamma_{31}[7,7,15] + \gamma_{36}[7,15,7] + \gamma_{40}[15,7,7] = 0,\\
&\gamma_{32}[7,7,15] + \gamma_{\{25,28,35,45\}}[7,15,7] + \gamma_{41}[15,7,7] = 0,\\
&\gamma_{33}[7,7,15] + \gamma_{\{25,30,37,45\}}[7,15,7] + \gamma_{\{39,44\}}[15,7,7] = 0,\\
&\gamma_{34}[7,7,15] + \gamma_{\{28,30,38,43,45\}}[7,15,7] + \gamma_{\{42,43,44\}}[15,7,7] = 0.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
\gamma_j = 0,\ j = 31, 32, 33, 34, 36, 40, 41,\\
\gamma_{\{25,28,35,45\}} = \gamma_{\{25,30,37,45\}} = 0,\\
\gamma_{\{39,44\}} = \gamma_{\{28,30,38,43,45\}} = \gamma_{\{42,43,44\}} = 0.
\end{cases} \tag{3.10.4}$$

Combining the relations (3.10.3) and (3.10.4) gives

$$\gamma_{25}[a_{4,25}] + \gamma_{28}[a_{4,28}] + \gamma_{30}[a_{4,30}] + \sum_{35\leqslant j\leqslant 45, j\neq 36,40,41} \gamma_j[a_{4,j}] = 0. \tag{3.10.5}$$

Under the homomorphism h, the image of the linear relation (3.10.5) is

$$\gamma_{\{25,28,30,35,37,38,45\}}[7,7,15] + \gamma_{\{39,42,44\}}[7,15,7] + \gamma_{43}[15,7,7] = 0.$$

From this, it implies

$$\gamma_{\{25,28,30,35,37,38,45\}} = \gamma_{\{39,42,44\}} = \gamma_{43} = 0. \tag{3.10.6}$$

Combining (3.10.2), (3.10.4) and (3.10.6), we obtain $\gamma_j = 0$ for $13 \leqslant j \leqslant 45$. The proposition follows. □

Proposition 3.11. *For any $s \geqslant 5$, the elements $[a_{s,j}], 13 \leqslant j \leqslant 45$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-3}$.*

Proof. Suppose there is a linear relation of the above elements

$$\sum_{j=13}^{45} \gamma_j [a_{s,j}] = 0, \tag{3.11.1}$$

where $\gamma_j \in \mathbb{F}_2, 13 \leqslant j \leqslant 45$.

Applying the homomorphisms $f_i, i = 1, 2, \ldots, 6$, to the relation (3.11.1), we get

$$\begin{aligned}
&\gamma_{13}v_{s,1} + \gamma_{14}v_{s,2} + \gamma_{27}v_{s,3} = 0, \quad \gamma_{15}v_{s,1} + \gamma_{17}v_{s,2} + \gamma_{26}v_{s,3} = 0,\\
&\gamma_{16}v_{s,1} + \gamma_{18}v_{s,2} + \gamma_{25}v_{s,3} = 0, \quad \gamma_{19}v_{s,1} + \gamma_{29}v_{s,2} + \gamma_{22}v_{s,3} = 0,\\
&\gamma_{20}v_{s,1} + \gamma_{28}v_{s,2} + \gamma_{23}v_{s,3} = 0, \quad \gamma_{30}v_{s,1} + \gamma_{21}v_{s,2} + \gamma_{24}v_{s,3} = 0.
\end{aligned}$$

From this, it implies $\gamma_j = 0,\ j = 13, 14, \ldots, 30$. So, the relation (3.11.1) becomes

$$\sum_{j=31}^{45} \gamma_j [a_{s,j}] = 0. \tag{3.11.2}$$

Applying the homomorphisms $g_i, i = 1, 2, 3, 4$, to (3.11.2), we obtain

$$\begin{aligned}
&\gamma_{31}v_{s,1} + \gamma_{36}v_{s,2} + \gamma_{40}v_{s,3} = 0, \quad \gamma_{32}v_{s,1} + \gamma_{35}v_{s,2} + \gamma_{41}v_{s,3} = 0,\\
&\gamma_{33}v_{s,1} + \gamma_{37}v_{s,2} + \gamma_{39}v_{s,3} = 0, \quad \gamma_{34}v_{s,1} + \gamma_{38}v_{s,2} + \gamma_{42}v_{s,3} = 0.
\end{aligned}$$

So, we obtain $\gamma_j = 0,\ j = 31, 32, \ldots, 42$. Hence, the relation (3.11.2) becomes

$$\gamma_{43}[a_{s,43}] + \gamma_{44}[a_{s,44}] + \gamma_{45}[a_{s,45}] = 0. \tag{3.11.3}$$

Now, apply the homomorphisms h to (3.11.3) and we get

$$\gamma_{44}v_{s,1} + \gamma_{45}v_{s,2} + \gamma_{43}v_{s,3} = 0.$$

From this, it implies $\gamma_{43} = \gamma_{44} = \gamma_{45} = 0$. The proposition follows. □

4. The indecomposables of P_4 in degree $2^{s+1} - 2$

According to [8], $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_2 = 3$ with a basis consisting of all the following classes: $w_{1,1} = [1, 1, 0],\ w_{1,2} = [1, 0, 1],\ w_{1,3} = [0, 1, 1]$.

For $s \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+1}-2}$ is an $\mathbb{F}_2$-vector space with a basis consisting of all the following classes:

For $s \geqslant 2$,

$$w_{s,1} = [1, 2^s - 2, 2^s - 1], \quad w_{s,2} = [1, 2^s - 1, 2^s - 2],$$
$$w_{s,3} = [2^s - 1, 1, 2^s - 2], \quad w_{s,4} = [0, 2^s - 1, 2^s - 1],$$
$$w_{s,5} = [2^s - 1, 0, 2^s - 1], \quad w_{s,6} = [2^s - 1, 2^s - 1, 0].$$

For $s \geqslant 3$, $w_{s,7} = [3, 2^s - 3, 2^s - 2]$.

Hence, we have

Proposition 4.1.

1. $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_2$ *is an* $\mathbb{F}_2$*-vector space of dimension 6 with a basis consisting of all the classes represented by the following monomials:*

$$b_{1,1} = (0, 0, 1, 1), \quad b_{1,2} = (0, 1, 0, 1), \quad b_{1,3} = (0, 1, 1, 0),$$
$$b_{1,4} = (1, 0, 0, 1), \quad b_{1,5} = (1, 0, 1, 0), \quad b_{1,6} = (1, 1, 0, 0).$$

2. For $s \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+1}-2}$ *is an* $\mathbb{F}_2$*-vector space with a basis consisting of all the classes represented by the following monomials:*

For $s \geqslant 2$.

$$b_{s,1} = (0, 1, 2^s - 2, 2^s - 1), \quad b_{s,2} = (0, 1, 2^s - 1, 2^s - 2),$$
$$b_{s,3} = (0, 2^s - 1, 1, 2^s - 2), \quad b_{s,4} = (1, 0, 2^s - 2, 2^s - 1),$$
$$b_{s,5} = (1, 0, 2^s - 1, 2^s - 2), \quad b_{s,6} = (1, 2^s - 2, 0, 2^s - 1),$$
$$b_{s,7} = (1, 2^s - 2, 2^s - 1, 0), \quad b_{s,8} = (1, 2^s - 1, 0, 2^s - 2),$$
$$b_{s,9} = (1, 2^s - 1, 2^s - 2, 0), \quad b_{s,10} = (2^s - 1, 0, 1, 2^s - 2),$$
$$b_{s,11} = (2^s - 1, 1, 0, 2^s - 2), \quad b_{s,12} = (2^s - 1, 1, 2^s - 2, 0),$$
$$b_{s,13} = (0, 0, 2^s - 1, 2^s - 1), \quad b_{s,14} = (0, 2^s - 1, 0, 2^s - 1),$$
$$b_{s,15} = (0, 2^s - 1, 2^s - 1, 0), \quad b_{s,16} = (2^s - 1, 0, 0, 2^s - 1),$$
$$b_{s,17} = (2^s - 1, 0, 2^s - 1, 0), \quad b_{s,18} = (2^s - 1, 2^s - 1, 0, 0),$$

For $s \geqslant 3$.

$$b_{s,19} = (0, 3, 2^s - 3, 2^s - 2), \quad b_{s,20} = (3, 0, 2^s - 3, 2^s - 2),$$
$$b_{s,21} = (3, 2^s - 3, 0, 2^s - 2), \quad b_{s,22} = (3, 2^s - 3, 2^s - 2, 0).$$

By Proposition 2.8, we need only to compute $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-2}$.

The main result of this section is

Theorem 4.2.

1. $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_6$ *is an* $\mathbb{F}_2$*-vector space of dimension 6 with a basis consisting of all the classes represented by the following monomials:*

$$b_{2,19} = (1, 1, 2, 2), \quad b_{2,20} = (1, 2, 1, 2), \quad b_{2,21} = (1, 1, 1, 3),$$
$$b_{2,22} = (1, 1, 3, 1), \quad b_{2,23} = (1, 3, 1, 1), \quad b_{2,24} = (3, 1, 1, 1).$$

2. For any $s \geqslant 3$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-2}$ *is an* $\mathbb{F}_2$*-vector space of dimension* $\mu_1(s-1) + 13$ *with a basis consisting of all the classes represented by the following monomials:*

i. $\phi(a_{s-1,j})$*, where the monomials* $a_{s-1,j}; s \geqslant 3, 1 \leqslant j \leqslant \mu_1(s-1)$*, are determined as in Section 3 and the homomorphism* ϕ *is defined in Definition 2.9. Recall that* $\mu_1(2) = 15, \mu_1(3) = 35, \mu_1(s-1) = 45, s \geqslant 5$.

ii. *The monomials* $b_{s,j}, 23 \leqslant j \leqslant 35$*, are determined as follows:*

For $s \geqslant 3$,

$$\begin{aligned} b_{s,23} &= (1,1,2^s-2,2^s-2), & b_{s,24} &= (1,2^s-2,1,2^s-2),\\ b_{s,25} &= (1,2,2^s-4,2^s-1), & b_{s,26} &= (1,2,2^s-1,2^s-4),\\ b_{s,27} &= (1,2^s-1,2,2^s-4), & b_{s,28} &= (2^s-1,1,2,2^s-4),\\ b_{s,29} &= (1,2,2^s-3,2^s-2), & b_{s,30} &= (1,3,2^s-4,2^s-2),\\ b_{s,31} &= (1,3,2^s-2,2^s-4), & b_{s,32} &= (3,1,2^s-4,2^s-2),\\ b_{s,33} &= (3,1,2^s-2,2^s-4), & b_{s,34} &= (3,2^s-3,2,2^s-4). \end{aligned}$$

For $s = 3$, $b_{3,35} = (3,3,4,4)$.
For $s \geqslant 4$, $b_{s,35} = (3,5,2^s-6,2^s-4)$.

Theorem 4.2 is proved by combining the following propositions.

Proposition 4.3.
1. The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_6$ is generated by the six elements $[b_{2,j}], 19 \leqslant j \leqslant 24$.
2. For any $s \geqslant 3$, the $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-2}$ is generated by the $\mu_1(s-1)+13$ elements listed in Theorem 4.2.

The proof of this proposition is based on some lemmas.

Lemma 4.4 (Kameko [8])**.** *Let M be an 2×3-matrix and x the monomial corresponding to M. If $\tau_1(x) < \tau_2(x)$ then M is strictly inadmissible.*

From the proof of Proposition 3.3, Lemma 2.6 and Lemma 4.4, we easily obtain

Lemma 4.5. *Let x be a monomial of degree 2^s-3 in P_4 with $s \geqslant 3$. If x is inadmissible then there is a strictly inadmissible matrix Δ such that $\Delta \triangleright x$.*

Lemma 4.6. *Let x be an admissible monomial of degree $2^{s+1}-2$ in P_4. We have*
1. If $s = 1$ then $\tau(x) = (2;0)$.
2. If $s \geqslant 2$ then $\tau(x)$ is one of the following sequences

$$(\underbrace{2;2;\ldots;2}_{s \text{ times}}),\ (4;\underbrace{3;3;\ldots;3}_{s-2 \text{ times}};1).$$

Proof. It is clear that the lemma holds for $s = 1, 2$. Suppose $s \geqslant 3$. Observe that $z = (2^s-1, 2^s-1, 0, 0)$ is the minimal spike of degree $2^{s+1}-2$ in P_4 and $\tau(z) = (\underbrace{2;2;\ldots;2}_{s \text{ times}})$. Since $2^{s+1}-2$ is even, using Theorem 2.11, we obtain $\tau_1(x) = 2$ or $\tau_1(x) = 4$. Suppose $\tau_1(x) = 2$. Then, using Theorem 2.13 and Proposition 2.7, we get either $\tau(x) = (\underbrace{2;2;\ldots;2}_{s \text{ times}})$ or $\tau(x) = (\underbrace{2;2;\ldots;2}_{a \text{ times}};3;\tau_{a+2}(x);\ldots)$, for some $a \geqslant 1$.

If $\tau(x) = (\underbrace{2;2;\ldots;2}_{a \text{ times}};3;\tau_{a+2}(x);\ldots)$ then

$$n = \deg x = 2^a + 2^{a+1}\big(2 + \sum_{i\geqslant a+2} 2^{i-a-2}\tau_i(x)\big) - 2.$$

From this, it implies $\alpha(n+2) \geqslant 2$ and we have a contradiction.

Suppose that $\tau_1(x) = 4$. Then $x = \phi(x')$ with x' a monomial of degree $2^s - 3$. Since x is admissible, using Lemma 4.5 and Theorem 2.5, we see that x' is also admissible. By Lemma 3.4, we have

$$\tau(x') = (\tau_2(x);\tau_3(x);\ldots;\tau_i(x);\ldots) = (\underbrace{3;3;\ldots;3}_{s-2 \text{ times}};1).$$

The lemma is proved. □

Lemma 4.7. *The following matrices are strictly inadmissible*

$$\Delta_{17} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}, \quad \Delta_{18} = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix},$$

$$\Delta_{19} = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}, \quad \Delta_{20} = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{pmatrix}.$$

Proof. By a direct calculation, we have

$$(1,2,2,1) = Sq^1(2,1,1,1) + Sq^2(1,1,1,1) + (1,2,1,2) + (1,1,2,2),$$
$$(2,1,1,2) = Sq^1(1,1,1,2) + (1,2,1,2) + (1,1,2,2),$$
$$(2,1,2,1) = Sq^1(1,1,2,1) + (1,2,2,1) + (1,1,2,2),$$
$$(2,2,1,1) = Sq^1(1,2,1,1) + (1,2,2,1) + (1,2,1,2).$$

The lemma is proved. □

Lemma 4.8. *The following matrices are strictly inadmissible*

$$\Delta_{21} = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \ \Delta_{22} = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}, \ \Delta_{23} = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix},$$

$$\Delta_{24} = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{pmatrix}, \ \Delta_{25} = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}, \ \Delta_{26} = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{pmatrix},$$

$$\Delta_{27} = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{pmatrix}, \ \Delta_{28} = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \end{pmatrix}, \ \Delta_{29} = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix},$$

$$\Delta_{30} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}, \ \Delta_{31} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 \end{pmatrix}, \ \Delta_{32} = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix}.$$

Proof. See [8]. □

Lemma 4.9. *The following matrices are strictly inadmissible*

$$\Delta_{33} = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \ \Delta_{34} = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \ \Delta_{35} = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}.$$

Proof. A direct calculation shows

$$\begin{aligned}(1,6,3,4) &= Sq^1\big((1,5,3,4)+(1,3,5,4)+(4,3,3,3)+(1,5,4,3)\\ &\quad +(1,3,4,5)+(1,4,5,3)+(1,4,3,5)\big)+Sq^2\big((2,3,3,4)\\ &\quad +(2,3,4,3)+(1,6,2,3)+(2,4,3,3)\big)+(1,3,6,4)+(1,6,2,5)\\ &\quad +(1,3,4,6)+(1,4,6,3)+(1,4,3,6) \mod \mathcal{L}_4(2;2;2),\\ (3,4,1,6) &= Sq^1\big((3,4,1,5)+(3,3,4,3)\big)+Sq^2\big((5,2,2,3)\\ &\quad +(2,3,4,3)\big)+Sq^4(3,2,2,3)+(3,2,4,5)+(3,3,4,4)\\ &\quad +(2,5,4,3)+(2,3,4,5) \mod \mathcal{L}_4(2;2;2),\\ (3,4,3,4) &= Sq^1\big((3,3,3,4)+(1,4,4,4)\big)+Sq^2(2,3,3,4)\\ &\quad +(2,5,3,4)+(2,3,5,4)+(3,3,4,4).\end{aligned}$$

The lemma follows. □

Lemma 4.10. *The following matrices are strictly inadmissible*

$$\Delta_{36}=\begin{pmatrix}0&1&0&1\\0&1&0&1\\0&0&1&1\end{pmatrix},\ \Delta_{37}=\begin{pmatrix}0&1&1&0\\0&1&1&0\\0&0&1&1\end{pmatrix},\ \Delta_{38}=\begin{pmatrix}0&1&1&0\\0&1&1&0\\0&1&0&1\end{pmatrix},$$

$$\Delta_{39}=\begin{pmatrix}1&0&0&1\\1&0&0&1\\0&0&1&1\end{pmatrix},\ \Delta_{40}=\begin{pmatrix}1&0&1&0\\1&0&1&0\\0&0&1&1\end{pmatrix},\ \Delta_{41}=\begin{pmatrix}1&0&0&1\\1&0&0&1\\0&1&0&1\end{pmatrix},$$

$$\Delta_{42}=\begin{pmatrix}1&0&1&0\\1&0&1&0\\0&1&1&0\end{pmatrix},\ \Delta_{43}=\begin{pmatrix}1&1&0&0\\1&1&0&0\\0&1&0&1\end{pmatrix},\ \Delta_{44}=\begin{pmatrix}1&1&0&0\\1&1&0&0\\0&1&1&0\end{pmatrix},$$

$$\Delta_{45}=\begin{pmatrix}1&0&1&0\\1&0&1&0\\1&0&0&1\end{pmatrix},\ \Delta_{46}=\begin{pmatrix}1&1&0&0\\1&1&0&0\\1&0&0&1\end{pmatrix},\ \Delta_{47}=\begin{pmatrix}1&1&0&0\\1&1&0&0\\1&0&1&0\end{pmatrix}.$$

Proof. See [8]. □

Lemma 4.11. *The following matrices are strictly inadmissible*

$$\Delta_{48}=\begin{pmatrix}1&1&0&0\\0&1&1&0\\0&1&0&1\\0&0&1&1\end{pmatrix},\ \Delta_{49}=\begin{pmatrix}1&1&0&0\\1&0&1&0\\1&0&0&1\\0&0&1&1\end{pmatrix},\ \Delta_{50}=\begin{pmatrix}1&1&0&0\\1&1&0&0\\0&0&1&1\\0&0&1&1\end{pmatrix},$$

$$\Delta_{51}=\begin{pmatrix}1&1&0&0\\1&0&0&1\\0&1&0&1\\0&0&1&1\end{pmatrix},\ \Delta_{52}=\begin{pmatrix}1&1&0&0\\1&0&1&0\\0&1&1&0\\0&0&1&1\end{pmatrix},\ \Delta_{53}=\begin{pmatrix}1&1&0&0\\1&1&0&0\\1&1&0&0\\0&0&1&1\end{pmatrix}.$$

Proof. By a direct calculation, we have

$$\begin{aligned}
(1,7,10,12) &= Sq^1\big((1,7,9,12)+(1,9,7,12)+(4,7,7,11)+(1,8,7,13)\\
&\quad +(1,7,8,13)\big)+Sq^2\big(2,7,7,12)+(2,8,7,11)+(2,7,8,11)\big)\\
&\quad +Sq^4\big((1,4,7,14)+(1,6,7,12)\big)+(1,6,11,12)+(1,4,11,14)\\
&\quad +(1,7,8,14) \mod \mathcal{L}_4(2;2;2;2),\\
(7,1,10,12) &= \overline{\varphi}_1(1,7,10,12),\\
(3,3,12,12) &= Sq^1\big((3,3,11,12)+(3,8,7,11)\big)\\
&\quad +Sq^2\big((2,3,11,12)+(2,8,7,11)\big)+Sq^4(3,4,7,12)+(2,5,11,12)\\
&\quad +(2,3,13,12)+(2,8,7,13) \mod \mathcal{L}_4(2;2;2;2),\\
(3,5,8,14) &= Sq^1\big((3,3,9,14)+(3,3,5,18)\big)\\
&\quad +Sq^2\big((2,3,9,14)+(5,3,6,14)\big)+Sq^4(3,3,6,14)\\
&\quad +(2,5,9,14)+(3,4,9,14) \mod \mathcal{L}_4(2;2;2;2),\\
(3,5,14,8) &= \overline{\varphi}_3(3,5,14,8),\\
(7,7,8,8) &= Sq^1(7,7,7,8)+Sq^2(7,6,7,8)+Sq^4\big((4,7,7,8)\\
&\quad +(5,6,7,8)\big)+(7,6,9,8)+(4,11,7,8)+(4,7,11,8)\\
&\quad +(5,10,7,8)+(5,6,11,8) \mod \mathcal{L}_4(2;2;2;2).
\end{aligned}$$

Here $\overline{\varphi}_1$ and $\overline{\varphi}_3$ are defined in Section 2.

The lemma is proved. □

Proof of Proposition 4.3. Let x is an admissible monomial of degree $2^{s+1}-2$ in P_4 for $s \geqslant 2$. Then either $\tau_1(x)=4$ or $\tau_1(x)=2$. If $\tau_1(x)=4$ then using Lemma 4.6 and Lemma 4.5, we get $x=\phi(x')$ with x' an admissible monomial of degree 2^s-3. By Proposition 3.1 and Theorem 3.2, $x'=a_{s-1,j}$, for some $1\leqslant j\leqslant \mu_1(s-1)$.

Suppose $\tau_1(x)=2$. By Lemma 4.6, $\tau_j(x)=2, j=1,2,\ldots,s$, and $\tau_j(x)=0$ for $j>s$. Then $x=b_{1,i}y^2$ with a monomial y of degree 2^s-2, $\tau_j(y)=\tau_{j+1}(x)=2$ for $j=1,2,\ldots,s-1$, and $1\leqslant i\leqslant 6$. Now prove the proposition by showing that if $x\neq b_{s,j}$, for all j then there is a strictly inadmissible matrix Δ such that $\Delta\triangleright x$. We prove this by induction on s.

For $s=2$, a direct computation shows

$$\begin{array}{llll}
b_{2,1}=b_{1,2}b_{1,1}^2, & b_{2,2}=b_{1,3}b_{1,1}^2, & b_{2,3}=b_{1,3}b_{1,2}^2, & b_{2,4}=b_{1,4}b_{1,1}^2,\\
b_{2,5}=b_{1,5}b_{1,1}^2, & b_{2,6}=b_{1,4}b_{1,2}^2, & b_{2,7}=b_{1,5}b_{1,3}^2, & b_{2,8}=b_{1,6}b_{1,2}^2,\\
b_{2,9}=b_{1,6}b_{1,3}^2, & b_{2,10}=b_{1,5}b_{1,4}^2, & b_{2,11}=b_{1,6}b_{1,4}^2, & b_{2,12}=b_{1,6}b_{1,5}^2,\\
b_{2,13}=b_{1,1}b_{1,1}^2, & b_{2,14}=b_{1,2}b_{1,2}^2, & b_{2,15}=b_{1,3}b_{1,3}^2, & b_{2,16}=b_{1,4}b_{1,4}^2,\\
b_{2,17}=b_{1,5}b_{1,5}^2, & b_{2,18}=b_{1,6}b_{1,6}^2, & b_{2,19}=b_{1,6}b_{1,1}^2, & b_{2,20}=b_{1,5}b_{1,2}^2,\\
b_{2,21}=\phi(a_{1,1}), & b_{2,22}=\phi(a_{1,2}), & b_{2,23}=\phi(a_{1,3}), & b_{2,24}=\phi(a_{1,4}).
\end{array}$$

We have $\Delta_{17}\triangleright b_{1,4}b_{1,3}^2$; $\Delta_{18}\triangleright b_{1,3}b_{1,4}^2$; $\Delta_{19}\triangleright b_{1,2}b_{1,5}^2$; $\Delta_{20}\triangleright b_{1,1}b_{1,6}^2$; $\Delta_{21}\triangleright b_{1,1}b_{1,2}^2$; $\Delta_{22}\triangleright b_{1,1}b_{1,3}^2$; $\Delta_{23}\triangleright b_{1,2}b_{1,3}^2$; $\Delta_{24}\triangleright b_{1,1}b_{1,4}^2$; $\Delta_{25}\triangleright b_{1,1}b_{1,5}^2$; $\Delta_{26}\triangleright b_{1,2}b_{1,4}^2$; $\Delta_{27}\triangleright b_{1,3}b_{1,5}^2$; $\Delta_{28}\triangleright b_{1,2}b_{1,6}^2$; $\Delta_{29}\triangleright b_{1,3}b_{1,6}^2$; $\Delta_{30}\triangleright b_{1,4}b_{1,5}^2$; $\Delta_{31}\triangleright b_{1,4}b_{1,6}^2$; $\Delta_{32}\triangleright b_{1,5}b_{1,6}^2$.

From these equalities, Lemmas 4.7, 4.8 and Theorem 2.5, we see that the case $s = 2$ is true.

For $s = 3$, by a direct computation, we obtain

$$\begin{array}{llll}
b_{3,1} = b_{1,2}b_{2,13}^2, & b_{3,2} = b_{1,3}b_{2,13}^2, & b_{3,3} = b_{1,3}b_{2,14}^2, & b_{3,4} = b_{1,4}b_{2,13}^2,\\
b_{3,5} = b_{1,5}b_{2,13}^2, & b_{3,6} = b_{1,4}b_{2,14}^2, & b_{3,7} = b_{1,5}b_{2,15}^2, & b_{3,8} = b_{1,6}b_{2,14}^2,\\
b_{3,9} = b_{1,6}b_{2,15}^2, & b_{3,10} = b_{1,5}b_{2,16}^2, & b_{3,11} = b_{1,6}b_{2,16}^2, & b_{3,12} = b_{1,6}b_{2,17}^2,\\
b_{3,13} = b_{1,1}b_{2,13}^2, & b_{3,14} = b_{1,2}b_{2,14}^2, & b_{3,15} = b_{1,3}b_{2,15}^2, & b_{3,16} = b_{1,4}b_{2,16}^2,\\
b_{3,17} = b_{1,5}b_{2,17}^2, & b_{3,18} = b_{1,6}b_{2,18}^2, & b_{3,19} = b_{1,3}b_{2,1}^2, & b_{3,20} = b_{1,5}b_{2,4}^2,\\
b_{3,21} = b_{1,6}b_{2,6}^2, & b_{3,22} = b_{1,6}b_{2,7}^2, & b_{3,23} = b_{1,6}b_{2,13}^2, & b_{3,24} = b_{1,5}b_{2,14}^2,\\
b_{3,25} = b_{1,4}b_{2,1}^2, & b_{3,26} = b_{1,5}b_{2,2}^2, & b_{3,27} = b_{1,6}b_{2,3}^2, & b_{3,28} = b_{1,6}b_{2,10}^2,\\
b_{3,29} = b_{1,5}b_{2,1}^2, & b_{3,30} = b_{1,6}b_{2,1}^2, & b_{3,31} = b_{1,6}b_{2,2}^2, & b_{3,32} = b_{1,6}b_{2,4}^2,\\
b_{3,33} = b_{1,6}b_{2,5}^2, & b_{3,34} = b_{1,6}b_{2,20}^2, & b_{3,35} = b_{1,6}b_{2,19}^2. &
\end{array}$$

We have $\Delta_{21}\triangleright b_{1,1}b_{2,j}^2$ for $j = 1, 14$; $\Delta_{17}\triangleright b_{1,4}b_{2,j}^2$; $\Delta_{22}\triangleright b_{1,1}b_{2,j}^2$; $\Delta_{23}\triangleright b_{1,2}b_{2,j}^2$ for $j = 2, 3, 15$; $\Delta_{18}\triangleright b_{1,3}b_{2,j}^2$; $\Delta_{24}\triangleright b_{1,1}b_{2,j}^2$; $\Delta_{26}\triangleright b_{1,2}b_{2,j}^2$ for $j = 4, 6, 16$; $\Delta_{19}\triangleright b_{1,2}b_{2,j}^2$; $\Delta_{25}\triangleright b_{1,1}b_{2,j}^2$; $\Delta_{27}\triangleright b_{1,3}b_{2,j}^2$; $\Delta_{30}\triangleright b_{1,4}b_{2,j}^2$ for $j = 5, 7, 10, 17, 20$; $\Delta_{20}\triangleright b_{1,1}b_{2,j}^2$; $\Delta_{28}\triangleright b_{1,2}b_{2,j}^2$; $\Delta_{29}\triangleright b_{1,3}b_{2,j}^2$; $\Delta_{31}\triangleright b_{1,4}b_{2,j}^2$; $\Delta_{32}\triangleright b_{1,5}b_{2,j}^2$ for $j = 8, 9, 11, 12, 18, 19$; $\Delta_{33}\triangleright b_{1,5}b_{2,3}^2$; $\Delta_{34}\triangleright b_{1,5}b_{2,6}^2$; $\Delta_{35}\triangleright b_{1,5}b_{2,20}^2$; $\Delta_{36}\triangleright b_{1,2}b_{2,1}^2$; $\Delta_{37}\triangleright b_{1,3}b_{2,2}^2$; $\Delta_{38}\triangleright b_{1,3}b_{2,3}^2$; $\Delta_{39}\triangleright b_{1,4}b_{2,4}^2$; $\Delta_{40}\triangleright b_{1,5}b_{2,5}^2$; $\Delta_{41}\triangleright b_{1,4}b_{2,6}^2$; $\Delta_{42}\triangleright b_{1,5}b_{2,7}^2$; $\Delta_{43}\triangleright b_{1,6}b_{2,8}^2$; $\Delta_{44}\triangleright b_{1,6}b_{2,9}^2$; $\Delta_{45}\triangleright b_{1,5}b_{2,10}^2$; $\Delta_{46}\triangleright b_{1,6}b_{2,11}^2$; $\Delta_{47}\triangleright b_{1,6}b_{2,12}^2$.

From the above equalities, Lemmas 4.7, 4.8, 4.9, 4.10 and Theorem 2.5, we see that if $x \neq b_{3,j}, 1 \leqslant j \leqslant 35$, then x is inadmissible. The case $s = 3$ holds.

Suppose $s \geqslant 4$ and the proposition holds for s. By the inductive hypothesis and Theorem 2.5, it suffices to consider monomials $x = b_{1,i}b_{s,j}^2$ with $1 \leqslant i \leqslant 6$ and $1 \leqslant j \leqslant 35$. We have

$$\begin{array}{llll}
b_{s+1,1} = b_{1,2}b_{s,13}^2, & b_{s+1,2} = b_{1,3}b_{s,13}^2, & b_{s+1,3} = b_{1,3}b_{s,14}^2, & b_{s+1,4} = b_{1,4}b_{s,13}^2,\\
b_{s+1,5} = b_{1,5}b_{s,13}^2, & b_{s+1,6} = b_{1,4}b_{s,14}^2, & b_{s+1,7} = b_{1,5}b_{s,15}^2, & b_{s+1,8} = b_{1,6}b_{s,14}^2,\\
b_{s+1,9} = b_{1,6}b_{s,15}^2, & b_{s+1,10} = b_{1,5}b_{s,16}^2, & b_{s+1,11} = b_{1,6}b_{s,16}^2, & b_{s+1,12} = b_{1,6}b_{s,17}^2,\\
b_{s+1,13} = b_{1,1}b_{s,13}^2, & b_{s+1,14} = b_{1,2}b_{s,14}^2, & b_{s+1,15} = b_{1,3}b_{s,15}^2, & b_{s+1,16} = b_{1,4}b_{s,16}^2,\\
b_{s+1,17} = b_{1,5}b_{s,17}^2, & b_{s+1,18} = b_{1,6}b_{s,18}^2, & b_{s+1,19} = b_{1,3}b_{s,1}^2, & b_{s+1,20} = b_{1,5}b_{s,4}^2,\\
b_{s+1,21} = b_{1,6}b_{s,6}^2, & b_{s+1,22} = b_{1,6}b_{s,7}^2, & b_{s+1,23} = b_{1,6}b_{s,13}^2, & b_{s+1,24} = b_{1,5}b_{s,14}^2,\\
b_{s+1,25} = b_{1,4}b_{s,1}^2, & b_{s+1,26} = b_{1,5}b_{s,2}^2, & b_{s+1,27} = b_{1,6}b_{s,3}^2, & b_{s+1,28} = b_{1,6}b_{s,10}^2,\\
b_{s+1,29} = b_{1,5}b_{s,1}^2, & b_{s+1,30} = b_{1,6}b_{s,1}^2, & b_{s+1,31} = b_{1,6}b_{s,2}^2, & b_{s+1,32} = b_{1,6}b_{s,4}^2,\\
b_{s+1,33} = b_{1,6}b_{s,5}^2, & b_{s+1,34} = b_{1,6}b_{s,24}^2, & b_{s+1,35} = b_{1,6}b_{s,29}^2. &
\end{array}$$

For $x \neq b_{s,j}, 1 \leqslant j \leqslant 35$, we have $\Delta_{21}\triangleright b_{1,1}b_{s,j}^2$ for $j = 1, 14$; $\Delta_{17}\triangleright b_{1,4}b_{s,j}^2$, $\Delta_{22}\triangleright b_{1,1}b_{s,j}^2$, $\Delta_{23}\triangleright b_{1,2}b_{s,j}^2$ for $j = 2, 3, 15, 19$; $\Delta_{18}\triangleright b_{1,3}b_{s,j}^2$, $\Delta_{24}\triangleright b_{1,1}b_{s,j}^2$, $\Delta_{26}\triangleright b_{1,2}b_{s,j}^2$ for $j = 4, 6, 16, 25$; $\Delta_{19}\triangleright b_{1,2}b_{s,j}^2$, $\Delta_{25}\triangleright b_{1,1}b_{s,j}^2$, $\Delta_{27}\triangleright b_{1,3}b_{s,j}^2$, $\Delta_{30}\triangleright b_{1,4}b_{s,j}^2$ for $j = 5, 7, 10, 17, 20, 24, 26, 29$; $\Delta_{20}\triangleright b_{1,1}b_{s,j}^2$, $\Delta_{28}\triangleright b_{1,2}b_{s,j}^2$, $\Delta_{29}\triangleright b_{1,3}b_{s,j}^2$, $\Delta_{31}\triangleright b_{1,4}b_{s,j}^2$, $\Delta_{32}\triangleright b_{1,5}b_{s,j}^2$ for $j = 8, 9, 11, 12, 18, 21, 22, 23, 27, 28, 30, 31, 32, 33, 34, 35$; $\Delta_{33}\triangleright b_{1,5}b_{s,j}^2$ for $j = 3, 19$; $\Delta_{34}\triangleright b_{1,5}b_{s,j}^2$ for $j = 6, 25$; $\Delta_{35}\triangleright b_{1,5}b_{s,j}^2$ for $j = 24, 29$; $\Delta_{36}\triangleright b_{1,2}b_{s,1}^2$; $\Delta_{37}\triangleright b_{1,3}b_{s,2}^2$; $\Delta_{38}\triangleright b_{1,3}b_{s,j}^2$ for $j = 3, 19$; $\Delta_{39}\triangleright b_{1,4}b_{s,4}^2$; $\Delta_{40}\triangleright b_{1,5}b_{s,5}^2$; $\Delta_{41}\triangleright b_{1,4}b_{s,j}^2$ for $j =$

6, 25; $\Delta_{42} \triangleright b_{1,5}b_{s,j}^2$ for $j = 7, 26$; $\Delta_{43} \triangleright b_{1,6}b_{s,j}^2$ for $j = 8, 30$; $\Delta_{44} \triangleright b_{1,6}b_{s,j}^2$ for $j = 9, 27, 31$; $\Delta_{45} \triangleright b_{1,5}b_{s,j}^2$ for $j = 10, 20$; $\Delta_{46} \triangleright b_{1,6}b_{s,j}^2$ for $j = 11, 21, 32$; $\Delta_{47} \triangleright b_{1,6}b_{s,j}^2$ for $j = 12, 22, 28, 33, 34$; $\Delta_{48} \triangleright b_{1,6}b_{s,19}^2$; $\Delta_{49} \triangleright b_{1,6}b_{s,20}^2$; $\Delta_{50} \triangleright b_{1,6}b_{s,23}^2$; $\Delta_{51} \triangleright b_{1,6}b_{s,25}^2$; $\Delta_{52} \triangleright b_{1,6}b_{s,26}^2$; $\Delta_{53} \triangleright b_{1,6}b_{3,35}^2$, for $s = 3$ and $\Delta_{47} \triangleright b_{1,6}b_{s,35}^2$, for $s \geqslant 4$.

From the above equalities, Lemmas 4.7, 4.8, 4.9, 4.10, 4.11, we see that the case $s + 1$ is true. The proof is completed. □

Now, we prove that the elements listed in Theorem 4.2 are linearly independent.

Proposition 4.12. *The elements* $[b_{2,i}], 19 \leqslant i \leqslant 24$, *are linearly independent.*

Proof. Suppose there is a linear relation

$$\sum_{19 \leqslant i \leqslant 24} \gamma_i [b_{2,i}] = 0,$$

with $\gamma_i \in \mathbb{F}_2$.

Since the monomials $b_{2,i}, i = 21, 22, 23, 24$, are the spikes, we get $\gamma_{21} = \gamma_{22} = \gamma_{23} = \gamma_{24} = 0$. Then, applying the homomorphisms f_1, f_2 to the above linear relation, we obtain

$$\gamma_{20}[3, 1, 2] = 0, \quad \gamma_{19}[3, 1, 2] = 0.$$

So, $\gamma_{19} = \gamma_{20} = 0$. The proposition follows. □

Proposition 4.13. *The elements* $[b_{3,i}]$ *for* $23 \leqslant i \leqslant 35$, *and* $[\phi(a_{2,j})]$ *for* $1 \leqslant j \leqslant 15$, *are linearly independent.*

Proof. Suppose that there is a linear relation

$$\sum_{23 \leqslant i \leqslant 35} \gamma_i [b_{3,i}] + \sum_{1 \leqslant j \leqslant 15} \eta_j [\phi(a_{2,j})] = 0, \tag{4.13.1}$$

with $\gamma_i, \eta_j \in \mathbb{F}_2$.

Apply the squaring operation Sq_*^0 to (4.13.1) and we get

$$\sum_{1 \leqslant j \leqslant 15} \eta_j [a_{2,j}] = 0.$$

Since $\{[a_{2,j}]; 1 \leqslant j \leqslant 15\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_5$, we obtain $\eta_j = 0, 1 \leqslant j \leqslant 15$. Hence, (4.13.1) becomes

$$\sum_{23 \leqslant i \leqslant 35} \gamma_i [b_{3,i}] = 0. \tag{4.13.2}$$

Now, we prove $\gamma_i = 0, 23 \leqslant i \leqslant 35$.

Applying the homomorphisms $f_t, t = 1, 2, \ldots, 5$, to the relation (4.13.2), we obtain

$$\gamma_{25}[1,6,7] + \gamma_{26}[1,7,6] + \gamma_{24}[7,1,6] + \gamma_{29}[3,5,6] = 0,$$
$$\gamma_{25}[1,6,7] + \gamma_{27}[1,7,6] + \gamma_{\{23,31,32,35\}}[7,1,6] + \gamma_{30}[3,5,6] = 0,$$
$$\gamma_{26}[1,6,7] + \gamma_{27}[1,7,6] + \gamma_{\{23,24,29,30,33,34,35\}}[7,1,6] + \gamma_{31}[3,5,6] = 0,$$
$$\gamma_{25}[1,6,7] + \gamma_{\{23,24,29,30,33,34,35\}}[1,7,6] + \gamma_{28}[7,1,6] + \gamma_{32}[3,5,6] = 0,$$
$$\gamma_{26}[1,6,7] + \gamma_{\{23,31,32,35\}}[1,7,6] + \gamma_{28}[7,1,6] + \gamma_{33}[3,5,6] = 0.$$

From the above relation, we get $\gamma_i = 0, i = 24, \ldots, 34$, and $\gamma_{23} = \gamma_{35}$. So, the relation (4.13.2) becomes

$$\gamma_{23}([1,1,6,6] + [3,3,4,4]) = 0.$$

Now, we prove that $[1,1,6,6] + [3,3,4,4] \neq 0$. Suppose the contrary, that the polynomial $(1,1,6,6)+(3,3,4,4)$ is hit. Then by the unstable property of the action of $\mathcal{A}$ on the polynomial algebra, we have

$$(1,1,6,6) + (3,3,4,4) = Sq^1(A) + Sq^2(B) + Sq^4(C),$$

for some polynomials $A \in (R_4)_{13}, B \in (R_4)_{12}, C \in (R_4)_{10}$. Let $Sq^2Sq^2Sq^2$ act on the both sides of the above equality. Since $Sq^2Sq^2Sq^2Sq^1 = 0$ and $Sq^2Sq^2Sq^2Sq^2 = 0$, we get

$$Sq^2Sq^2Sq^2((1,1,6,6) + (3,3,4,4)) = Sq^2Sq^2Sq^2Sq^4(C).$$

On the other hand, by a direct computation, it is not difficult to check that

$$Sq^2Sq^2Sq^2((1,1,6,6) + (3,3,4,4)) \neq Sq^2Sq^2Sq^2Sq^4(C),$$

for all $C \in (R_4)_{10}$. This is a contradiction. Hence, $[1,1,6,6]+[3,3,4,4] \neq 0$ and $\gamma_{23} = \gamma_{35} = 0$. The proposition is proved. □

Proposition 4.14. *For $s \geqslant 4$, the $\mu_1(s-1) + 13$ elements listed in Theorem 4.2 are linearly independent.*

Proof. Suppose that there is a linear relation

$$\sum_{23 \leqslant i \leqslant 35} \gamma_i[b_{s,i}] + \sum_{1 \leqslant j \leqslant \mu_1(s-1)} \eta_j[\phi(a_{s-1,j})] = 0, \tag{4.14.1}$$

with $\gamma_i, \eta_j \in \mathbb{F}_2$.

Applying the squaring operation Sq^0_* to (4.14.1), we get

$$\sum_{1 \leqslant j \leqslant \mu_1(s-1)} \eta_j[a_{s-1,j}] = 0.$$

Since $\{[a_{s-1,j}]; 1 \leqslant j \leqslant \mu_1(s-1)\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{2^s-3}$, we obtain $\eta_j = 0, 1 \leqslant j \leqslant \mu_1(s-1)$. Hence, (4.14.1) becomes

$$\sum_{23 \leqslant i \leqslant 35} \gamma_i[b_{s,i}] = 0. \tag{4.14.2}$$

Apply the homomorphisms $f_t, t = 1, 2, \ldots, 6$, to the relation (4.14.2) and we obtain

$$\begin{aligned}
&\gamma_{25}w_{s,1} + \gamma_{26}w_{s,2} + \gamma_{24}w_{s,3} + \gamma_{29}w_{s,7} = 0,\\
&\gamma_{25}w_{s,1} + \gamma_{27}w_{s,2} + \gamma_{\{23,31,32\}}w_{s,3} + \gamma_{30}w_{s,7} = 0,\\
&\gamma_{26}w_{s,1} + \gamma_{27}w_{s,2} + \gamma_{\{23,24,29,30,33,34,35\}}w_{s,3} + \gamma_{31}w_{s,7} = 0,\\
&\gamma_{25}w_{s,1} + \gamma_{\{23,24,29,30,33,34,35\}}w_{s,2} + \gamma_{28}w_{s,3} + \gamma_{32}w_{s,7} = 0,\\
&\gamma_{26}w_{s,1} + \gamma_{\{23,31,32\}}w_{s,2} + \gamma_{28}w_{s,3} + \gamma_{33}w_{s,7} = 0,\\
&\gamma_{24}w_{s,1} + \gamma_{27}w_{s,2} + \gamma_{28}w_{s,3} + \gamma_{34}w_{s,7} = 0.
\end{aligned}$$

From these relations, we get $\gamma_i = 0, i = 23, 24, \ldots, 35$.

The proposition follows. □

Remark 4.15. By a direct calculation, we can easily show that

$$[1, 1, 6, 6] + [3, 3, 4, 4]$$

is an $GL_4(\mathbb{F}_2)$-invariant in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{14}$.

5. The indecomposables of P_4 in degree $2^{s+1} - 1$

First of all, we recall a result in [8] on the dimension of the $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+1}-1}$.

Set $\rho_1(1) = 7, \rho_1(2) = 10, \rho_1(3) = 13$ and $\rho_1(s) = 14$ for $s \geqslant 4$.

According to Kameko [8], $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+1}-1}$ is an $\mathbb{F}_2$-vector space of dimension $\rho_1(s)$ with a basis consisting of all the following classes:

For $s \geqslant 1$,

$u_{s,1} = [0, 1, 2^{s+1} - 2]$, $u_{s,2} = [1, 0, 2^{s+1} - 2]$, $u_{s,3} = [1, 2^{s+1} - 2, 0]$,
$u_{s,4} = [0, 0, 2^{s+1} - 1]$, $u_{s,5} = [0, 2^{s+1} - 1, 0]$, $u_{s,6} = [2^{s+1} - 1, 0, 0]$.

For $s = 1$, $u_{1,7} = [1, 1, 1]$.

For $s \geqslant 2$,

$u_{s,7} = [1, 2, 2^{s+1} - 4]$, $u_{s,8} = [1, 2^s - 1, 2^s - 1]$,
$u_{s,9} = [2^s - 1, 1, 2^s - 1]$, $u_{s,10} = [2^s - 1, 2^s - 1, 1]$.

For $s \geqslant 3$,

$u_{s,11} = [3, 2^s - 3, 2^s - 1]$, $u_{s,12} = [3, 2^s - 1, 2^s - 3]$,
$u_{s,13} = [2^s - 1, 3, 2^s - 3]$.

For $s \geqslant 4$, $u_{s,14} = [7, 2^s - 5, 2^s - 3]$.

Set $\rho_2(1) = 14, \rho_2(2) = 26, \rho_2(3) = 38$ and $\rho_2(s) = 42$ for $s \geqslant 4$. From this result we easily obtain

Proposition 5.1. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+1}-1}$ is an $\mathbb{F}_2$-vector space of dimension $\rho_2(s)$ with a basis consisting of all the classes represented by the following monomials:*

For $s \geqslant 1$,

$$\begin{aligned} c_{s,1} &= (0,0,1,2^{s+1}-2), & c_{s,2} &= (0,1,0,2^{s+1}-2),\\ c_{s,3} &= (0,1,2^{s+1}-2,0), & c_{s,4} &= (1,0,0,2^{s+1}-2),\\ c_{s,5} &= (1,0,2^{s+1}-2,0), & c_{s,6} &= (1,2^{s+1}-2,0,0),\\ c_{s,7} &= (0,0,0,2^{s+1}-1), & c_{s,8} &= (0,0,2^{s+1}-1,0),\\ c_{s,9} &= (0,2^{s+1}-1,0,0), & c_{s,10} &= (2^{s+1}-1,0,0,0). \end{aligned}$$

For $s = 1$,

$$c_{1,11} = (0,1,1,1),\ c_{1,12} = (1,0,1,1),\ c_{1,13} = (1,1,0,1),\ c_{1,14} = (1,1,1,0).$$

For $s \geqslant 2$,

$$\begin{aligned} c_{s,11} &= (0,1,2,2^{s+1}-4), & c_{s,12} &= (1,0,2,2^{s+1}-4),\\ c_{s,13} &= (1,2,0,2^{s+1}-4), & c_{s,14} &= (1,2,2^{s+1}-4,0),\\ c_{s,15} &= (0,1,2^s-1,2^s-1), & c_{s,16} &= (0,2^s-1,1,2^s-1),\\ c_{s,17} &= (0,2^s-1,2^s-1,1), & c_{s,18} &= (1,0,2^s-1,2^s-1),\\ c_{s,19} &= (1,2^s-1,0,2^s-1), & c_{s,20} &= (1,2^s-1,2^s-1,0),\\ c_{s,21} &= (2^s-1,0,1,2^s-1), & c_{s,22} &= (2^s-1,0,2^s-1,1),\\ c_{s,23} &= (2^s-1,1,0,2^s-1), & c_{s,24} &= (2^s-1,1,2^s-1,0),\\ c_{s,25} &= (2^s-1,2^s-1,0,1), & c_{s,26} &= (2^s-1,2^s-1,1,0). \end{aligned}$$

For $s \geqslant 3$,

$$\begin{aligned} c_{s,27} &= (0,3,2^s-3,2^s-1), & c_{s,28} &= (0,3,2^s-1,2^s-3),\\ c_{s,29} &= (0,2^s-1,3,2^s-3), & c_{s,30} &= (3,0,2^s-3,2^s-1),\\ c_{s,31} &= (3,0,2^s-1,2^s-3), & c_{s,32} &= (3,2^s-3,0,2^s-1),\\ c_{s,33} &= (3,2^s-3,2^s-1,0), & c_{s,34} &= (3,2^s-1,0,2^s-3),\\ c_{s,35} &= (3,2^s-1,2^s-3,0), & c_{s,36} &= (2^s-1,0,3,2^s-3),\\ c_{s,37} &= (2^s-1,3,0,2^s-3), & c_{s,38} &= (2^s-1,3,2^s-3,0). \end{aligned}$$

For $s \geqslant 4$,

$$\begin{aligned} c_{s,39} &= (0,7,2^s-5,2^s-3), & c_{s,40} &= (7,0,2^s-5,2^s-3),\\ c_{s,41} &= (7,2^s-5,0,2^s-3), & c_{s,42} &= (7,2^s-5,2^s-3,0). \end{aligned}$$

Observe that $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_3 = 0$. So we need only to determine the space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-1}$ for $s \geqslant 2$.

Set $\rho_3(2) = 9, \rho_3(3) = 37, \rho_3(4) = 47$ and $\rho_3(s) = 43$ for $s \geqslant 5$. We have

Theorem 5.2. *For any $s \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-1}$ is an $\mathbb{F}_2$-vector space of dimension $\rho_3(s)$ with a basis consisting of all the classes represented by the following monomials:*

For $s \geqslant 2$,

$$\begin{aligned} d_{s,1} &= (1,1,2^s-2,2^s-1), & d_{s,2} &= (1,1,2^s-1,2^s-2),\\ d_{s,3} &= (1,2^s-2,1,2^s-1), & d_{s,4} &= (1,2^s-2,2^s-1,1),\\ d_{s,5} &= (1,2^s-1,1,2^s-2), & d_{s,6} &= (1,2^s-1,2^s-2,1),\\ d_{s,7} &= (2^s-1,1,1,2^s-2), & d_{s,8} &= (2^s-1,1,2^s-2,1). \end{aligned}$$

For $s = 2$, $d_{2,9} = (1,2,2,2)$.

For $s \geqslant 3$,

$$\begin{aligned}
&d_{s,9} = (1, 2, 2^s - 3, 2^s - 1), && d_{s,10} = (1, 2, 2^s - 1, 2^s - 3),\\
&d_{s,11} = (1, 2^s - 1, 2, 2^s - 3), && d_{s,12} = (2^s - 1, 1, 2, 2^s - 3),\\
&d_{s,13} = (1, 3, 2^s - 4, 2^s - 1), && d_{s,14} = (1, 3, 2^s - 1, 2^s - 4),\\
&d_{s,15} = (1, 2^s - 1, 3, 2^s - 4), && d_{s,16} = (3, 1, 2^s - 4, 2^s - 1),\\
&d_{s,17} = (3, 1, 2^s - 1, 2^s - 4), && d_{s,18} = (3, 2^s - 1, 1, 2^s - 4),\\
&d_{s,19} = (2^s - 1, 1, 3, 2^s - 4), && d_{s,20} = (2^s - 1, 3, 1, 2^s - 4),\\
&d_{s,21} = (1, 3, 2^s - 3, 2^s - 2), && d_{s,22} = (1, 3, 2^s - 2, 2^s - 3),\\
&d_{s,23} = (1, 2^s - 2, 3, 2^s - 3), && d_{s,24} = (3, 1, 2^s - 3, 2^s - 2),\\
&d_{s,25} = (3, 1, 2^s - 2, 2^s - 3), && d_{s,26} = (3, 2^s - 3, 1, 2^s - 2),\\
&d_{s,27} = (3, 2^s - 3, 2^s - 2, 1), && d_{s,28} = (3, 2^s - 3, 2, 2^s - 3),\\
&d_{s,29} = (3, 3, 2^s - 4, 2^s - 3), && d_{s,30} = (3, 3, 2^s - 3, 2^s - 4),\\
&d_{s,31} = (3, 2^s - 3, 3, 2^s - 4).
\end{aligned}$$

For $s = 3$,

$$\begin{aligned}
&d_{3,32} = (3, 4, 1, 7), && d_{3,33} = (3, 4, 7, 1), && d_{3,34} = (3, 7, 4, 1),\\
&d_{3,35} = (7, 3, 4, 1), && d_{3,36} = (3, 4, 3, 5), && d_{3,37} = (1, 2, 4, 8).
\end{aligned}$$

For $s \geqslant 4$,

$$\begin{aligned}
&d_{s,32} = (1, 6, 2^s - 5, 2^s - 3), && d_{s,33} = (1, 7, 2^s - 6, 2^s - 3),\\
&d_{s,34} = (7, 1, 2^s - 6, 2^s - 3), && d_{s,35} = (1, 7, 2^s - 5, 2^s - 4),\\
&d_{s,36} = (7, 1, 2^s - 5, 2^s - 4), && d_{s,37} = (7, 2^s - 5, 1, 2^s - 4),\\
&d_{s,38} = (3, 4, 2^s - 5, 2^s - 3), && d_{s,39} = (3, 5, 2^s - 6, 2^s - 3),\\
&d_{s,40} = (3, 5, 2^s - 5, 2^s - 4), && d_{s,41} = (3, 7, 2^s - 7, 2^s - 4),\\
&d_{s,42} = (7, 3, 2^s - 7, 2^s - 4), && d_{s,43} = (1, 2, 4, 2^{s+1} - 8).
\end{aligned}$$

For $s = 4$,

$$\begin{aligned}
&d_{4,44} = (3, 7, 8, 13), && d_{4,45} = (7, 3, 8, 13),\\
&d_{4,46} = (7, 7, 8, 9), && d_{4,47} = (7, 7, 9, 8).
\end{aligned}$$

We prove Theorem 5.2 by proving some propositions.

Proposition 5.3. *For any $s \geqslant 2$, the $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-1}$ is generated by the $\rho_3(s)$ elements listed in Theorem 5.2.*

This proposition is proved by combining the following lemmas. First, we determine the τ-sequence of an admissible monomial of degree $2^{s+1} - 1$ in P_4.

Lemma 5.4. *Let x be an admissible monomial of degree $2^{s+1} - 1$ in P_4. We have*

1. *If $s = 1$ then either $\tau(x) = (1; 1)$ or $\tau(x) = (3; 0)$.*
2. *If $s = 2$ then either $\tau(x) = (1; 1; 1)$ or $\tau(x) = (1; 3)$ or $\tau(x) = (3; 2)$.*
3. *If $s \geqslant 3$ then $\tau(x)$ is one of the following sequences*

$$(\underbrace{1; 1; \ldots; 1}_{s+1 \text{ times}}), \quad (3; \underbrace{2; 2; \ldots; 2}_{s-1 \text{ times}}).$$

We need the following for the proof of Lemma 5.4.

From the proof of Proposition 4.3, Lemma 2.6, Lemma 4.4 and Lemma 4.5, we get the following

Lemma 5.5. *Let x be a monomial of degree $2^s - 2$ in P_4 with $s \geqslant 2$. If x is inadmissible then there is a strictly inadmissible matrix Δ such that $\Delta \triangleright x$.*

Lemma 5.6. *Let M be an 2×4-matrix and x the monomial corresponding to M. If either $\tau(x) = (1;2)$ or $\tau(x) = (1;3)$, $x \neq (1,2,2,2)$ then M is strictly inadmissible.*

Proof. If $\tau(x) = (1;2)$ the lemma follows from Lemma 4.4, since one of the columns of M is zero. We prove the lemma for the case $\tau(x) = (1;3)$ and $x \neq (1,2,2,2)$.

If one of the columns of M is zero then the lemma follows from Lemma 4.4. So, we assume that all columns of M are non-zero. Since $x \neq (1,2,2,2)$, x is one of the monomials $(2,1,2,2), (2,2,1,2), (2,2,2,1)$. It is easy to see that

$$\begin{aligned}(2,1,2,2) &= Sq^1(1,1,2,2) + (1,2,2,2),\\ (2,2,1,2) &= Sq^1(1,2,1,2) + (1,2,2,2),\\ (2,2,2,1) &= Sq^1(1,2,2,1) + (1,2,2,2).\end{aligned}$$

Hence, the lemma is proved. □

Lemma 5.7. *Let $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4 \in \{0,1\}$. If $\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4 < 4$ then the following matrix is strictly inadmissible*

$$M = \begin{pmatrix} \varepsilon_1 & \varepsilon_2 & \varepsilon_3 & \varepsilon_4\\ 1 & 0 & 0 & 0\\ 0 & 1 & 1 & 1\end{pmatrix}.$$

Proof. The monomial corresponding to the matrix M is $x = (\varepsilon_1 + 2, \varepsilon_2 + 4, \varepsilon_3 + 4, \varepsilon_4 + 4)$. If $\varepsilon_1 = 0$ then

$$\begin{aligned}x = Sq^1(1, \varepsilon_2 + 4, \varepsilon_3 + 4, \varepsilon_4 + 4) &+ \varepsilon_2(1, \varepsilon_2 + 5, \varepsilon_3 + 4, \varepsilon_4 + 4)\\ &+ \varepsilon_3(1, \varepsilon_2 + 4, \varepsilon_3 + 5, \varepsilon_4 + 4) + \varepsilon_4(1, \varepsilon_2 + 4, \varepsilon_3 + 4, \varepsilon_4 + 5).\end{aligned}$$

Hence, the lemma is true. We assume that $\varepsilon_1 = 1$.

If $\varepsilon_2 + \varepsilon_3 + \varepsilon_4 = 0$ then

$$x = (3,4,4,4) = Sq^1(3,3,4,4) + Sq^2(2,3,4,4) + (2,5,4,4).$$

If $\varepsilon_2 + \varepsilon_3 + \varepsilon_4 = 1$ then there is an $\mathcal{A}$-homomorphism $f : P_4 \to P_4$ induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = f(3,5,4,4)$. We have

$$\begin{aligned}(3,5,4,4) = Sq^1\big((3,6,2,4) + (3,4,4,4) + (3,3,1,8)\big)&\\ + Sq^2(5,3,2,4) + Sq^4(3,3,2,4) + (4,3,1,8) + (3,4,1,8).&\end{aligned}$$

If $\varepsilon_2+\varepsilon_3+\varepsilon_4 = 2$ then there is an $\mathcal{A}$-homomorphism $f' : P_4 \to P_4$ induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = f'(3,5,5,4)$. We have

$$\begin{aligned}(3,5,5,4) &= Sq^1\big((4,5,5,2) + (5,5,5,1) + (6,3,5,2) + (3,6,5,2)\\ &\quad + (3,5,6,2)\big) + Sq^2\big((3,5,5,2) + (6,3,5,1) + (3,6,5,1) + (3,5,6,1)\big)\\ &\quad + (3,6,6,2) + (6,4,6,1) + (8,3,5,1) + (3,8,5,1) + (3,5,8,1).\end{aligned}$$

The homomorphisms f, f' send monomials to monomials and preserve the associated τ-sequences. The lemma follows. □

Lemma 5.8. *Let $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4 \in \{0,1\}$. If $\varepsilon_1+\varepsilon_2+\varepsilon_3+\varepsilon_4 > 0$ then the following matrix is strictly inadmissible*

$$M = \begin{pmatrix} 1 & 0 & 0 & 0\\ 0 & 1 & 1 & 1\\ \varepsilon_1 & \varepsilon_2 & \varepsilon_3 & \varepsilon_4\end{pmatrix}.$$

Proof. The monomial corresponding to the matrix M is $x = (1+4\varepsilon_1, 2+4\varepsilon_2, 2+4\varepsilon_3, 2+4\varepsilon_4)$. If $\varepsilon_1 = 1$ then

$$\begin{aligned}x &= Sq^2(3, 2+4\varepsilon_2, 2+4\varepsilon_3, 2+4\varepsilon_4) + (3, 4\varepsilon_2+4, 4\varepsilon_3+2, 4\varepsilon_4+2)\\ &\quad + (3, 4\varepsilon_2+2, 4\varepsilon_3+4, 4\varepsilon_4+2) + (3, 4\varepsilon_2+2, 4\varepsilon_3+2, 4\varepsilon_4+4).\end{aligned}$$

Hence, the lemma holds. Suppose that $\varepsilon_1 = 0$.

If $\varepsilon_2+\varepsilon_3+\varepsilon_4 = 1$ then there is an $\mathcal{A}$-homomorphism $f : P_4 \to P_4$ induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = f(1,6,2,2)$. A direct computation shows

$$(1,6,2,2) = Sq^1(2,6,1,1) + Sq^2(1,6,1,1) + Sq^4(1,4,1,1).$$

If $\varepsilon_2+\varepsilon_3+\varepsilon_4 = 2$ then there is an $\mathcal{A}$-homomorphism $f' : P_4 \to P_4$ induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = f'(1,6,6,2)$. We have

$$\begin{aligned}(1,6,6,2) &= Sq^1\big((2,5,5,2) + (2,3,5,4)\big)\\ &\quad + Sq^2\big((1,5,5,2) + (1,3,5,4)\big) + (1,4,6,4).\end{aligned}$$

If $\varepsilon_2+\varepsilon_3+\varepsilon_4 = 3$ then

$$\begin{aligned}x = (1,6,6,6) &= Sq^1\big((2,5,5,6) + (2,3,5,8)\big)\\ &\quad + Sq^2\big((1,5,5,6) + (1,3,5,8)\big) + (1,4,6,8).\end{aligned}$$

Hence, the lemma is proved. □

Proof of Lemma 5.4. Obviously, the first part of the lemma is true. Suppose $s \geqslant 2$. Since $2^{s+1}-1$ is odd, we have either $\tau_1(x) = 1$ or $\tau_1(x) = 3$.

If $\tau_1(x) = 1$ then using Lemmas 5.6, 5.7, 5.8, we get either $\tau(x) = (1;3)$, for $s = 2$, or $\tau(x) = (\underbrace{1;1;\ldots;1}_{s+1 \text{ times}})$, for $s \geqslant 2$.

Suppose that $\tau_1(x) = 3$. Then $x = z_i y^2$, $1 \leqslant i \leqslant 4$, where z_1, z_2, z_3, z_4 are defined as in the proof of Proposition 3.3 and y is a monomial of degree

$2^s - 2$. Since x is admissible, using Lemma 5.5 and Theorem 2.5, we see that y is also admissible and $\tau_i(y) = \tau_{i+1}(x), i \geqslant 1$. Using Lemma 4.6 and Proposition 2.7, we obtain

$$\tau(y) = (\tau_2(x); \tau_3(x); \ldots; \tau_i(x); \ldots) = (\underbrace{2; 2; \ldots; 2}_{s-1 \text{ times}}).$$

The lemma is proved. □

Lemma 5.9. *The following matrices are strictly inadmissible*

$$\Delta_{54} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \Delta_{55} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \Delta_{56} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix},$$
$$\Delta_{57} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \Delta_{58} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \Delta_{59} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

Proof. See [8]. □

Lemma 5.10. *The following matrices are strictly inadmissible*

$$\Delta_{60} = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{pmatrix}, \quad \Delta_{61} = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix},$$
$$\Delta_{62} = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{pmatrix}, \quad \Delta_{63} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{pmatrix}.$$

Proof. A direct computation shows

$$(2,1,1,3) = Sq^1(1,1,1,3) + (1,2,1,3) + (1,1,2,3) + (1,1,1,4),$$
$$(2,1,3,1) = Sq^1(1,1,3,1) + (1,2,3,1) + (1,1,3,2) + (1,1,4,1),$$
$$(2,3,1,1) = Sq^1(1,3,1,1) + (1,3,2,1) + (1,3,1,2) + (1,4,1,1),$$
$$(3,2,1,1) = Sq^1(3,1,1,1) + (3,1,2,1) + (3,1,1,2) + (4,1,1,1).$$

The lemma follows. □

Lemma 5.11. *The following matrices are strictly inadmissible*

$$\Delta_{64} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \ \Delta_{65} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \ \Delta_{66} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix},$$
$$\Delta_{67} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \ \Delta_{68} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \ \Delta_{69} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

Proof. See [8]. □

Lemma 5.12. *The following matrices are strictly inadmissible*

$$\Delta_{70} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \ \Delta_{71} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{pmatrix}, \ \Delta_{72} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{pmatrix},$$

$$\Delta_{73} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{pmatrix}, \ \Delta_{74} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \ \Delta_{75} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix},$$

$$\Delta_{76} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \ \Delta_{77} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}, \ \Delta_{78} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix},$$

$$\Delta_{79} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \ \Delta_{80} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \ \Delta_{81} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix},$$

$$\Delta_{82} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{pmatrix}, \ \Delta_{83} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \ \Delta_{84} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix},$$

$$\Delta_{85} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \ \Delta_{86} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \ \Delta_{87} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix},$$

$$\Delta_{88} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \quad \Delta_{89} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix},$$

$$\Delta_{90} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}, \quad \Delta_{91} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(3,12,1,15),(3,12,15,1),(3,15,12,1),(15,3,12,1),(3,4,9,15),$$
$$(3,4,15,9),(3,15,4,9),(15,3,4,9),(3,5,8,15),(3,5,15,8),$$
$$(3,15,5,8),(15,3,5,8),(7,11,12,1),(3,12,3,13),(3,5,9,14),$$
$$(3,5,14,9),(3,7,13,8),(7,3,13,8),(3,7,12,9),(7,3,12,9),$$
$$(7,11,4,9),(7,11,5,8).$$

If x is one of the four monomials $(3,12,1,15),(3,12,15,1),(3,15,12,1)$, $(15,3,12,1)$ then there is a homomorphism $\psi_1 : P_4 \to P_4$ of $\mathcal{A}$-modules

induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = \psi_1(3, 12, 1, 15)$. By a direct calculation, we get

$$\begin{aligned}(3,12,1,15) &= Sq^1\big((3,11,1,15) + (3,7,1,19)\big)\\ &\quad + Sq^2\big((2,11,1,15) + (5,7,2,15)\big) + Sq^4(3,7,2,15)\\ &\quad + (2,13,1,15) + (3,9,4,15) \mod \mathcal{L}_4(3;2;2;2),\end{aligned}$$

where $\tau(2, 13, 1, 15) = \tau(3, 9, 4, 15) = \tau(x)$ and $\sigma(2, 13, 1, 15), \sigma(3, 9, 4, 15) < \sigma(x)$.

If x' is one of the four monomials $(3, 4, 9, 15), (3, 4, 15, 9), (3, 15, 4, 9), (15, 3, 4, 9)$ then there is a homomorphism $\psi_2 : P_4 \to P_4$ of $\mathcal{A}$-modules induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x' = \psi_2(3, 4, 9, 15)$. We have

$$\begin{aligned}(3,4,9,15) &= Sq^1\big((3,1,11,15) + (3,1,7,19)\big)\\ &\quad + Sq^2\big((2,1,11,15) + (5,2,7,15)\big) + Sq^4(3,2,7,15)\\ &\quad + (2,1,13,15) + (3,1,12,15) \mod \mathcal{L}_4(3;2;2;2),\end{aligned}$$

where $\tau(2, 1, 13, 15) = \tau(3, 1, 12, 15) = \tau(x')$, $\sigma(2, 1, 13, 15), \sigma(3, 1, 12, 15) < \sigma(x')$.

If x'' is one of the four monomials $(3, 5, 8, 15), (3, 5, 15, 8), (3, 15, 5, 8), (15, 3, 5, 8)$ then there is a homomorphism $\psi_3 : P_4 \to P_4$ of $\mathcal{A}$-modules induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x'' = \varphi_3(3, 5, 8, 15)$. A direct computation shows

$$\begin{aligned}(3,5,8,15) &= Sq^1\big((3,3,9,15) + (3,3,5,19)\big)\\ &\quad + Sq^2\big((2,3,9,15) + (5,3,6,15)\big) + Sq^4(3,3,6,15)\\ &\quad + (2,5,9,15) + (3,4,9,15) \mod \mathcal{L}_4(3;2;2;2).\end{aligned}$$

Here $\tau(2, 5, 9, 15) = \tau(3, 4, 9, 15) = \tau(x'')$ and $\sigma(2, 5, 9, 15), \sigma(3, 4, 9, 15) < \sigma(x'')$.

Similarly, we have

$$\begin{aligned}(7,11,12,1) &= Sq^1(7,13,9,1) + Sq^2\big((7,11,10,1)\\ &\quad + (7,14,7,1) + (7,11,9,2)\big) + Sq^4(5,14,7,1)\\ &\quad + (5,14,11,1) + (7,11,9,4) \mod \mathcal{L}_4(3;2;2;2),\\ (3,12,3,13) &= Sq^1\big((3,11,3,13) + (3,7,3,17)\big) + Sq^2\big((5,7,3,14)\\ &\quad + (2,11,3,13)\big) + Sq^4(3,7,3,14) + (2,13,3,13) + (2,11,5,13)\\ &\quad + (3,9,5,14) + (3,11,4,13) \mod \mathcal{L}_4(3;2;2;2),\\ (3,5,9,14) &= Sq^1\big((3,3,11,13) + (3,3,7,17)\big) + Sq^2\big((5,3,7,14)\\ &\quad + (2,3,11,13)\big) + Sq^4(3,3,7,14) + (3,4,11,13) + (3,3,12,13)\\ &\quad + (2,5,11,13) + (2,3,13,13) \mod \mathcal{L}_4(3;2;2;2),\end{aligned}$$

$$\begin{aligned}(3,7,13,8) &= Sq^1(3,7,11,9) + Sq^2\big((5,7,11,6) + (5,7,7,10)\\ &\quad + (2,7,11,9)\big) + Sq^4\big((3,11,7,6) + (3,5,7,12) + (3,5,13,6)\big)\\ &\quad + Sq^8(3,7,7,6) + (3,7,9,12) + (3,7,12,9) + (2,7,13,9)\\ &\quad + (3,5,11,12) + (3,5,13,10) \mod \mathcal{L}_4(3;2;2;2),\\ (3,7,12,9) &= Sq^1(3,7,9,11) + Sq^2\big((5,7,10,7) + (5,7,6,11)\\ &\quad + (2,7,9,11)\big) + Sq^4\big((3,11,6,7) + (3,5,12,7) + (3,5,6,13)\big)\\ &\quad + Sq^8(3,7,6,7) + (3,7,8,13) + (3,7,9,12) + (2,7,9,13)\\ &\quad + (3,5,12,11) + (3,5,10,13) \mod \mathcal{L}_4(3;2;2;2),\\ (7,11,4,9) &= Sq^1(7,13,1,9) + Sq^2\big((7,11,2,9)\\ &\quad + (7,14,1,7) + (7,11,1,10)\big) + Sq^4(5,14,1,7)\\ &\quad + (7,11,1,12) + (5,14,1,11) \mod \mathcal{L}_4(3;2;2;2),\\ (7,11,5,8) &= Sq^1(7,11,3,9) + Sq^2\big((7,13,3,6) + (7,7,3,12)\\ &\quad + (7,10,3,9)\big) + Sq^4\big((11,7,3,6) + (5,11,5,6) + (5,7,5,10)\big)\\ &\quad + Sq^8(7,7,3,6) + (7,9,5,10) + (5,11,9,6) + (5,7,9,10)\\ &\quad + (7,11,4,9) + (7,10,5,9) \mod \mathcal{L}_4(3;2;2;2).\end{aligned}$$

The lemma is proved. □

Lemma 5.13. *The following matrices are strictly inadmissible*

$$\Delta_{92} = \begin{pmatrix}1&1&0&1\\1&1&0&0\\0&1&0&1\\0&0&1&1\\0&0&1&1\end{pmatrix}, \quad \Delta_{93} = \begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&0&1\\0&0&1&1\\0&0&1&1\end{pmatrix}.$$

Proof. A direct computation shows

$$\begin{aligned}(7,3,24,29) &= Sq^1\big((7,3,23,29) + (7,3,15,37) + (7,3,19,33)\big)\\ &\quad + Sq^2\big((7,2,23,29) + (7,5,19,30) + (7,3,15,36) + (7,4,15,35)\big)\\ &\quad + Sq^4\big((4,3,23,29) + (5,2,23,29) + (11,3,15,30) + (5,3,21,30)\big)\\ &\quad + Sq^8(7,3,15,30) + (4,3,27,29) + (7,2,25,29) + (5,2,27,29)\\ &\quad + (5,3,25,30) \mod \mathcal{L}_4(3;2;2;2;2).\\ (3,7,24,29) &= \overline{\varphi}_1(7,3,24,29).\end{aligned}$$

The lemma follows. □

Now we prove Proposition 5.3.

Proof of Proposition 5.3. Suppose x is an admissible monomial of degree $2^{s+1}-1$ in P_4. For $s=2$, using Lemma 5.4 we have either $\tau(x) = (1;1;1)$ or $\tau(x) = (1;3)$ or $\tau(x) = (3;2)$. By Lemma 5.5 and Proposition 4.4, it

suffices to consider $x = a_{1,i}c_{1,j}^2, i = 1, 2, 3, 4,\ 1 \leqslant j \leqslant 14$ or $x = z_i b_{1,j}^2, i = 1, 2, 3, 4,\ j = 1, 2, 3, 4, 5, 6$. A direct computation shows

$$\begin{array}{llll}
c_{2,1} = c_{1,2}c_{1,7}^2, & c_{2,2} = c_{1,3}c_{1,7}^2, & c_{2,3} = c_{1,3}c_{1,8}^2, & c_{2,4} = c_{1,4}c_{1,7}^2,\\
c_{2,5} = c_{1,4}c_{1,8}^2, & c_{2,6} = c_{1,4}c_{1,9}^2, & c_{2,7} = c_{1,1}c_{1,7}^2, & c_{2,8} = c_{1,2}c_{1,8}^2,\\
c_{2,9} = c_{1,3}c_{1,9}^2, & c_{2,10} = c_{1,4}c_{1,10}^2, & c_{2,11} = c_{1,3}c_{1,1}^2, & c_{2,12} = c_{1,4}c_{1,1}^2,\\
c_{2,13} = c_{1,4}c_{1,2}^2, & c_{2,14} = c_{1,4}c_{1,3}^2, & c_{2,15} = z_1b_{1,1}^2, & c_{2,16} = z_1b_{1,2}^2,\\
c_{2,17} = z_1b_{1,3}^2, & c_{2,18} = z_2b_{1,1}^2, & c_{2,19} = z_3b_{1,2}^2, & c_{2,20} = z_4b_{1,3}^2,\\
c_{2,21} = z_2b_{1,4}^2, & c_{2,22} = z_2b_{1,5}^2, & c_{2,23} = z_3b_{1,4}^2, & c_{2,24} = z_4b_{1,5}^2,\\
c_{2,25} = z_3b_{1,6}^2, & c_{2,26} = z_4b_{1,6}^2, & d_{2,1} = z_3b_{1,1}^2, & d_{2,2} = z_4b_{1,1}^2,\\
d_{2,3} = z_2b_{1,2}^2, & d_{2,4} = z_2b_{1,3}^2, & d_{2,5} = z_4b_{1,2}^2, & d_{2,6} = z_3b_{1,3}^2,\\
d_{2,7} = z_4b_{1,4}^2, & d_{2,8} = z_3b_{1,5}^2, & d_{2,9} = a_{1,4}c_{1,11}^2. &
\end{array}$$

For $x \neq c_{2,i}, 1 \leqslant i \leqslant 26$, and $x \neq d_{2,j}, 1 \leqslant j \leqslant 9$, we have $\Delta_{54} \triangleright a_{1,1}c_{1,j}^2$ for $j = 1,\ 8$; $\Delta_{55} \triangleright a_{1,1}c_{1,j}^2$, $\Delta_{56} \triangleright a_{1,2}c_{1,j}^2$ for $j = 2,\ 3,\ 9$; $\Delta_{57} \triangleright a_{1,1}c_{1,j}^2$, $\Delta_{58} \triangleright a_{1,2}c_{1,j}^2$, $\Delta_{59} \triangleright a_{1,3}c_{1,j}^2$ for $j = 4,\ 5,\ 6,\ 10$; $\Delta_{60} \triangleright z_1b_{1,4}^2$; $\Delta_{61} \triangleright z_1b_{1,5}^2$; $\Delta_{62} \triangleright z_1b_{1,6}^2$; $\Delta_{63} \triangleright z_2b_{1,6}^2$; $\Delta_{64} \triangleright a_{1,2}c_{1,1}^2$; $\Delta_{65} \triangleright a_{1,3}c_{1,2}^2$; $\Delta_{66} \triangleright a_{1,3}c_{1,3}^2$; $\Delta_{67} \triangleright a_{1,4}c_{1,4}^2$; $\Delta_{68} \triangleright a_{1,4}c_{1,5}^2$; $\Delta_{69} \triangleright a_{1,4}c_{1,6}^2$. Using Lemma 5.6, we see that if $x = a_{1,i}c_{1,j}^2, i = 1, 2, 3, 4,\ j = 11, 12, 13, 14$ and $x \neq (1, 2, 2, 2)$ then x is inadmissible. These equalities, Lemmas 5.9, 5.10, 5.11 and Theorem 2.5 imply that the case $s = 2$ is true.

Suppose $s \geqslant 3$. Using Lemma 5.4, we get either $\tau_1(x) = 1$ or $\tau_1(x) = 3$. If $\tau_1(x) = 1$ then $\tau_i(x) = 1, 1 \leqslant i \leqslant s+1$. So, $x = a_{1,i}y_1^2$, where $i = 1, 2, 3, 4$ and y_1 is a monomial with $\tau_i(y_1) = 1, i = 1, 2, \ldots, s$. If $\tau_1(x) = 3$ then $\tau_i(x) = 2, 2 \leqslant i \leqslant s-1$. Hence $x = z_i y_2^2$, where y_2 is a monomial of degree $2^s - 2$. Now we prove that if $x \neq c_{s,i}$ and $x \neq d_{s,j}$ for all i, j then there is a strictly inadmissible matrix Δ such that $\Delta \triangleright x$. The proof proceeds by induction on s.

Let $s = 3$. Since the proposition holds for $s = 2$, using Lemma 5.5, it suffices to consider $y_1 = c_{2,i}$ and $y_2 = b_{2,j}$ for some i, j. A direct computation shows

$$\begin{array}{llll}
c_{3,1} = a_{1,2}c_{2,7}^2, & c_{3,2} = a_{1,3}c_{2,7}^2, & c_{3,3} = a_{1,3}c_{2,8}^2, & c_{3,4} = a_{1,4}c_{2,7}^2,\\
c_{3,5} = a_{1,4}c_{2,8}^2, & c_{3,6} = a_{1,4}c_{2,9}^2, & c_{3,7} = a_{1,1}c_{2,7}^2, & c_{3,8} = a_{1,2}c_{2,8}^2,\\
c_{3,9} = a_{1,3}c_{2,9}^2, & c_{3,10} = a_{1,4}c_{2,10}^2, & c_{3,11} = a_{1,3}c_{2,1}^2, & c_{3,12} = a_{1,4}c_{2,1}^2,\\
c_{3,13} = a_{1,4}c_{2,2}^2, & c_{3,14} = a_{1,4}c_{2,3}^2, & c_{3,15} = z_1b_{2,13}^2, & c_{3,16} = z_1b_{2,14}^2,\\
c_{3,17} = z_1b_{2,15}^2, & c_{3,18} = z_2b_{2,13}^2, & c_{3,19} = z_3b_{2,14}^2, & c_{3,20} = z_4b_{2,15}^2,\\
c_{3,21} = z_2b_{2,16}^2, & c_{3,22} = z_2b_{2,17}^2, & c_{3,23} = z_3b_{2,16}^2, & c_{3,24} = z_4b_{2,17}^2,\\
c_{3,25} = z_3b_{2,18}^2, & c_{3,26} = z_4b_{2,18}^2, & c_{3,27} = z_1b_{2,1}^2, & c_{3,28} = z_1b_{2,2}^2,\\
c_{3,29} = z_1b_{2,3}^2, & c_{3,30} = z_2b_{2,4}^2, & c_{3,31} = z_2b_{2,5}^2, & c_{3,32} = z_3b_{2,6}^2,\\
c_{3,33} = z_4b_{2,7}^2, & c_{3,34} = z_3b_{2,8}^2, & c_{3,35} = z_4b_{2,9}^2, & c_{3,36} = z_2b_{2,10}^2,\\
c_{3,37} = z_3b_{2,11}^2, & c_{3,38} = z_4b_{2,12}^2, & d_{3,1} = z_3b_{2,13}^2, & d_{3,2} = z_4b_{2,13}^2,\\
d_{3,3} = z_2b_{2,14}^2, & d_{3,4} = z_2b_{2,15}^2, & d_{3,5} = z_4b_{2,14}^2, & d_{3,6} = z_3b_{2,15}^2,\\
d_{3,7} = z_4b_{2,16}^2, & d_{3,8} = z_3b_{2,17}^2, & d_{3,9} = z_2b_{2,1}^2, & d_{3,10} = z_2b_{2,2}^2,\\
d_{3,11} = z_3b_{2,3}^2, & d_{3,12} = z_3b_{2,10}^2, & d_{3,13} = z_3b_{2,1}^2, & d_{3,14} = z_4b_{2,2}^2,\\
d_{3,15} = z_4b_{2,3}^2, & d_{3,16} = z_3b_{2,4}^2, & d_{3,17} = z_4b_{2,5}^2, & d_{3,18} = z_4b_{2,8}^2,
\end{array}$$

$$\begin{array}{llll} d_{3,19} = z_4b_{2,10}^2, & d_{3,20} = z_4b_{2,11}^2, & d_{3,21} = z_4b_{2,1}^2, & d_{3,22} = z_3b_{2,2}^2, \\ d_{3,23} = z_2b_{2,3}^2, & d_{3,24} = z_4b_{2,4}^2, & d_{3,25} = z_3b_{2,5}^2, & d_{3,26} = z_4b_{2,6}^2, \\ d_{3,27} = z_3b_{2,7}^2, & d_{3,28} = z_3b_{2,20}^2, & d_{3,29} = z_3b_{2,19}^2, & d_{3,30} = z_4b_{2,19}^2, \\ d_{3,31} = z_4b_{2,20}^2, & d_{3,32} = z_2b_{2,6}^2, & d_{3,33} = z_2b_{2,7}^2, & d_{3,34} = z_3b_{2,9}^2, \\ d_{3,35} = z_3b_{2,12}^2, & d_{3,36} = z_2b_{2,20}^2, & d_{3,37} = a_{1,4}c_{2,11}^2. & \end{array}$$

For $x \neq c_{3,i}, 1 \leqslant i \leqslant 38$ and $x \neq d_{3,j}, 1 \leqslant j \leqslant 37$, we have $\Delta_{54} \triangleright a_{1,1}c_{2,j}^2$ for $j = 1, 8$; $\Delta_{55} \triangleright a_{1,1}c_{2,j}^2$, $\Delta_{56} \triangleright a_{1,2}c_{2,j}^2$ for $j = 2, 3, 9, 11$; $\Delta_{57} \triangleright a_{1,1}c_{2,j}^2$, $\Delta_{58} \triangleright a_{1,2}c_{2,j}^2$, $\Delta_{59} \triangleright a_{1,3}c_{2,j}^2$ for $j = 4, 5, 6, 10, 12, 13, 14$; $\Delta_{60} \triangleright z_1b_{2,j}^2$ for $j = 4, 6, 16$; $\Delta_{61} \triangleright z_1b_{2,j}^2$ for $j = 5, 7, 10, 17, 20$; $\Delta_{62} \triangleright z_1b_{2,j}^2$; $\Delta_{63} \triangleright z_2b_{2,j}^2$ for $j = 8, 9, 11, 12, 18, 19$; $\Delta_{64} \triangleright a_{1,2}c_{2,1}^2$; $\Delta_{65} \triangleright a_{1,3}c_{2,2}^2$; $\Delta_{66} \triangleright a_{1,3}c_{2,j}^2$ for $j = 3, 11$; $\Delta_{67} \triangleright a_{1,4}c_{2,4}^2$; $\Delta_{68} \triangleright a_{1,4}c_{2,j}^2$ for $j = 5, 12$; $\Delta_{69} \triangleright a_{1,4}c_{2,j}^2$ for $j = 6, 13, 14$. So, the case $s = 3$ is true.

Suppose that $s \geqslant 3$ and the proposition holds for s. We have

$$\begin{array}{llll} c_{s+1,1} = a_{1,2}c_{s,7}^2, & c_{s+1,2} = a_{1,3}c_{s,7}^2, & c_{s+1,3} = a_{1,3}c_{s,8}^2, & c_{s+1,4} = a_{1,4}c_{s,7}^2, \\ c_{s+1,5} = a_{1,4}c_{s,8}^2, & c_{s+1,6} = a_{1,4}c_{s,9}^2, & c_{s+1,7} = a_{1,1}c_{s,7}^2, & c_{s+1,8} = a_{1,2}c_{s,8}^2, \\ c_{s+1,9} = a_{1,3}c_{s,9}^2, & c_{s+1,10} = a_{1,4}c_{s,10}^2, & c_{s+1,11} = a_{1,3}c_{s,1}^2, & c_{s+1,12} = a_{1,4}c_{s,1}^2, \\ c_{s+1,13} = a_{1,4}c_{s,2}^2, & c_{s+1,14} = a_{1,4}c_{s,3}^2, & c_{s+1,15} = z_4b_{s,13}^2, & c_{s+1,16} = z_4b_{s,14}^2, \\ c_{s+1,17} = z_4b_{s,15}^2, & c_{s+1,18} = z_3b_{s,13}^2, & c_{s+1,19} = z_2b_{s,14}^2, & c_{s+1,20} = z_1b_{s,15}^2, \\ c_{s+1,21} = z_3b_{s,16}^2, & c_{s+1,22} = z_3b_{s,17}^2, & c_{s+1,23} = z_2b_{s,16}^2, & c_{s+1,24} = z_1b_{s,17}^2, \\ c_{s+1,25} = z_2b_{s,18}^2, & c_{s+1,26} = z_1b_{s,18}^2, & c_{s+1,27} = z_4b_{s,1}^2, & c_{s+1,28} = z_4b_{s,2}^2, \\ c_{s+1,29} = z_4b_{s,3}^2, & c_{s+1,30} = z_3b_{s,4}^2, & c_{s+1,31} = z_3b_{s,5}^2, & c_{s+1,32} = z_2b_{s,6}^2, \\ c_{s+1,33} = z_1b_{s,7}^2, & c_{s+1,34} = z_2b_{s,8}^2, & c_{s+1,35} = z_1b_{s,9}^2, & c_{s+1,36} = z_3b_{s,10}^2, \\ c_{s+1,37} = z_2b_{s,11}^2, & c_{s+1,38} = z_1b_{s,12}^2, & c_{s+1,39} = z_4b_{s,19}^2, & c_{s+1,40} = z_3b_{s,20}^2, \\ c_{s+1,41} = z_2b_{s,21}^2, & c_{s+1,42} = z_1b_{s,22}^2, & d_{s+1,1} = z_3b_{s,13}^2, & d_{s+1,2} = z_4b_{s,13}^2, \\ d_{s+1,3} = z_2b_{s,14}^2, & d_{s+1,4} = z_2b_{s,15}^2, & d_{s+1,5} = z_4b_{s,14}^2, & d_{s+1,6} = z_3b_{s,15}^2, \\ d_{s+1,7} = z_4b_{s,16}^2, & d_{s+1,8} = z_3b_{s,17}^2, & d_{s+1,9} = z_2b_{s,1}^2, & d_{s+1,10} = z_2b_{s,2}^2, \\ d_{s+1,11} = z_3b_{s,3}^2, & d_{s+1,12} = z_3b_{s,10}^2, & d_{s+1,13} = z_3b_{s,1}^2, & d_{s+1,14} = z_4b_{s,2}^2, \\ d_{s+1,15} = z_4b_{s,3}^2, & d_{s+1,16} = z_3b_{s,4}^2, & d_{s+1,17} = z_4b_{s,5}^2, & d_{s+1,18} = z_4b_{s,8}^2, \\ d_{s+1,19} = z_4b_{s,10}^2, & d_{s+1,20} = z_4b_{s,11}^2, & d_{s+1,21} = z_4b_{s,1}^2, & d_{s+1,22} = z_3b_{s,2}^2, \\ d_{s+1,23} = z_2b_{s,3}^2, & d_{s+1,24} = z_4b_{s,4}^2, & d_{s+1,25} = z_3b_{s,5}^2, & d_{s+1,26} = z_4b_{s,6}^2, \\ d_{s+1,27} = z_3b_{s,7}^2, & d_{s+1,28} = z_3b_{s,24}^2, & d_{s+1,29} = z_3b_{s,23}^2, & d_{s+1,30} = z_4b_{s,23}^2, \\ d_{s+1,31} = z_4b_{s,24}^2, & d_{s+1,32} = z_2b_{s,19}^2, & d_{s+1,33} = z_3b_{s,19}^2, & d_{s+1,34} = z_3b_{s,20}^2, \\ d_{s+1,35} = z_4b_{s,19}^2, & d_{s+1,36} = z_4b_{s,20}^2, & d_{s+1,37} = z_4b_{s,21}^2, & d_{s+1,38} = z_2b_{s,29}^2, \\ d_{s+1,39} = z_3b_{s,29}^2, & d_{s+1,40} = z_4b_{s,29}^2, & d_{s+1,41} = z_4b_{s,30}^2, & d_{s+1,42} = z_4b_{s,32}^2, \\ d_{s+1,43} = a_{1,4}c_{s,11}^2. & & & \end{array}$$

For $s = 3$, $d_{4,44} = z_3b_{3,30}^2$, $d_{4,45} = z_3b_{3,32}^2$, $d_{4,46} = z_3b_{3,35}^2$, $d_{4,47} = z_4b_{3,35}^2$.

If $x \neq c_{s,i}, 1 \leqslant i \leqslant 42$ and $x \neq d_{s,j}$ for all j then $\Delta_{54} \triangleright a_{1,1}c_{s,j}^2$ for $j = 1, 8$; $\Delta_{55} \triangleright a_{1,1}c_{s,j}^2$, $\Delta_{56} \triangleright a_{1,2}c_{s,j}^2$ for $j = 2, 3, 9, 11$; $\Delta_{57} \triangleright a_{1,1}c_{s,j}^2$, $\Delta_{58} \triangleright a_{1,2}c_{s,j}^2$, $\Delta_{59} \triangleright a_{1,3}c_{s,j}^2$ for $j = 4, 5, 6, 10, 12, 13, 14$; $\Delta_{57} \triangleright a_{1,1}c_{s,37}^2$ for $s = 3$, $\Delta_{57} \triangleright a_{1,1}c_{s,43}^2$ for $s > 3$; $\Delta_{58} \triangleright a_{1,2}c_{s,37}^2$ for $s = 3$, $\Delta_{58} \triangleright a_{1,2}c_{s,43}^2$ for $s > 3$ $\Delta_{59} \triangleright a_{1,3}c_{s,37}^2$ for $s = 3$, $\Delta_{59} \triangleright a_{1,3}c_{s,43}^2$ for $j > 3$; $\Delta_{60} \triangleright z_1b_{s,j}^2$ for $j = 4, 6, 16, 25$; $\Delta_{61} \triangleright z_1b_{s,j}^2$ for $j = 5, 7, 10, 17, 20, 24, 26, 29$; $\Delta_{62} \triangleright z_1b_{s,j}^2$,

$\Delta_{63} \triangleright z_2 b_{s,j}^2$ for $j = 8, 9, 11, 12, 18, 21, 22, 23, 27, 28, 30, 31, 32, 33, 34, 35$; $\Delta_{64} \triangleright a_{1,2} c_{s,1}^2$; $\Delta_{65} \triangleright a_{1,3} c_{s,2}^2$; $\Delta_{66} \triangleright a_{1,3} c_{s,j}^2$ for $j = 3, 11$; $\Delta_{67} \triangleright a_{1,4} c_{s,4}^2$; $\Delta_{68} \triangleright z_1 b_{s,j}^2$ for $j = 5, 12$; $\Delta_{69} \triangleright z_1 b_{s,j}^2$ for $j = 6, 13, 14$; $\Delta_{69} \triangleright z_1 b_{s,37}^2$ for $s = 3$, $\Delta_{69} \triangleright z_1 b_{s,43}^2$ for $s > 3$; $\Delta_{70} \triangleright z_2 b_{s,6}^2$; $\Delta_{71} \triangleright z_2 b_{s,7}^2$; $\Delta_{72} \triangleright z_3 b_{s,9}^2$; $\Delta_{73} \triangleright z_3 b_{s,12}^2$; $\Delta_{74} \triangleright z_2 b_{s,25}^2$; $\Delta_{75} \triangleright z_2 b_{s,26}^2$; $\Delta_{76} \triangleright z_3 b_{s,27}^2$; $\Delta_{77} \triangleright z_3 b_{s,28}^2$; $\Delta_{78} \triangleright z_3 b_{s,25}^2$; $\Delta_{79} \triangleright z_4 b_{s,26}^2$; $\Delta_{80} \triangleright z_4 b_{s,27}^2$; $\Delta_{81} \triangleright z_4 b_{s,28}^2$; $\Delta_{82} \triangleright z_3 b_{s,22}^2$; $\Delta_{83} \triangleright z_2 b_{s,24}^2$; $\Delta_{84} \triangleright z_4 b_{s,25}^2$; $\Delta_{85} \triangleright z_3 b_{s,26}^2$; $\Delta_{86} \triangleright z_4 b_{s,31}^2$; $\Delta_{87} \triangleright z_4 b_{s,33}^2$; $\Delta_{88} \triangleright z_3 b_{s,31}^2$; $\Delta_{89} \triangleright z_3 b_{s,33}^2$; $\Delta_{90} \triangleright z_3 b_{s,34}^2$, for $s = 3$, $\Delta_{90} \triangleright z_3 b_{s,j}^2$ for $j = 34, 35$ and $s > 3$; $\Delta_{91} \triangleright z_4 b_{s,34}^2$ for $s = 3$, $\Delta_{91} \triangleright z_4 b_{s,j}^2$ for $j = 34, 35$ and $s > 3$; $\Delta_{92} \triangleright z_3 b_{s,30}^2$, $\Delta_{93} \triangleright z_3 b_{s,32}^2$ for $s > 3$.

The proposition is completely proved. □

Now, we prove the $\rho_3(s)$ elements listed in Theorem 5.2 are linearly independent.

Proposition 5.14. *The elements $[d_{2,j}], 1 \leqslant j \leqslant 9$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_7$.*

Proof. Suppose that there is a linear relation

$$\sum_{1 \leqslant j \leqslant 9} \gamma_j [d_{2,j}] = 0, \tag{5.14.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Under the homomorphisms $f_i, 1 \leqslant i \leqslant 5$, the images of (5.14.1) respectively are

$$\begin{aligned}
&\gamma_{\{1,2,5,6,7,8,9\}}[1,2,4] + \gamma_3[3,1,3] + \gamma_4[3,3,1] = 0,\\
&\gamma_{\{2,3,4,5,7,8,9\}}[1,2,4] + \gamma_1[3,1,3] + \gamma_6[3,3,1] = 0,\\
&\gamma_{\{1,3,4,6,7,8,9\}}[1,2,4] + \gamma_2[3,1,3] + \gamma_5[3,3,1] = 0,\\
&\gamma_{\{2,4,5,6,7,9\}}[1,2,4] + \gamma_{\{1,3\}}[1,3,3] + \gamma_8[3,3,1] = 0,\\
&\gamma_{\{1,3,5,6,8,9\}}[1,2,4] + \gamma_{\{2,4\}}[1,3,3] + \gamma_7[3,3,1] = 0.
\end{aligned}$$

From these above equalities, we get $\gamma_j = 0, 1 \leqslant j \leqslant 9$. The proposition is proved. □

Proposition 5.15. *The elements $[d_{3,j}], 1 \leqslant j \leqslant 37$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{15}$.*

Proof. Suppose that there is a linear relation

$$\sum_{1 \leqslant j \leqslant 37} \gamma_j [d_{3,j}] = 0, \tag{5.15.1}$$

with $\gamma_j \in \mathbb{F}_2$. We need to prove that $\gamma_j = 0$ for all j.

Under the homomorphisms $f_i, i = 1, 2, \ldots, 6$, the images of (5.15.1) respectively are

$$\begin{aligned}
&a_1[1,2,12] + \gamma_{\{3,32\}}[7,1,7] + \gamma_{\{4,33\}}[7,7,1]\\
&\quad + \gamma_9[3,5,7] + \gamma_{10}[3,7,5] + \gamma_{\{23,36\}}[7,3,5] = 0,
\end{aligned}$$

$$
\begin{aligned}
&a_2[1,2,12]+\gamma_{\{1,16\}}[7,1,7]+\gamma_{\{6,34\}}[7,7,1]\\
&\quad+\gamma_{13}[3,5,7]+\gamma_{11}[3,7,5]+\gamma_{\{22,29\}}[7,3,5]=0,\\
&a_3[1,2,12]+\gamma_{\{2,17\}}[7,1,7]+\gamma_{\{5,18\}}[7,7,1]\\
&\quad+\gamma_{14}[3,5,7]+\gamma_{15}[3,7,5]+\gamma_{\{21,30,31\}}[7,3,5]=0,\\
&a_4[1,2,12]+\gamma_{\{1,3,9,13\}}[1,7,7]+\gamma_{\{8,35\}}[7,7,1]\\
&\quad+\gamma_{\{16,32\}}[3,5,7]+\gamma_{\{25,28,29,36\}}[3,7,5]+\gamma_{12}[7,3,5]=0,\\
&a_5[1,2,12]+\gamma_{\{2,4,10,14\}}[1,7,7]+\gamma_{\{7,20\}}[7,7,1]\\
&\quad+\gamma_{\{17,33\}}[3,5,7]+\gamma_{\{24,30\}}[3,7,5]+\gamma_{19}[7,3,5]=0,\\
&a_6[1,2,12]+\gamma_{\{5,6,11,15\}}[1,7,7]+\gamma_{\{7,8,12,19\}}[7,1,7]\\
&\quad+\gamma_{\{26,27,28,31\}}[3,5,7]+\gamma_{\{18,34\}}[3,7,5]+\gamma_{\{20,35\}}[7,3,5]=0.
\end{aligned}
$$

where

$$
\begin{aligned}
&a_1=\gamma+\gamma_{\{3,4,9,10,23,32,33,36\}},\ a_3=\gamma+\gamma_{\{2,5,14,15,17,18,21,30,31\}},\\
&a_2=\gamma+\gamma_{\{1,6,11,13,16,22,29,34\}},\ a_4=\gamma+\gamma_{\{1,3,8,9,12,13,16,25,28,29,32,35,36\}},\\
&a_5=\gamma+\gamma_{\{2,4,7,10,14,17,19,20,24,30,33\}},\\
&a_6=\gamma+\gamma_{\{5,6,7,8,11,12,15,18,19,20,26,27,28,31,34,35\}}.
\end{aligned}
$$

with $\gamma=\sum_{j=1}^{37}\gamma_j$.

Computing directly from the above equalities, we get

$$
\begin{cases}
a_i=0,\ i=1,2,\dots,6,\ \gamma_j=0, j=9,10,\dots,15,19,\\
\gamma_{\{3,32\}}=\gamma_{\{4,33\}}=\gamma_{\{23,36\}}=\gamma_{\{1,16\}}=\gamma_{\{6,34\}}=0,\\
\gamma_{\{22,29\}}=\gamma_{\{2,17\}}=\gamma_{\{5,18\}}=\gamma_{\{21,30,31\}}=0,\\
\gamma_{\{1,3\}}=\gamma_{\{8,35\}}=\gamma_{\{16,32\}}=\gamma_{\{25,28,29,36\}}=0,\\
\gamma_{\{2,4\}}=\gamma_{\{7,20\}}=\gamma_{\{17,33\}}=\gamma_{\{24,30\}}=\gamma_{\{5,6\}}=0,\\
\gamma_{\{7,8\}}=\gamma_{\{26,27,28,31\}}=\gamma_{\{18,34\}}=\gamma_{\{20,35\}}=0.
\end{cases}
\tag{5.15.2}
$$

With the aid of (5.15.2), the homomorphisms g_1, g_2 send (5.15.1) to

$$
\begin{aligned}
&\gamma_{\{25,27,28\}}[1,2,12]+\gamma_{27}[7,7,1]+\gamma_{25}[3,7,5]+\gamma_{28}[7,3,5]=0,\\
&\gamma_{\{21,23,26\}}[1,2,12]+\gamma_{26}[7,7,1]+\gamma_{\{21,23,31\}}[3,7,5]+\gamma_{31}[7,3,5]=0.
\end{aligned}
$$

Computing directly from these equalities gives

$$
\begin{cases}
\gamma_j=0,\ j=25,26,27,28,31,\\
\gamma_{\{21,23\}}=0.
\end{cases}
\tag{5.15.3}
$$

With the aid of (5.15.2) and (5.15.3), the homomorphisms g_3, g_4 send (5.15.1) respectively to

$$
\begin{aligned}
&\gamma_{\{21,22,24\}}[1,2,12]+\gamma_{24}[7,7,1]+\gamma_{\{21,22,24\}}[3,7,5]+\gamma_{24}[7,3,5]=0,\\
&\gamma_{\{21,22,24\}}[1,2,12]+\gamma_{\{21,22,23\}}[7,7,1]+\gamma_{22}[3,7,5]+\gamma_{\{22,23,24\}}[7,3,5]=0.
\end{aligned}
$$

These equalities imply

$$\gamma_j = 0,\ j = 21, 22, 23, 24. \tag{5.15.4}$$

Combining (5.15.2), (5.15.3) and (5.15.4), we get

$$\begin{cases} \gamma_j = 0,\ j \neq 1, 2, \ldots, 8, 16, 17, 18, 20, 32, 33, 34, 35, \\ \gamma_1 = \gamma_3 = \gamma_{16} = \gamma_{32}, \quad \gamma_2 = \gamma_4 = \gamma_{17} = \gamma_{33}, \\ \gamma_5 = \gamma_6 = \gamma_{18} = \gamma_{34}, \quad \gamma_7 = \gamma_8 = \gamma_{20} = \gamma_{35}. \end{cases} \tag{5.15.5}$$

Substituting (5.15.5) into the relation (5.15.1), we obtain

$$\gamma_1[\theta_1] + \gamma_2[\theta_2] + \gamma_5[\theta_3] + \gamma_7[\theta_4] = 0, \tag{5.15.6}$$

where

$$\theta_1 = d_{3,1} + d_{3,3} + d_{3,16} + d_{3,32}, \quad \theta_2 = d_{3,2} + d_{3,4} + d_{3,17} + d_{3,33},$$
$$\theta_3 = d_{3,5} + d_{3,6} + d_{3,18} + d_{3,34}, \quad \theta_4 = d_{3,7} + d_{3,8} + d_{3,20} + d_{3,35}.$$

Now, we prove that $\gamma_1 = \gamma_2 = \gamma_5 = \gamma_7 = 0$. The proof is divided into 4 steps.

Step 1. Consider the homomorphisms $\varphi_i, i = 1, 2, 3, 4$, defined in Section 2. Under the homomorphism φ_4, the image of (5.15.6) is

$$\gamma_1[\theta_1] + \gamma_2[\theta_2] + \gamma_5[\theta_3] + \gamma_7[\theta_4] + \gamma_7[\theta_3] = 0. \tag{5.15.7}$$

Combining (3.15.6) and (3.15.7), we get

$$\gamma_7[\theta_3] = 0. \tag{5.15.8}$$

If $[\theta_3] = 0$ then the polynomial θ_3 is hit. Hence, we have

$$\theta_3 = Sq^1(A) + Sq^2(B) + Sq^4(C),$$

for some polynomials $A \in (R_4)_{14}, B \in (R_4)_{13}, C \in (R_4)_{11}$. Let $Sq^2Sq^2Sq^2$ act on the both sides of this equality. We get

$$Sq^2Sq^2Sq^2(\theta_3) = Sq^2Sq^2Sq^2Sq^4(C),$$

By a direct calculation, we see that the monomial $(8, 7, 4, 2)$ is a term of $Sq^2Sq^2Sq^2(\theta_3)$. This monomial is not a term of $Sq^2Sq^2Sq^2Sq^4(y)$ for all monomials $y \in (R_4)_{11}$. Hence,

$$Sq^2Sq^2Sq^2(\theta_3) \neq Sq^2Sq^2Sq^2Sq^4(C),$$

for all $C \in (R_4)_{11}$ and we have a contradiction. So, $[\theta_3] \neq 0$ and the relation (5.15.8) implies $\gamma_7 = 0$.

Step 2. Since $\gamma_7 = 0$, the homomorphism φ_1 sends (5.15.6) to

$$\gamma_1[\theta_1] + \gamma_2[\theta_2] + \gamma_5[\theta_4] = 0. \tag{5.15.9}$$

Using the relation (5.15.9) and by the same argument as given in Step 1, we get $\gamma_5 = 0$.

Step 3. Since $\gamma_7 = \gamma_5 = 0$, the homomorphism φ_2 sends (5.15.6) to

$$\gamma_1[\theta_1] + \gamma_2[\theta_3] = 0. \tag{5.15.10}$$

Using the relation (5.15.10) and by the same argument as given in Step 2, we obtain $\gamma_2 = 0$.

Step 4. Since $\gamma_7 = \gamma_5 = \gamma_2 = 0$, the homomorphism φ_3 sends (5.15.6) to

$$\gamma_1[\theta_2] = 0.$$

Using this relation and by the same argument as given in Step 3, we obtain $\gamma_1 = 0$.

The proposition is completely proved. □

Proposition 5.16. *The elements* $[d_{4,j}], 1 \leqslant j \leqslant 47$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{31}$.

Proof. Suppose that there is a linear relation

$$\sum_{1\leqslant j\leqslant 47} \gamma_j[d_{4,j}] = 0, \tag{5.16.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_i, i = 1, 2, 3, 4, 5, 6$, to the linear relation (5.16.1), we get

$$\begin{aligned}
&a_1[1,2,28] + 1\gamma_3[15,1,15] + \gamma_4[15,15,1] + \gamma_9[3,13,15]\\
&\quad + \gamma_{10}[3,15,13] + \gamma_{23}[15,3,13] + \gamma_{\{32,38\}}[7,11,13] = 0,\\
&a_2[1,2,28] + \gamma_{\{1,16\}}[15,1,15] + \gamma_6[15,15,1] + \gamma_{13}[3,13,15]\\
&\quad + \gamma_{11}[3,15,13] + \gamma_{\{22,29,45,46\}}[15,3,13] + \gamma_{\{33,44\}}[7,11,13] = 0,\\
&a_3[1,2,28] + \gamma_{\{2,17\}}[15,1,15] + \gamma_{\{5,18\}}[15,15,1] + \gamma_{14}[3,13,15]\\
&\quad + \gamma_{15}[3,15,13] + \gamma_{\{21,30,31,40,41,47\}}[15,3,13] + \gamma_{35}[7,11,13] = 0,\\
&a_4[1,2,28] + \gamma_{\{1,3,9,13\}}[1,15,15] + \gamma_8[15,15,1] + \gamma_{16}[3,13,15]\\
&\quad + \gamma_{\{25,28,29,38,39,44,46\}}[3,15,13] + \gamma_{12}[15,3,13] + \gamma_{\{34,45\}}[7,11,13] = 0,\\
&a_5[1,2,28] + \gamma_{\{2,4,10,14\}}[1,15,15] + \gamma_{\{7,20\}}[15,15,1] + \gamma_{17}[3,13,15]\\
&\quad + \gamma_{\{24,30,42,47\}}[3,15,13] + \gamma_{19}[15,3,13] + \gamma_{36}[7,11,13] = 0,\\
&a_6[1,2,28] + \gamma_{\{5,6,11,15\}}[1,15,15] + \gamma_{\{7,8,12,19\}}[15,1,15] + \gamma_{18}[3,15,13]\\
&\quad + \gamma_{\{26,27,28,31\}}[3,13,15] + \gamma_{20}[15,3,13] + \gamma_{37}[7,11,13] = 0.
\end{aligned}$$

where

$$\begin{aligned}
a_1 &= \gamma + \gamma_{\{3,4,9,10,23,32,38\}},\\
a_2 &= \gamma + \gamma_{\{1,6,11,13,16,22,29,33,44,45,46\}},\\
a_3 &= \gamma + \gamma_{\{2,5,14,15,17,18,21,30,31,35,40,41,47\}},\\
a_4 &= \gamma + \gamma_{\{1,3,8,9,12,13,16,25,28,29,34,38,39,44,45,46\}},\\
a_5 &= \gamma + \gamma_{\{2,4,7,10,14,17,19,20,24,30,36,42,47\}},\\
a_6 &= \gamma + \gamma_{\{5,6,7,8,11,12,15,18,19,20,26,27,28,31,37\}},
\end{aligned}$$

with $\gamma = \sum_{1\leqslant j\leqslant 47} \gamma_j$.

Computing directly from the above equalities, we get

$$\begin{cases} a_i = 0,\ i = 1, 2 \ldots, 6, \\ \gamma_j = 0,\ j = 1, 2, \ldots, 20, 23, 35, 36, 37, \\ \gamma_{\{32,38\}} = \gamma_{\{22,29,34,46\}} = \gamma_{\{33,44\}} = 0, \\ \gamma_{\{21,30,31,40,41,47\}} = \gamma_{\{25,28,29,32,33,39,46\}} = 0, \\ \gamma_{\{34,45\}} = \gamma_{\{24,30,42,47\}} = \gamma_{\{26,27,28,31\}} = 0. \end{cases} \tag{5.16.2}$$

With the aid of (5.16.2), the homomorphisms g_1, g_2 send (5.16.1) respectively to

$$\begin{aligned} &\gamma_{\{22,25,27,28,29,39,46\}}[1,2,28] + \gamma_{27}[15,15,1] \\ &\quad + \gamma_{\{22,25,29,46\}}[3,15,13] + \gamma_{28}[15,3,13] + \gamma_{39}[7,11,13] = 0, \\ &\gamma_{\{21,24,26,30,32,41,42,47\}}[1,2,28] + \gamma_{26}[15,15,1] \\ &\quad + \gamma_{\{21,24,30,31,32,40,41,42,47\}}[3,15,13] + \gamma_{31}[15,3,13] + \gamma_{40}[7,11,13] = 0. \end{aligned}$$

Computing directly from the above equalities and (5.16.2), we have

$$\begin{cases} \gamma_j = 0,\ j = 26, 27, 28, 31, 39, 40, \\ \gamma_{\{22,25,29,46\}} = \gamma_{\{21,24,30,32,41,42,47\}} = 0. \end{cases} \tag{5.16.3}$$

With the aid of (5.16.2) and (5.16.3), the homomorphisms g_3, g_4 send (5.16.1) respectively to

$$\begin{aligned} &\gamma_{\{21,22,24,25,33,34\}}[1,2,28] + \gamma_{\{24,25,34\}}[15,15,1] + \gamma_{41}[7,11,13] \\ &\quad + \gamma_{\{21,29,30,33,34,41,42,46,47\}}[3,15,13] + \gamma_{\{22,29,30,34,42,46,47\}}[15,3,13] = 0, \\ &\gamma_{\{21,22,24,25,33,34\}}[1,2,28] + \gamma_{\{21,22,32,33\}}[15,15,1] + \gamma_{42}[7,11,13] \\ &\quad + \gamma_{\{24,29,30,33,34,41,42,46,47\}}[3,15,13] + \gamma_{\{25,29,30,32,33,41,46,47\}}[15,3,13] = 0. \end{aligned}$$

From these equalities, it implies

$$\begin{cases} \gamma_{41} = \gamma_{42} = \gamma_{\{21,22,24,25,33,34\}} = \gamma_{\{21,29,30,33,34,46,47\}} = 0, \\ \gamma_{\{24,25,34\}} = \gamma_{\{22,29,30,34,46,47\}} = \gamma_{\{21,22,24,25,33,34\}} = 0, \\ \gamma_{\{21,22,32,33\}} = \gamma_{\{24,29,30,33,34,46,47\}} = \gamma_{\{25,29,30,32,33,46,47\}} = 0. \end{cases} \tag{5.16.4}$$

Combining (5.16.2), (5.16.3) and (5.16.4), we get

$$\begin{cases} \gamma_j = 0, \text{ for } j \neq 22, 25, 29, 30, 33, 34, 44, 45, 46, 47, \\ \gamma_{22} = \gamma_{29} = \gamma_{44},\ \ \gamma_{25} = \gamma_{34} = \gamma_{45}, \\ \gamma_{\{30,47\}} = 0,\ \ \gamma_{\{22,25,29,46\}} = 0. \end{cases}$$

Substituting the above equalities into the linear relation (5.16.1), we have

$$\gamma_{22}[\theta_1] + \gamma_{25}[\theta_2] + \gamma_{46}[\theta_3] + \gamma_{47}[\theta_4] = 0, \tag{5.16.5}$$

where

$$\theta_1 = d_{4,22} + d_{4,29} + d_{4,33} + d_{4,44},$$
$$\theta_2 = d_{4,25} + d_{4,29} + d_{4,34} + d_{4,45},$$
$$\theta_3 = d_{4,29} + d_{4,46}, \quad \theta_4 = d_{4,30} + d_{4,47}.$$

Now, we prove $\gamma_{22} = \gamma_{25} = \gamma_{46} = \gamma_{47} = 0$. The proof is divided into 4 steps.

Step 1. The homomorphism φ_4 sends (5.16.5) to

$$\gamma_{22}[\theta_1] + \gamma_{25}[\theta_2] + \gamma_{46}[\theta_3] + \gamma_{47}[\theta_4] + \gamma_{25}[\theta_1] = 0. \tag{5.16.6}$$

Combining (5.16.5) and (5.16.6) gives

$$\gamma_{25}[\theta_1] = 0. \tag{5.16.7}$$

By an analogous argument as given in the proof of Proposition 5.15, we can show that the polynomial θ_1 is non-hit. So, $[\theta_1] \neq 0$ and $\gamma_{25} = 0$.

Step 2. Applying the homomorphism φ_1 to (5.16.5), we obtain

$$\gamma_{22}[\theta_2] + \gamma_{46}[\theta_3] + \gamma_{47}[\theta_4] = 0. \tag{5.16.8}$$

Using (5.16.8) and by a same argument as given in Step 1, we get $\gamma_{22} = 0$.

Step 3. Under the homomorphism φ_2, the image of (5.16.5) is

$$\gamma_{46}[\theta_2] + \gamma_{47}[\theta_4] = 0. \tag{5.16.9}$$

Using (5.16.9) and by a same argument as given in Step 3, we obtain $\gamma_{46} = 0$.

Step 4. Since $\gamma_{22} = \gamma_{25} = \gamma_{46} = 0$, the homomorphism φ_3 sends (5.16.5) to

$$\gamma_{47}[\theta_3] = 0.$$

From this equality and by a same argument as given in Step 3, we get $\gamma_{47} = 0$. The proposition is proved. □

Proposition 5.17. *For $s \geqslant 5$, the elements $[d_{s,j}], 1 \leqslant j \leqslant 43$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}-1}$.*

Proof. Suppose that there is a linear relation

$$\sum_{1 \leqslant j \leqslant 43} \gamma_j [d_{s,j}] = 0, \tag{5.17.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Under the homomorphisms $f_i, i = 1, 2, 3, 4, 5, 6$, the images of (5.17.1) respectively are

$$\begin{aligned}
&a_1u_{s,7} + \gamma_3u_{s,9} + \gamma_4u_{s,10} + \gamma_9u_{s,11}\\
&\quad + \gamma_{10}u_{s,12} + \gamma_{23}u_{s,13} + \gamma_{\{32,38\}}u_{s,14} = 0,\\
&a_2u_{s,7} + \gamma_{\{1,16\}}u_{s,9} + \gamma_6u_{s,10} + \gamma_{13}u_{s,11}\\
&\quad + \gamma_{11}u_{s,12} + \gamma_{\{22,29\}}u_{s,13} + \gamma_{33}u_{s,14} = 0,\\
&a_3u_{s,7} + \gamma_{\{2,17\}}u_{s,9} + \gamma_{\{5,18\}}u_{s,10} + \gamma_{14}u_{s,11}\\
&\quad + \gamma_{15}u_{s,12} + \gamma_{\{21,30,31,40,41\}}u_{s,13} + \gamma_{35}u_{s,14} = 0,\\
&a_4u_{s,7} + \gamma_{\{1,3,9,13\}}u_{s,8} + \gamma_8u_{s,10} + \gamma_{16}u_{s,11}\\
&\quad + \gamma_{\{25,28,29,38,39\}}u_{s,12} + \gamma_{12}u_{s,13} + \gamma_{34}u_{s,14} = 0,\\
&a_5u_{s,7} + \gamma_{\{2,4,10,14\}}u_{s,8} + \gamma_{\{7,20\}}u_{s,10} + \gamma_{17}u_{s,11}\\
&\quad + \gamma_{\{24,30,42\}}u_{s,12} + \gamma_{19}u_{s,13} + \gamma_{36}u_{s,14} = 0,\\
&a_6u_{s,7} + \gamma_{\{5,6,11,15\}}u_{s,8} + \gamma_{\{7,8,12,19\}}u_{s,9} + \gamma_{18}u_{s,12}\\
&\quad + \gamma_{\{26,27,28,31\}}u_{s,11} + \gamma_{20}u_{s,13} + \gamma_{37}u_{s,14} = 0.
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma + \gamma_{\{3,4,9,10,23,32,38\}}, \quad a_2 = \gamma + \gamma_{\{1,6,11,13,16,22,29,33\}},\\
&a_3 = \gamma + \gamma_{\{2,5,14,15,17,18,21,30,31,35,40,41\}},\\
&a_4 = \gamma + \gamma_{\{1,3,8,9,12,13,16,25,28,29,34,38,39\}},\\
&a_5 = \gamma + \gamma_{\{2,4,7,10,14,17,19,20,24,30,36,42\}},\\
&a_6 = \gamma + \gamma_{\{5,6,7,8,11,12,15,18,19,20,26,27,28,31,37\}},
\end{aligned}$$

with $\gamma = \sum_{1\leqslant j\leqslant 43}\gamma_j$.

Computing directly from the above equalities gives

$$\begin{cases}
a_i = 0, \ i = 1, 2, 3, 4, 5, 6,\\
\gamma_j = 0, \ j = 1, 2, \ldots, 20, 23, 33, \ldots, 37,\\
\gamma_{\{32,38\}} = \gamma_{\{22,29\}} = \gamma_{\{21,30,31,40,41\}} = 0,\\
\gamma_{\{22,25,28,32,39\}} = \gamma_{\{24,30,42\}} = \gamma_{\{26,27,28,31\}} = 0.
\end{cases} \tag{5.17.2}$$

With the aid of (5.17.2), the homomorphisms g_1, g_2 send (5.17.1) respectively to

$$\begin{aligned}
&\gamma_{\{25,27,28,39\}}u_{s,7} + \gamma_{27}u_{s,10} + \gamma_{25}u_{s,12} + \gamma_{28}u_{s,13} + \gamma_{39}u_{s,14} = 0,\\
&\gamma_{\{21,24,26,30,32,41,42\}}u_{s,7} + \gamma_{26}u_{s,10}\\
&\quad + \gamma_{\{21,24,30,31,32,40,41,42\}}u_{s,12} + \gamma_{31}u_{s,13} + \gamma_{40}u_{s,14} = 0.
\end{aligned}$$

From the above equalities, we obtain

$$\begin{cases}
\gamma_j = 0, \ j = 25, 26, 27, 28, 31, 39, 40,\\
\gamma_{\{21,24,30,32,41,42\}} = 0.
\end{cases} \tag{5.17.3}$$

With the aid of (5.17.2) and (5.17.3), the homomorphisms g_3, g_4 send (5.17.1) respectively to

$$\gamma_{\{21,22,24\}}u_{s,7} + \gamma_{24}u_{s,10} + \gamma_{\{21,22,30,41,42\}}u_{s,12} + \gamma_{\{30,42\}}u_{s,13} + \gamma_{41}u_{s,14} = 0,$$
$$\gamma_{\{21,22,24\}}u_{s,7} + \gamma_{\{21,22,32\}}u_{s,10} + \gamma_{\{22,24,30,41,42\}}u_{s,12} + \gamma_{\{22,30,32,41\}}u_{s,13} + \gamma_{42}u_{s,14} = 0.$$

From these equalities, it implies

$$\gamma_j = 0,\ j = 21, 22, 24, 30, 32, 41, 42. \tag{5.17.4}$$

Combining (5.17.2), (5.17.3) and (5.17.4), we get $\gamma_j = 0$ for all j. The proposition is proved. □

Remark 5.18. The $\mathbb{F}_2$-subspace of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{15}$ generated by the elements $[\theta_1], [\theta_2], [\theta_3], [\theta_4]$, which are defined as in the proof of Proposition 5.15, is an $GL_4(\mathbb{F}_2)$-submodule of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{15}$.

The $\mathbb{F}_2$-subspace of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{31}$ generated by $[\theta_1], [\theta_2], [\theta_3], [\theta_4]$, which are defined as in the proof of Proposition 5.16, is an $GL_4(\mathbb{F}_2)$-submodule of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{31}$.

6. The indecomposables of P_4 in degree $2^{s+t+1} + 2^{s+1} - 3$

6.1. The τ-sequence of the admissible monomials.

First of all, we determine the τ-sequence of an admissible monomial of degree $2^{s+t+1} + 2^{s+1} - 3$ for any positive integers s, t.

Lemma 6.1.1. *Let x be a monomial of degree $2^{s+t+1} + 2^{s+1} - 3$ in P_4 with s, t are the positive integers. If x is admissible then $\tau(x)$ is one of the following sequences*

$$(\underbrace{3; 3; \ldots; 3}_{s\ times}; \underbrace{1; 1; \ldots; 1}_{t+1\ times}), \quad (\underbrace{3; 3; \ldots; 3}_{s+1\ times}; \underbrace{2; 2; \ldots; 2}_{t-1\ times}).$$

From the proof of Proposition 5.3 and Lemmas 2.6, 4.4, 4.5, we easily obtain the following

Lemma 6.1.2. *Let x be a monomial of degree $2^s - 1$ in P_4 with $s \geqslant 2$. If x is inadmissible then there is a strictly inadmissible matrix Δ such that $\Delta \triangleright x$.*

Proof of Proposition 6.1.1. Observe that $z = (2^{s+t+1} - 1, 2^s - 1, 2^s - 1, 0)$ is the minimal spike of degree $2^{s+t+1} + 2^{s+1} - 3$ in P_4 and

$$\tau(z) = (\underbrace{3; 3; \ldots; 3}_{s\ \text{times}}; \underbrace{1; 1; \ldots; 1}_{t+1\ \text{times}}).$$

Since x is admissible and $2^{s+t+1} + 2^{s+1} - 3$ is odd, using Theorem 2.11, we obtain $\tau_1(x) = 3$. Using Theorem 2.13 and Proposition 2.7, we get $\tau_i(x) = 3$ for $i = 1, 2, \ldots, s$.

Suppose that $M = (\varepsilon_{ij}(x)), i \geqslant 1, 1 \leqslant j \leqslant 4$, is the matrix associated with x. We set $M' = (\varepsilon_{ij}(x)), i > s, 1 \leqslant j \leqslant 4$ and denote by x' the monomial corresponding to M'. Then $\tau_i(x') = \tau_{i+s}(x), i \geqslant 1$. We have

$$\begin{aligned} 2^{s+t+1} + 2^{s+1} - 3 = \deg x &= \sum_{i\geqslant 1} 2^{i-1}\tau_i(x) \\ &= 3(2^s - 1) + 2^s \sum_{j\geqslant 1} 2^{j-1}\tau_{j+s}(x) \\ &= 2^{s+1} + 2^s - 3 + 2^s \deg(x'). \end{aligned}$$

From this, it implies $\deg(x') = 2^{t+1} - 1$. Since x is admissible, using Lemma 6.1.2 and Theorem 2.5, we see that x' is also admissible. By Lemmas 5.6, 5.7, 5.4, $\tau(x')$ is one of the following sequences

$$(\underbrace{1; 1; \ldots; 1}_{t+1 \text{ times}}), \quad (3; \underbrace{2; 2; \ldots; 2}_{t-1 \text{ times}}).$$

The lemma is proved. □

6.2. **The case $t = 1$.**

According to Kameko [8], $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+2}+2^{s+1}-3} = 7$ with a basis given by the classes

$$\begin{aligned} &w_{1,s,1} = [2^s - 1, 2^s - 1, 2^{s+2} - 1], && w_{1,s,2} = [2^s - 1, 2^{s+2} - 1, 2^s - 1], \\ &w_{1,s,3} = [2^{s+2} - 1, 2^s - 1, 2^s - 1], && w_{1,s,4} = [2^s - 1, 2^{s+1} - 1, 3.2^s - 1], \\ &w_{1,s,5} = [2^{s+1} - 1, 2^s - 1, 3.2^s - 1], && w_{1,s,6} = [2^{s+1} - 1, 3.2^s - 1, 2^s - 1], \\ &w_{1,s,7} = [2^{s+1} - 1, 2^{s+1} - 1, 2^{s+1} - 1]. \end{aligned}$$

Hence, we have

Proposition 6.2.1. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+2}+2^{s+1}-3}$ is an $\mathbb{F}_2$-vector space of dimension 28 with a basis consisting of all the classes represented by the monomials $a_{1,s,j}, 1 \leqslant j \leqslant 28$, which are determined as follows:*

1. $(0, 2^{s+1} - 1, 2^{s+1} - 1, 2^{s+1} - 1)$,
2. $(2^{s+1} - 1, 0, 2^{s+1} - 1, 2^{s+1} - 1)$,
3. $(2^{s+1} - 1, 2^{s+1} - 1, 0, 2^{s+1} - 1)$,
4. $(2^{s+1} - 1, 2^{s+1} - 1, 2^{s+1} - 1, 0)$,
5. $(0, 2^s - 1, 2^s - 1, 2^{s+2} - 1)$,
6. $(0, 2^s - 1, 2^{s+2} - 1, 2^s - 1)$,
7. $(0, 2^{s+2} - 1, 2^s - 1, 2^s - 1)$,
8. $(2^s - 1, 0, 2^s - 1, 2^{s+2} - 1)$,
9. $(2^s - 1, 0, 2^{s+2} - 1, 2^s - 1)$,
10. $(2^s - 1, 2^s - 1, 0, 2^{s+2} - 1)$,
11. $(2^s - 1, 2^s - 1, 2^{s+2} - 1, 0)$,
12. $(2^s - 1, 2^{s+2} - 1, 0, 2^s - 1)$,
13. $(2^s - 1, 2^{s+2} - 1, 2^s - 1, 0)$,
14. $(2^{s+2} - 1, 0, 2^s - 1, 2^s - 1)$,
15. $(2^{s+2} - 1, 2^s - 1, 0, 2^s - 1)$,
16. $(2^{s+2} - 1, 2^s - 1, 2^s - 1, 0)$,
17. $(0, 2^s - 1, 2^{s+1} - 1, 3.2^s - 1)$,
18. $(0, 2^{s+1} - 1, 2^s - 1, 3.2^s - 1)$,
19. $(0, 2^{s+1} - 1, 3.2^s - 1, 2^s - 1)$,
20. $(2^s - 1, 0, 2^{s+1} - 1, 3.2^s - 1)$,
21. $(2^s - 1, 2^{s+1} - 1, 0, 3.2^s - 1)$,
22. $(2^s - 1, 2^{s+1} - 1, 3.2^s - 1, 0)$,
23. $(2^{s+1} - 1, 0, 2^s - 1, 3.2^s - 1)$,
24. $(2^{s+1} - 1, 0, 3.2^s - 1, 2^s - 1)$,
25. $(2^{s+1} - 1, 2^s - 1, 0, 3.2^s - 1)$,
26. $(2^{s+1} - 1, 2^s - 1, 3.2^s - 1, 0)$,
27. $(2^{s+1} - 1, 3.2^s - 1, 0, 2^s - 1)$,
28. $(2^{s+1} - 1, 3.2^s - 1, 2^s - 1, 0)$.

By Proposition 2.8, we need only to determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}-3}$.

Set $\mu_2(1) = 46, \mu_2(2) = 94$ and $\mu_2(s) = 105$ for $s \geqslant 3$. We have

Theorem 6.2.2. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}-3}$ is an $\mathbb{F}_2$-vector space of dimension $\mu_2(s) - 28$ with a basis consisting of all the classes represented by the monomials $a_{1,s,j}, 29 \leqslant j \leqslant \mu_2(s)$, which are determined as follows:*

For $s \geqslant 1$,

29. $(1, 2^{s+1} - 2, 2^{s+1} - 1, 2^{s+1} - 1)$,
30. $(1, 2^{s+1} - 1, 2^{s+1} - 2, 2^{s+1} - 1)$,
31. $(1, 2^{s+1} - 1, 2^{s+1} - 1, 2^{s+1} - 2)$,
32. $(2^{s+1} - 1, 1, 2^{s+1} - 2, 2^{s+1} - 1)$,
33. $(2^{s+1} - 1, 1, 2^{s+1} - 1, 2^{s+1} - 2)$,
34. $(2^{s+1} - 1, 2^{s+1} - 1, 1, 2^{s+1} - 2)$,
35. $(1, 2^s - 1, 2^s - 1, 2^{s+2} - 2)$,
36. $(1, 2^s - 1, 2^{s+2} - 2, 2^s - 1)$,
37. $(1, 2^{s+2} - 2, 2^s - 1, 2^s - 1)$,
38. $(1, 2^s - 1, 2^{s+1} - 2, 3.2^s - 1)$,
39. $(1, 2^{s+1} - 2, 2^s - 1, 3.2^s - 1)$,
40. $(1, 2^{s+1} - 2, 3.2^s - 1, 2^s - 1)$.

For $s = 1$,

41. $(1, 1, 3, 4)$,
42. $(1, 3, 1, 4)$,
43. $(1, 3, 4, 1)$,
44. $(3, 1, 1, 4)$,
45. $(3, 1, 4, 1)$,
46. $(3, 4, 1, 1)$.

For $s \geqslant 2$,

41. $(2^s - 1, 1, 2^{s+1} - 2, 3.2^s - 1)$,
42. $(2^s - 1, 1, 2^s - 1, 2^{s+2} - 2)$,
43. $(2^s - 1, 1, 2^{s+2} - 2, 2^s - 1)$,
44. $(2^s - 1, 2^s - 1, 1, 2^{s+2} - 2)$,
45. $(3, 2^{s+1} - 3, 2^{s+1} - 2, 2^{s+1} - 1)$,
46. $(3, 2^{s+1} - 3, 2^{s+1} - 1, 2^{s+1} - 2)$,
47. $(3, 2^{s+1} - 1, 2^{s+1} - 3, 2^{s+1} - 2)$,
48. $(2^{s+1} - 1, 3, 2^{s+1} - 3, 2^{s+1} - 2)$,
49. $(1, 2^s - 2, 2^s - 1, 2^{s+2} - 1)$,
50. $(1, 2^s - 2, 2^{s+2} - 1, 2^s - 1)$,
51. $(1, 2^s - 1, 2^s - 2, 2^{s+2} - 1)$,
52. $(1, 2^s - 1, 2^{s+2} - 1, 2^s - 2)$,
53. $(1, 2^{s+2} - 1, 2^s - 2, 2^s - 1)$,
54. $(1, 2^{s+2} - 1, 2^s - 1, 2^s - 2)$,
55. $(2^s - 1, 1, 2^s - 2, 2^{s+2} - 1)$,
56. $(2^s - 1, 1, 2^{s+2} - 1, 2^s - 2)$,
57. $(2^s - 1, 2^{s+2} - 1, 1, 2^s - 2)$,
58. $(2^{s+2} - 1, 1, 2^s - 2, 2^s - 1)$,
59. $(2^{s+2} - 1, 1, 2^s - 1, 2^s - 2)$,
60. $(2^{s+2} - 1, 2^s - 1, 1, 2^s - 2)$,
61. $(1, 2^s - 2, 2^{s+1} - 1, 3.2^s - 1)$,
62. $(1, 2^{s+1} - 1, 2^s - 2, 3.2^s - 1)$,
63. $(1, 2^{s+1} - 1, 3.2^s - 1, 2^s - 2)$,
64. $(2^{s+1} - 1, 1, 2^s - 2, 3.2^s - 1)$,
65. $(2^{s+1} - 1, 1, 3.2^s - 1, 2^s - 2)$,
66. $(2^{s+1} - 1, 3.2^s - 1, 1, 2^s - 2)$,
67. $(1, 2^s - 1, 2^{s+1} - 1, 3.2^s - 2)$,
68. $(1, 2^{s+1} - 1, 2^s - 1, 3.2^s - 2)$,
69. $(1, 2^{s+1} - 1, 3.2^s - 2, 2^s - 1)$,
70. $(2^s - 1, 1, 2^{s+1} - 1, 3.2^s - 2)$,
71. $(2^s - 1, 2^{s+1} - 1, 1, 3.2^s - 2)$,
72. $(2^{s+1} - 1, 1, 2^s - 1, 3.2^s - 2)$,
73. $(2^{s+1} - 1, 1, 3.2^s - 2, 2^s - 1)$,
74. $(2^{s+1} - 1, 2^s - 1, 1, 3.2^s - 2)$,
75. $(3, 2^s - 1, 2^{s+2} - 3, 2^s - 2)$,
76. $(3, 2^{s+2} - 3, 2^s - 2, 2^s - 1)$,
77. $(3, 2^{s+2} - 3, 2^s - 1, 2^s - 2)$,
78. $(3, 2^{s+1} - 3, 2^s - 2, 3.2^s - 1)$,
79. $(3, 2^{s+1} - 3, 3.2^s - 1, 2^s - 2)$,
80. $(3, 2^{s+1} - 1, 3.2^s - 3, 2^s - 2)$,
81. $(2^{s+1} - 1, 3, 3.2^s - 3, 2^s - 2)$,
82. $(3, 2^s - 1, 2^{s+1} - 3, 3.2^s - 2)$,
83. $(3, 2^{s+1} - 3, 2^s - 1, 3.2^s - 2)$,
84. $(3, 2^{s+1} - 3, 3.2^s - 2, 2^s - 1)$.

For $s = 2$,

85. $(3,3,3,12)$, 86. $(3,3,12,3)$, 87. $(7,9,2,3)$,
88. $(7,9,3,2)$, 89. $(3,3,4,11)$, 90. $(3,3,7,8)$,
91. $(3,7,3,8)$, 92. $(3,7,8,3)$, 93. $(7,3,3,8)$,
94. $(7,3,8,3)$.

For $s \geqslant 3$,

85. $(3, 2^s-3, 2^s-2, 2^{s+2}-1)$, 86. $(3, 2^s-3, 2^{s+2}-1, 2^s-2)$,
87. $(3, 2^{s+2}-1, 2^s-3, 2^s-2)$, 88. $(2^{s+2}-1, 3, 2^s-3, 2^s-2)$,
89. $(3, 2^s-3, 2^s-1, 2^{s+2}-2)$, 90. $(3, 2^s-3, 2^{s+2}-2, 2^s-1)$,
91. $(3, 2^s-1, 2^s-3, 2^{s+2}-2)$, 92. $(2^s-1, 3, 2^s-3, 2^{s+2}-2)$,
93. $(2^s-1, 3, 2^{s+2}-3, 2^s-2)$, 94. $(3, 2^s-3, 2^{s+1}-2, 3.2^s-1)$,
95. $(3, 2^s-3, 2^{s+1}-1, 3.2^s-2)$, 96. $(3, 2^{s+1}-1, 2^s-3, 3.2^s-2)$,
97. $(2^{s+1}-1, 3, 2^s-3, 3.2^s-2)$, 98. $(2^s-1, 3, 2^{s+1}-3, 3.2^s-2)$,
99. $(7, 2^{s+2}-5, 2^s-3, 2^s-2)$, 100. $(7, 2^{s+1}-5, 2^s-3, 3.2^s-2)$,
101. $(7, 2^{s+1}-5, 3.2^s-3, 2^s-2)$, 102. $(7, 2^{s+1}-5, 2^{s+1}-3, 2^{s+1}-2)$.

For $s = 3$,

103. $(7,7,7,24)$, 104. $(7,7,9,22)$, 105. $(7,7,25,6)$.

For $s \geqslant 4$,

103. $(7, 2^s-5, 2^s-3, 2^{s+2}-2)$, 104. $(7, 2^s-5, 2^{s+1}-3, 3.2^s-2)$,
105. $(7, 2^s-5, 2^{s+2}-3, 2^s-2)$.

This theorem is proved by combining some propositions.

Proposition 6.2.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}-3}$ is generated by the $\mu_2(s) - 28$ elements listed in Theorem 6.2.2.*

Lemma 6.2.4. *The following matrix is strictly inadmissible*

$$\begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

Proof. By a direct computation, we have

$$\begin{aligned}(7,8,3,3) &= Sq^1\big((7,5,5,3) + (7,5,3,5) + (7,3,5,5)\big)\\ &\quad + Sq^2\big((7,6,3,3) + (7,3,6,3) + (7,3,3,6)\big) + Sq^4\big((5,6,3,3)\\ &\quad + (5,3,6,3) + (5,3,3,6)\big) + (5,10,3,3) + (5,3,10,3)\\ &\quad + (5,3,3,10) + (7,3,8,3) + (7,3,3,8) \mod \mathcal{L}_4(3;3;1;1).\end{aligned}$$

The lemma is proved. □

Lemma 6.2.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices are

$$(15,19,5,6), (7,15,17,6), (15,7,17,6), (7,7,15,16),$$
$$(7,15,7,16), (7,15,16,7), (15,7,7,16), (15,7,16,7).$$

We prove the lemma for the matrices associated to the monomials

$$(15,19,5,6), (15,7,17,6), (15,7,7,16), (15,7,16,7).$$

The others can be obtained by a similar calculation. We have

$$\begin{aligned}
(15,19,5,6) &= Sq^1\big((15,15,7,7) + (15,15,5,9)\big) + Sq^2\big((15,15,7,6)\\
&\quad + (15,15,6,7)\big) + Sq^4(15,15,5,6) + Sq^8\big((11,15,5,6)\\
&\quad + (9,15,7,6) + (9,15,6,7) + (8,15,7,7)\big) + (11,23,5,6)\\
&\quad + (9,23,7,6) + (9,23,6,7) + (8,23,7,7) + (15,17,7,6)\\
&\quad + (15,17,6,7) + (15,16,7,7) \mod \mathcal{L}_4(3;3;3;1;1),\\
(15,7,17,6) &= Sq^1\big((15,5,15,9) + (15,7,15,7)\big) + Sq^2(15,7,15,6)\\
&\quad + (15,6,15,7) + Sq^4(15,5,15,6) + Sq^8\big((9,7,15,6)\\
&\quad + (11,5,15,6) + (8,7,15,7) + (9,6,15,7)\big) + (9,7,23,6)\\
&\quad + (11,5,23,6) + (8,7,23,7) + (9,6,23,7) + (15,5,19,6)\\
&\quad + (15,6,17,7) + (15,7,16,7) \mod \mathcal{L}_4(3;3;3;1;1),\\
(15,7,16,7) &= Sq^1\big((15,7,13,9) + (15,7,9,13) + (15,7,11,11)\\
&\quad + (15,5,11,13) + (15,5,13,11)\big) + Sq^2\big((15,7,14,7)\\
&\quad + (15,7,11,10) + (15,7,10,11) + (15,7,7,14) + (15,6,11,11)\big)\\
&\quad + Sq^4\big((15,5,14,7) + (15,5,7,14)\big) + Sq^8\big((9,7,14,7)\\
&\quad + (11,5,14,7) + (9,7,7,14) + (11,5,7,14)\big) + (9,7,22,7)\\
&\quad + (15,5,18,7) + (11,5,22,7) + (9,7,7,22) + (15,7,7,16)\\
&\quad + (15,5,7,18) + (11,5,7,22) \mod \mathcal{L}_4(3;3;3;1;1),\\
(15,7,7,16) &= Sq^1\big((15,7,7,15) + (15,3,11,15)\big) + Sq^2\big((15,7,6,15)\\
&\quad + (15,3,10,15)\big) + Sq^4\big((15,4,7,15) + (15,5,6,15)\big)\\
&\quad + Sq^8\big((8,7,7,15) + (11,4,7,15) + (9,7,6,15) + (11,5,6,15)\big)\\
&\quad + (8,7,7,23) + (11,4,7,23) + (15,4,7,19) + (9,7,6,23)\\
&\quad + (15,7,6,17) + (15,5,6,19) + (11,5,6,23) \mod \mathcal{L}_4(3;3;3;1;1).
\end{aligned}$$

The lemma follows. □

Combining Theorem 2.5, the lemmas in Sections 3, 5 and the Lemmas 6.1.1, 6.2.4, 6.2.5, we get Proposition 6.2.3.

Now, we show that $[a_{1,s,j}], 29 \leqslant j \leqslant \mu_2(s)$, are linearly independent in the space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}-3}$.

Proposition 6.2.6. *The elements* $[a_{1,1,j}],\ 29 \leqslant j \leqslant 46$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_9$.

Proof. Suppose there is a linear relation

$$\sum_{j=29}^{46} \gamma_j [a_{1,1,j}] = 0, \tag{6.2.6.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Consider the homomorphisms $f_i,\ i = 1, 2, 3, 4, 5, 6$. Under these homomorphisms, the images of the above relation respectively are

$$\begin{aligned}
&\gamma_{\{37,46\}}[7,1,1] + \gamma_{39}[3,1,5] + \gamma_{40}[3,5,1] + \gamma_{29}[3,3,3] = 0,\\
&\gamma_{\{36,45\}}[7,1,1] + \gamma_{\{32,38\}}[3,1,5] + \gamma_{43}[3,5,1] + \gamma_{30}[3,3,3] = 0,\\
&\gamma_{\{35,44\}}[7,1,1] + \gamma_{\{33,41\}}[3,1,5] + \gamma_{\{34,42\}}[3,5,1] + \gamma_{31}[3,3,3] = 0,\\
&\gamma_{\{36,37,40,43\}}[1,7,1]\\
&\quad + \gamma_{\{29,30,38,39\}}[1,3,5] + \gamma_{\{45,46\}}[3,5,1] + \gamma_{32}[3,3,3] = 0,\\
&\gamma_{\{35,37,39,42\}}[1,7,1]\\
&\quad + \gamma_{\{29,31,40,41\}}[1,3,5] + \gamma_{\{34,44,46\}}[3,5,1] + \gamma_{33}[3,3,3] = 0,\\
&\gamma_{\{35,36,38,41\}}[1,1,7]\\
&\quad + \gamma_{\{30,31,42,43\}}[1,3,5] + \gamma_{\{32,33,44,45\}}[3,1,5] + \gamma_{34}[3,3,3] = 0.
\end{aligned}$$

From the above equalities, we get

$$\begin{cases} \gamma_j = 0, \text{ for } j \neq 35, 36, 37, 44, 45, 46,\\ \gamma_{35} = \gamma_{36} = \gamma_{37} = \gamma_{44} = \gamma_{45} = \gamma_{46}. \end{cases} \tag{6.2.6.2}$$

Substituting (6.2.6.2) into the relation (6.2.6.1), we get

$$\gamma_{35}[\theta] = 0, \tag{6.2.6.3}$$

where $\theta = a_{1,1,35} + a_{1,1,36} + a_{1,1,37} + a_{1,1,44} + a_{1,1,45} + a_{1,1,46}$.

We prove $\gamma_{35} = 0$ by showing $[\theta] \neq 0$. Suppose $[\theta] = 0$. Then θ is hit and we get

$$\theta = Sq^1(A) + Sq^2(B) + Sq^4(C),$$

for some polynomials $A \in (R_4)_8, B \in (R_4)_7, C \in (R_4)_5$. Let $(Sq^2)^3 = Sq^2Sq^2Sq^2$ act on the both sides of this equality. It is easy to check that $(Sq^2)^3Sq^4(C) = 0$ for all $C \in (R_4)_5$. Since $(Sq^2)^3$ annihilates Sq^1 and Sq^2, the right hand side is sent to zero. On the other hand, a direct computation shows

$$(Sq^2)^3(\theta) = (1, 2, 4, 8) + \text{symmetries} \neq 0.$$

Hence, we have a contradiction. So, $[\theta] \neq 0$ and the relation (6.2.6.3) implies $\gamma_{35} = 0$. The proposition is proved. □

Proposition 6.2.7. *The elements* $[a_{1,2,j}],\ 29 \leqslant j \leqslant 94$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{21}$.

Proof. Suppose there is a linear relation

$$\sum_{j=29}^{94} \gamma_j [a_{1,2,j}] = 0, \tag{6.2.7.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms $f_i, i = 1, 2, \ldots, 6$, to the relation (6.2.7.1) and we get

$$\begin{aligned}
&\gamma_{49}[3,3,15] + \gamma_{50}[3,15,3] + \gamma_{37}[15,3,3] + \gamma_{61}[3,7,11] \\
&\quad + \gamma_{39}[7,3,11] + \gamma_{40}[7,11,3] + \gamma_{29}[7,7,7] = 0,\\
&\gamma_{51}[3,3,15] + \gamma_{53}[3,15,3] + \gamma_{\{36,86,94\}}[15,3,3] + \gamma_{62}[3,7,11] \\
&\quad + \gamma_{\{38,89\}}[7,3,11] + \gamma_{\{69,92\}}[7,11,3] + \gamma_{30}[7,7,7] = 0,\\
&\gamma_{52}[3,3,15] + \gamma_{54}[3,15,3] + \gamma_{\{35,85,93\}}[15,3,3] + \gamma_{63}[3,7,11] \\
&\quad + \gamma_{\{67,90\}}[7,3,11] + \gamma_{\{68,91\}}[7,11,3] + \gamma_{31}[7,7,7] = 0,\\
&\gamma_{55}[3,3,15] + \gamma_{\{43,76,84,86,92\}}[3,15,3] + \gamma_{58}[15,3,3] + \gamma_{64}[7,3,11] \\
&\quad + \gamma_{\{41,45,78,89\}}[3,7,11] + \gamma_{\{73,87,94\}}[7,11,3] + \gamma_{32}[7,7,7] = 0,\\
&\gamma_{56}[3,3,15] + \gamma_{\{42,77,83,85,91\}}[3,15,3] + \gamma_{\{46,70,79,90\}}[3,7,11] \\
&\quad + \gamma_{59}[15,3,3] + \gamma_{65}[7,3,11] + \gamma_{\{72,88,93\}}[7,11,3] + \gamma_{33}[7,7,7] = 0,\\
&\gamma_{\{44,75,82,85,86,89,90\}}[3,3,15] + \gamma_{57}[3,15,3] + \gamma_{60}[15,3,3] + \gamma_{66}[7,11,3] \\
&\quad + \gamma_{\{47,71,80,91,92\}}[3,7,11] + \gamma_{\{48,74,81,93,94\}}[7,3,11] + \gamma_{34}[7,7,7] = 0.
\end{aligned}$$

From the above equalities, we obtain

$$\begin{cases}
\gamma_j = 0,\ 29 \leqslant j \leqslant 34, \text{or } j = 37, 39, 40, \text{or } 49 \leqslant j \leqslant 66,\\
\gamma_{\{36,86,94\}} = \gamma_{\{38,89\}} = \gamma_{\{69,92\}} = \gamma_{\{35,85,93\}} = \gamma_{\{67,90\}} = 0,\\
\gamma_{\{68,91\}} = \gamma_{\{43,76,84,86,92\}} = \gamma_{\{41,45,78,89\}} = \gamma_{\{73,87,94\}} = 0,\\
\gamma_{\{42,77,83,85,91\}} = \gamma_{\{46,70,79,90\}} = \gamma_{\{72,88,93\}} = 0,\\
\gamma_{\{44,75,82,85,86,89,90\}} = \gamma_{\{47,71,80,91,92\}} = \gamma_{\{48,74,81,93,94\}} = 0.
\end{cases} \tag{6.2.7.2}$$

With the aid of (6.2.7.2), the homomorphisms g_1, g_2 send (6.2.7.1) respectively to

$$\begin{aligned}
&\gamma_{\{36,43,73,86,94\}}[3,15,3] + \gamma_{\{76,87\}}[15,3,3] + \gamma_{41}[3,7,11] \\
&\quad + \gamma_{78}[7,3,11] + \gamma_{\{73,84,87,94\}}[7,11,3] + \gamma_{45}[7,7,7] = 0,\\
&\gamma_{\{35,42,72,85,93\}}[3,15,3] + \gamma_{\{77,88\}}[15,3,3] + \gamma_{70}[3,7,11] \\
&\quad + \gamma_{79}[7,3,11] + \gamma_{\{72,83,88,93\}}[7,11,3] + \gamma_{46}[7,7,7] = 0.
\end{aligned}$$

Computing directly from the above equalities, we obtain

$$\begin{cases} \gamma_j = 0,\ j = 41, 45, 46, 70, 78, 79, \\ \gamma_{\{36,43,73,86,94\}} = \gamma_{\{76,87\}} = \gamma_{\{35,42,72,85,93\}} = 0, \\ \gamma_{\{77,88\}} = \gamma_{\{73,84,87,94\}} = \gamma_{\{72,83,88,93\}} = 0. \end{cases} \tag{6.2.7.3}$$

With the aid of (6.2.7.2) and (6.2.7.3), the homomorphisms g_3, g_4 send (6.2.7.1) respectively to

$$\begin{aligned} &a_1[3,15,3] + a_2[15,3,3] + \gamma_{71}[3,7,11] \\ &\quad + \gamma_{80}[7,3,11] + a_3[7,11,3] + \gamma_{47}[7,7,7] = 0, \\ &a_4[3,15,3] + a_5[15,3,3] + a_6[3,7,11] \\ &\quad + a_7[7,3,11] + a_8[7,11,3] + \gamma_{48}[7,7,7] = 0, \end{aligned}$$

where

$$\begin{aligned} &a_1 = \gamma_{\{35,44,74,85,93\}}, \quad a_2 = \gamma_{\{36,75,81,86,94\}}, \\ &a_3 = \gamma_{\{74,81,82,93,94\}}, \quad a_4 = \gamma_{\{42,44,68,71,76,77,83,85\}}, \\ &a_5 = \gamma_{\{43,69,75,76,77,80,84,86\}}, \quad a_6 = \gamma_{\{72,74,87,88,93\}}, \\ &a_7 = \gamma_{\{73,81,87,88,94\}}, \quad a_8 = \gamma_{\{38,67,68,69,71,80,82,83,84\}}. \end{aligned}$$

From the above equalities, we obtain

$$\begin{cases} \gamma_{47} = \gamma_{48} = \gamma_{71} = \gamma_{80} = 0, \\ a_i = 0,\ 1 \leqslant i \leqslant 8. \end{cases} \tag{6.2.7.4}$$

Combining (6.2.7.2), (6.2.7.3) and (6.2.7.4), we get

$$\begin{cases} \gamma_j = 0,\ j \neq 35, 36, 42, 43, 44, 68, 69, 72, 73, 74, 75, \\ \qquad\qquad 76, 77, 81, 85, 86, 87, 88, 91, 92, 93, 94, \\ \gamma_{35} = \gamma_{36} = \gamma_{68} = \gamma_{69} = \gamma_{91} = \gamma_{92}, \quad \gamma_{42} = \gamma_{43} = \gamma_{72} = \gamma_{73}, \\ \gamma_{44} = \gamma_{74} = \gamma_{76} = \gamma_{87},\ \gamma_{75} = \gamma_{77} = \gamma_{81} = \gamma_{88}, \\ \gamma_{\{42,44,94\}} = \gamma_{\{42,75,93\}} = \gamma_{\{35,42,44,86\}} = \gamma_{\{35,42,75,85\}} = 0. \end{cases} \tag{6.2.7.5}$$

Substituting (6.2.7.5) into the relation (6.2.7.1), we have

$$\gamma_{35}[\theta_1] + \gamma_{42}[\theta_2] + \gamma_{44}[\theta_3] + \gamma_{75}[\theta_4] = 0, \tag{6.2.7.6}$$

where

$$\begin{aligned} \theta_1 &= a_{1,2,35} + a_{1,2,36} + a_{1,2,68} + a_{1,2,69} \\ &\quad + a_{1,2,85} + a_{1,2,86} + a_{1,2,91} + a_{1,2,92}, \\ \theta_2 &= a_{1,2,42} + a_{1,2,43} + a_{1,2,72} + a_{1,2,73} \\ &\quad + a_{1,2,85} + a_{1,2,86} + a_{1,2,93} + a_{1,2,94}, \\ \theta_3 &= a_{1,2,44} + a_{1,2,74} + a_{1,2,76} + a_{1,2,86} + a_{1,2,87} + a_{1,2,94}, \\ \theta_4 &= a_{1,2,75} + a_{1,2,77} + a_{1,2,81} + a_{1,2,85} + a_{1,2,88} + a_{1,2,93}. \end{aligned}$$

We need to show that $\gamma_{35} = \gamma_{42} = \gamma_{44} = \gamma_{75} = 0$. The proof is divided into 4 steps.

Step 1. The homomorphism φ_4 sends (6.2.7.6) to

$$(\gamma_{35}+\gamma_{42})[\theta_1]+\gamma_{42}[\theta_2]+\gamma_{44}[\theta_3]+\gamma_{75}[\theta_4]=0. \tag{6.2.7.7}$$

Combining (6.2.7.6) and (6.2.7.7) gives

$$\gamma_{42}[\theta_1]=0.$$

We prove $[\theta_1]\neq 0$. Suppose $[\theta_1]=0$. Then, the polynomial θ_1 is hit and we have

$$\theta_1=Sq^1(A)+Sq^2(B)+Sq^4(C)+Sq^8(D),$$

for some polynomials $A\in(R_4)_{20}, B\in(R_4)_{19}, C\in(R_4)_{17}, D\in(R_4)_{13}$.

Let $(Sq^2)^3$ act on the both sides of this equality. Since $(Sq^2)^3Sq^1=0$ and $(Sq^2)^3Sq^2=0$, we get

$$(Sq^2)^3(\theta_1)=(Sq^2)^3Sq^4(C)+(Sq^2)^3Sq^8(D).$$

By a direct computation, we see that the monomial $(8,7,8,3)$ is a term of $(Sq^2)^3(\theta_1)$. This monomial is not a term of $(Sq^2)^3Sq^4(u)$ for $u\neq(3,7,4,3),(4,7,3,3)$. It is also not a term of $(Sq^2)^3Sq^8(D)$ for all polynomial D. So, either $(3,7,4,3)$ or $(4,7,3,3)$ is a term of C.

If $(3,7,4,3)$ is a term of C then

$$(Sq^2)^3(\theta_1+Sq^4(3,7,4,3))=(Sq^2)^3(Sq^4(C')+Sq^8(D)),$$

where $C'=C+(3,7,4,3)$. The monomial $(4,16,4,3)$ is a term of the polynomial $(Sq^2)^3(\theta_1+Sq^4(3,7,4,3))$. This monomial is not a term of the polynomial $(Sq^2)^3Sq^4(u)$ for $u\neq(3,7,4,3),(4,7,3,3)$. It is also not a term of $(Sq^2)^3Sq^8(D)$ for all $D\in(R_4)_{13}$. So, either $(3,7,4,3)$ or $(4,7,3,3)$ is a term of C'. Since $(3,7,4,3)$ is not a term of C', the monomial $(4,7,3,3)$ is a term of C'. Then we have

$$(Sq^2)^3(\theta_1+Sq^4((3,7,4,3)+(4,7,3,3)))=(Sq^2)^3(Sq^4(C'')+Sq^8(D)),$$

where $C''=C'+(4,7,3,3)=C+(3,7,4,3)+(4,7,3,3)$. Now the monomial $(8,7,8,3)$ is a term of $(Sq^2)^3(\theta_1+Sq^4((3,7,4,3)+(4,7,3,3)))$. Hence, either $(3,7,4,3)$ or $(4,7,3,3)$ is a term of C''. On the other hand, the two monomials $(3,7,4,3)$ and $(4,7,3,3)$ are not a term of C'', so we have a contradiction.

By a same argument, if $(4,7,3,3)$ is a term of C then we also have a contradiction. Hence, $[\theta_1]\neq 0$ and $\gamma_{42}=0$.

Step 2. Since $\gamma_{42}=0$, the homomorphism φ_1 sends (6.2.7.6) to

$$\gamma_{35}[\theta_2]+\gamma_{44}[\theta_3]+\gamma_{75}[\theta_4]=0. \tag{6.2.7.8}$$

Using (6.2.7.8) and by an analogous argument as given in Step 1, we get $\gamma_{35}=0$.

Step 3. The homomorphism φ_2 sends (6.2.7.8) to

$$\gamma_{44}[\theta_1]+\gamma_{75}[\theta_4]=0. \tag{6.2.7.9}$$

Using the relation (6.2.7.9) and by a same argument as given in Step 2, we obtain $\gamma_{44}=0$.

Step 4. The homomorphism φ_3 sends (6.2.7.9) to

$$\gamma_{75}[\theta_3] = 0.$$

Using this equality and by a same argument as given in Step 3, we get $\gamma_{75} = 0$. The proposition is proved. □

Proposition 6.2.8. *For $s \geqslant 3$, the elements $[a_{1,s,j}]$, $29 \leqslant j \leqslant 105$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}-3}$.*

Proof. Suppose there is a linear relation

$$\sum_{j=29}^{105} \gamma_j[a_{1,s,j}] = 0, \tag{6.2.8.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_i, i = 1, 2, 3, 4, 5, 6$, to (6.2.8.1), we have

$$\begin{aligned}
&\gamma_{49}w_{1,s,1} + \gamma_{50}w_{1,s,2} + \gamma_{37}w_{1,s,3} + \gamma_{61}w_{1,s,4}\\
&\qquad + \gamma_{39}w_{1,s,5} + \gamma_{40}w_{1,s,6} + \gamma_{29}w_{1,s,7} = 0,\\
&\gamma_{51}w_{1,s,1} + \gamma_{53}w_{1,s,2} + \gamma_{36}w_{1,s,3} + \gamma_{62}w_{1,s,4}\\
&\qquad + \gamma_{38}w_{1,s,5} + \gamma_{69}w_{1,s,6} + \gamma_{30}w_{1,s,7} = 0,\\
&\gamma_{52}w_{1,s,1} + \gamma_{54}w_{1,s,2} + a_1w_{1,s,3} + \gamma_{63}w_{1,s,4}\\
&\qquad + \gamma_{67}w_{1,s,5} + \gamma_{68}w_{1,s,6} + \gamma_{31}w_{1,s,7} = 0,\\
&\gamma_{55}w_{1,s,1} + \gamma_{43}w_{1,s,2} + \gamma_{58}w_{1,s,3} + \gamma_{41}w_{1,s,4}\\
&\qquad + \gamma_{64}w_{1,s,5} + \gamma_{73}w_{1,s,6} + \gamma_{32}w_{1,s,7} = 0,\\
&\gamma_{56}w_{1,s,1} + a_2w_{1,s,2} + \gamma_{59}w_{1,s,3} + \gamma_{70}w_{1,s,4}\\
&\qquad + \gamma_{65}w_{1,s,5} + \gamma_{72}w_{1,s,6} + \gamma_{33}w_{1,s,7} = 0,\\
&a_3w_{1,s,1} + \gamma_{57}w_{1,s,2} + \gamma_{60}w_{1,s,3} + \gamma_{71}w_{1,s,4}\\
&\qquad + \gamma_{74}w_{1,s,5} + \gamma_{66}w_{1,s,6} + \gamma_{34}w_{1,s,7} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases} \gamma_{\{35,103\}}, & s = 3,\\ \gamma_{35}, & s \geqslant 4, \end{cases} \quad a_2 = \begin{cases} \gamma_{\{42,103\}}, & s = 3,\\ \gamma_{42}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{44,103,104,105\}}, & s = 3,\\ \gamma_{44}, & s \geqslant 4. \end{cases}$$

From these equalities, we get

$$\begin{cases} a_1 = a_2 = a_3 = 0,\\ \gamma_j = 0,\ j = 29, \ldots, 34, 36, \ldots, 41, 43, 49, \ldots, 74. \end{cases} \tag{6.2.8.2}$$

With the aid of (6.2.8.2), the homomorphisms g_1, g_2, g_3, g_4 send (6.2.8.1) respectively to

$$\begin{aligned}
&\gamma_{85}w_{1,s,1} + \gamma_{90}w_{1,s,2} + \gamma_{76}w_{1,s,3} + \gamma_{94}w_{1,s,4}\\
&\qquad + \gamma_{78}w_{1,s,5} + \gamma_{84}w_{1,s,6} + \gamma_{45}w_{1,s,7} = 0,\\
&\gamma_{86}w_{1,s,1} + a_4w_{1,s,2} + \gamma_{77}w_{1,s,3} + \gamma_{95}w_{1,s,4}\\
&\qquad + \gamma_{79}w_{1,s,5} + \gamma_{83}w_{1,s,6} + \gamma_{46}w_{1,s,7} = 0,\\
&\gamma_{87}w_{1,s,1} + a_5w_{1,s,2} + a_6w_{1,s,3} + \gamma_{96}w_{1,s,4}\\
&\qquad + \gamma_{80}w_{1,s,5} + a_7w_{1,s,6} + \gamma_{47}w_{1,s,7} = 0,\\
&\gamma_{88}w_{1,s,1} + a_8w_{1,s,2} + a_9w_{1,s,3} + \gamma_{97}w_{1,s,4}\\
&\qquad + \gamma_{81}w_{1,s,5} + a_{10}w_{1,s,6} + \gamma_{48}w_{1,s,7} = 0,
\end{aligned}$$

where

$$a_4 = \begin{cases} \gamma_{\{35,89\}}, & s = 3,\\ \gamma_{89}, & s \geqslant 4, \end{cases} \quad a_5 = \begin{cases} \gamma_{\{44,91\}}, & s = 3,\\ \gamma_{91}, & s \geqslant 4, \end{cases}$$

$$a_6 = \begin{cases} \gamma_{\{75,105\}}, & s = 3,\\ \gamma_{75}, & s \geqslant 4, \end{cases} \quad a_7 = \begin{cases} \gamma_{\{82,104\}}, & s = 3,\\ \gamma_{82}, & s \geqslant 4, \end{cases}$$

$$a_8 = \begin{cases} \gamma_{\{44,92,99,100\}}, & s = 3,\\ \gamma_{92}, & s \geqslant 4, \end{cases} \quad a_9 = \begin{cases} \gamma_{\{93,99,101,105\}}, & s = 3,\\ \gamma_{93}, & s \geqslant 4, \end{cases}$$

$$a_{10} = \begin{cases} \gamma_{\{98,100,101,104\}}, & s = 3,\\ \gamma_{98}, & s \geqslant 4. \end{cases}$$

These equalities imply

$$\begin{cases} a_i = 0,\ i = 4, 5, \ldots, 10,\\ \gamma_j = 0,\ j = 45, \ldots, 48, 76, \ldots, 81,\\ 83, \ldots, 88, 90, 94, \ldots, 97. \end{cases} \tag{6.2.8.3}$$

With the aid of (6.2.8.2) and (6.2.8.3), the homomorphism h sends (6.2.8.1) to

$$\begin{aligned}
&a_{11}w_{1,s,1} + a_{12}w_{1,s,2} + \gamma_{99}w_{1,s,3} + a_{13}w_{1,s,4}\\
&\qquad + \gamma_{100}w_{1,s,5} + \gamma_{101}w_{1,s,6} + \gamma_{102}w_{1,s,7} = 0,
\end{aligned}$$

where

$$a_{11} = \begin{cases} \gamma_{92}, & s = 3,\\ \gamma_{103}, & s \geqslant 4, \end{cases} \quad a_{12} = \begin{cases} \gamma_{93}, & s = 3,\\ \gamma_{105}, & s \geqslant 4, \end{cases} \quad a_{13} = \begin{cases} \gamma_{98}, & s = 3,\\ \gamma_{104}, & s \geqslant 4. \end{cases}$$

From this, we get

$$\begin{cases} a_{11} = a_{12} = a_{13} = 0,\\ \gamma_j = 0,\ j = 99, 100, 101, 102. \end{cases} \tag{6.2.8.4}$$

Combining (6.2.8.2), 6.2.8.3) and (6.2.8.4), we obtain $\gamma_j = 0$ for $29 \leqslant j \leqslant 105$. The proposition is proved. □

Remark 6.2.9. The element $[\theta]$ defined as in the proof of Proposition 6.2.6 is an $GL_4(\mathbb{F}_2)$-invariant in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_9$.

The $\mathbb{F}_2$-subspace of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{21}$ generated by $[\theta_1], [\theta_2], [\theta_3], [\theta_4]$, which are defined as in the proof of Proposition 6.2.7, is an $GL_4(\mathbb{F}_2)$-submodule of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{21}$.

6.3. **The case $t = 2$.**

Recall that from Kameko [8], $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+3}+2^{s+1}-3} = 10$ with a basis consisting of all the classes $w_{2,s,i}, 1 \leqslant i \leqslant 10$, which are represented by the following monomials:

1. $(2^s - 1, 2^s - 1, 2^{s+3} - 1)$,
2. $(2^s - 1, 2^{s+3} - 1, 2^s - 1)$,
3. $(2^{s+3} - 1, 2^s - 1, 2^s - 1)$,
4. $(2^s - 1, 2^{s+1} - 1, 7.2^s - 1)$,
5. $(2^{s+1} - 1, 2^s - 1, 7.2^s - 1)$,
6. $(2^{s+1} - 1, 7.2^s - 1, 2^s - 1)$,
7. $(2^{s+1} - 1, 3.2^s - 1, 5.2^s - 1)$,
8. $(2^{s+1} - 1, 2^{s+2} - 1, 2^{s+2} - 1)$,
9. $(2^{s+2} - 1, 2^{s+1} - 1, 2^{s+2} - 1)$,
10. $(2^{s+2} - 1, 2^{s+2} - 1, 2^{s+1} - 1)$.

Hence, we easily obtain

Proposition 6.3.1. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+3}+2^{s+1}-3}$ is an $\mathbb{F}_2$-vector space of dimension 40 with a basis consisting of all the classes represented by the monomials $a_{2,s,j}, 1 \leqslant j \leqslant 40$, which are determined as follows:*

1. $(0, 2^{s+1} - 1, 2^{s+2} - 1, 2^{s+2} - 1)$,
2. $(0, 2^{s+2} - 1, 2^{s+1} - 1, 2^{s+2} - 1)$,
3. $(0, 2^{s+2} - 1, 2^{s+2} - 1, 2^{s+1} - 1)$,
4. $(2^{s+1} - 1, 0, 2^{s+2} - 1, 2^{s+2} - 1)$,
5. $(2^{s+1} - 1, 2^{s+2} - 1, 0, 2^{s+2} - 1)$,
6. $(2^{s+1} - 1, 2^{s+2} - 1, 2^{s+2} - 1, 0)$,
7. $(2^{s+2} - 1, 0, 2^{s+1} - 1, 2^{s+2} - 1)$,
8. $(2^{s+2} - 1, 0, 2^{s+2} - 1, 2^{s+1} - 1)$,
9. $(2^{s+2} - 1, 2^{s+1} - 1, 0, 2^{s+2} - 1)$,
10. $(2^{s+2} - 1, 2^{s+1} - 1, 2^{s+2} - 1, 0)$,
11. $(2^{s+2} - 1, 2^{s+2} - 1, 0, 2^{s+1} - 1)$,
12. $(2^{s+2} - 1, 2^{s+2} - 1, 2^{s+1} - 1, 0)$,
13. $(0, 2^s - 1, 2^s - 1, 2^{s+3} - 1)$,
14. $(0, 2^s - 1, 2^{s+3} - 1, 2^s - 1)$,
15. $(0, 2^{s+3} - 1, 2^s - 1, 2^s - 1)$,
16. $(2^s - 1, 0, 2^s - 1, 2^{s+3} - 1)$,
17. $(2^s - 1, 0, 2^{s+3} - 1, 2^s - 1)$,
18. $(2^s - 1, 2^s - 1, 0, 2^{s+3} - 1)$,
19. $(2^s - 1, 2^s - 1, 2^{s+3} - 1, 0)$,
20. $(2^s - 1, 2^{s+3} - 1, 0, 2^s - 1)$,
21. $(2^s - 1, 2^{s+3} - 1, 2^s - 1, 0)$,
22. $(2^{s+3} - 1, 0, 2^s - 1, 2^s - 1)$,
23. $(2^{s+3} - 1, 2^s - 1, 0, 2^s - 1)$,
24. $(2^{s+3} - 1, 2^s - 1, 2^s - 1, 0)$,
25. $(0, 2^s - 1, 2^{s+1} - 1, 7.2^s - 1)$,
26. $(0, 2^{s+1} - 1, 2^s - 1, 7.2^s - 1)$,
27. $(0, 2^{s+1} - 1, 7.2^s - 1, 2^s - 1)$,
28. $(2^s - 1, 0, 2^{s+1} - 1, 7.2^s - 1)$,
29. $(2^s - 1, 2^{s+1} - 1, 0, 7.2^s - 1)$,
30. $(2^s - 1, 2^{s+1} - 1, 7.2^s - 1, 0)$,
31. $(2^{s+1} - 1, 0, 2^s - 1, 7.2^s - 1)$,
32. $(2^{s+1} - 1, 0, 7.2^s - 1, 2^s - 1)$,
33. $(2^{s+1} - 1, 2^s - 1, 0, 7.2^s - 1)$,
34. $(2^{s+1} - 1, 2^s - 1, 7.2^s - 1, 0)$,
35. $(2^{s+1} - 1, 7.2^s - 1, 0, 2^s - 1)$,
36. $(2^{s+1} - 1, 7.2^s - 1, 2^s - 1, 0)$,
37. $(0, 2^{s+1} - 1, 3.2^s - 1, 5.2^s - 1)$,
38. $(2^{s+1} - 1, 0, 3.2^s - 1, 5.2^s - 1)$,
39. $(2^{s+1} - 1, 3.2^s - 1, 0, 5.2^s - 1)$,
40. $(2^{s+1} - 1, 3.2^s - 1, 5.2^s - 1, 0)$.

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+1}-3}$.

Set $\mu_3(1) = 87, \mu_3(2) = 135$ and $\mu_3(s) = 150$ for $s \geqslant 3$.

The main result of this subsection is

Theorem 6.3.2. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+1}-3}$ is an $\mathbb{F}_2$-vector space of dimension $\mu_2(s) - 40$ with a basis consisting of all the classes represented by the monomials $a_{2,s,j}, 41 \leqslant j \leqslant \mu_3(s)$, which are determined as follows:*

For $s \geqslant 1$,

41. $(1, 2^{s+1} - 2, 2^{s+2} - 1, 2^{s+2} - 1)$,
42. $(1, 2^{s+2} - 1, 2^{s+1} - 2, 2^{s+2} - 1)$,
43. $(1, 2^{s+2} - 1, 2^{s+2} - 1, 2^{s+1} - 2)$,
44. $(2^{s+2} - 1, 1, 2^{s+1} - 2, 2^{s+2} - 1)$,
45. $(2^{s+2} - 1, 1, 2^{s+2} - 1, 2^{s+1} - 2)$,
46. $(2^{s+2} - 1, 2^{s+2} - 1, 1, 2^{s+1} - 2)$,
47. $(1, 2^{s+1} - 1, 2^{s+2} - 2, 2^{s+2} - 1)$,
48. $(1, 2^{s+1} - 1, 2^{s+2} - 1, 2^{s+2} - 2)$,
49. $(1, 2^{s+2} - 2, 2^{s+1} - 1, 2^{s+2} - 1)$,
50. $(1, 2^{s+2} - 2, 2^{s+2} - 1, 2^{s+1} - 1)$,
51. $(1, 2^{s+2} - 1, 2^{s+1} - 1, 2^{s+2} - 2)$,
52. $(1, 2^{s+2} - 1, 2^{s+2} - 2, 2^{s+1} - 1)$,
53. $(2^{s+1} - 1, 1, 2^{s+2} - 2, 2^{s+2} - 1)$,
54. $(2^{s+1} - 1, 1, 2^{s+2} - 1, 2^{s+2} - 2)$,
55. $(2^{s+1} - 1, 2^{s+2} - 1, 1, 2^{s+2} - 2)$,
56. $(2^{s+2} - 1, 1, 2^{s+1} - 1, 2^{s+2} - 2)$,
57. $(2^{s+2} - 1, 1, 2^{s+2} - 2, 2^{s+1} - 1)$,
58. $(2^{s+2} - 1, 2^{s+1} - 1, 1, 2^{s+2} - 2)$,
59. $(3, 2^{s+2} - 3, 2^{s+1} - 2, 2^{s+2} - 1)$,
60. $(3, 2^{s+2} - 3, 2^{s+2} - 1, 2^{s+1} - 2)$,
61. $(3, 2^{s+2} - 1, 2^{s+2} - 3, 2^{s+1} - 2)$,
62. $(2^{s+2} - 1, 3, 2^{s+2} - 3, 2^{s+1} - 2)$,
63. $(3, 2^{s+1} - 1, 2^{s+2} - 3, 2^{s+2} - 2)$,
64. $(3, 2^{s+2} - 3, 2^{s+1} - 1, 2^{s+2} - 2)$,
65. $(3, 2^{s+2} - 3, 2^{s+2} - 2, 2^{s+1} - 1)$,
66. $(1, 2^s - 1, 2^s - 1, 2^{s+3} - 2)$,
67. $(1, 2^s - 1, 2^{s+3} - 2, 2^s - 1)$,
68. $(1, 2^{s+3} - 2, 2^s - 1, 2^s - 1)$,
69. $(1, 2^s - 1, 2^{s+1} - 2, 7.2^s - 1)$,
70. $(1, 2^{s+1} - 2, 2^s - 1, 7.2^s - 1)$,
71. $(1, 2^{s+1} - 2, 7.2^s - 1, 2^s - 1)$,
72. $(1, 2^{s+1} - 2, 3.2^s - 1, 5.2^s - 1)$.

For $s = 1$,

73. $(3, 3, 4, 7)$,
74. $(3, 3, 7, 4)$,
75. $(3, 7, 3, 4)$,
76. $(7, 3, 3, 4)$,
77. $(1, 1, 3, 12)$,
78. $(1, 3, 1, 12)$,
79. $(1, 3, 12, 1)$,
80. $(3, 1, 1, 12)$,
81. $(3, 1, 12, 1)$,
82. $(1, 3, 4, 9)$,
83. $(3, 1, 4, 9)$,
84. $(1, 3, 5, 8)$,
85. $(3, 1, 5, 8)$,
86. $(3, 5, 1, 8)$,
87. $(3, 5, 8, 1)$.

For $s \geqslant 2$,

73. $(2^{s+1} - 1, 3, 2^{s+2} - 3, 2^{s+2} - 2)$,
74. $(3, 2^{s+1} - 3, 2^{s+2} - 2, 2^{s+2} - 1)$,
75. $(3, 2^{s+1} - 3, 2^{s+2} - 1, 2^{s+2} - 2)$,
76. $(3, 2^{s+2} - 1, 2^{s+1} - 3, 2^{s+2} - 2)$,
77. $(2^{s+2} - 1, 3, 2^{s+1} - 3, 2^{s+2} - 2)$,
78. $(7, 2^{s+2} - 5, 2^{s+1} - 3, 2^{s+2} - 2)$,
79. $(7, 2^{s+2} - 5, 2^{s+2} - 3, 2^{s+1} - 2)$,
80. $(1, 2^s - 2, 2^s - 1, 2^{s+3} - 1)$,
81. $(1, 2^s - 2, 2^{s+3} - 1, 2^s - 1)$,
82. $(1, 2^s - 1, 2^s - 2, 2^{s+3} - 1)$,
83. $(1, 2^s - 1, 2^{s+3} - 1, 2^s - 2)$,
84. $(1, 2^{s+3} - 1, 2^s - 2, 2^s - 1)$,
85. $(1, 2^{s+3} - 1, 2^s - 1, 2^s - 2)$,
86. $(2^s - 1, 1, 2^s - 2, 2^{s+3} - 1)$,
87. $(2^s - 1, 1, 2^{s+3} - 1, 2^s - 2)$,
88. $(2^s - 1, 2^{s+3} - 1, 1, 2^s - 2)$,
89. $(2^{s+3} - 1, 1, 2^s - 2, 2^s - 1)$,
90. $(2^{s+3} - 1, 1, 2^s - 1, 2^s - 2)$,
91. $(2^{s+3} - 1, 2^s - 1, 1, 2^s - 2)$,
92. $(2^s - 1, 1, 2^s - 1, 2^{s+3} - 2)$,
93. $(2^s - 1, 1, 2^{s+3} - 2, 2^s - 1)$,
94. $(2^s - 1, 2^s - 1, 1, 2^{s+3} - 2)$,
95. $(1, 2^s - 2, 2^{s+1} - 1, 7.2^s - 1)$,
96. $(1, 2^{s+1} - 1, 2^s - 2, 7.2^s - 1)$,

97. $(1, 2^{s+1}-1, 7.2^s-1, 2^s-2)$, 98. $(2^{s+1}-1, 1, 2^s-2, 7.2^s-1)$,
99. $(2^{s+1}-1, 1, 7.2^s-1, 2^s-2)$, 100. $(2^{s+1}-1, 7.2^s-1, 1, 2^s-2)$,
101. $(2^s-1, 1, 2^{s+1}-2, 7.2^s-1)$, 102. $(1, 2^s-1, 2^{s+1}-1, 7.2^s-2)$,
103. $(1, 2^{s+1}-1, 2^s-1, 7.2^s-2)$, 104. $(1, 2^{s+1}-1, 7.2^s-2, 2^s-1)$,
105. $(2^s-1, 1, 2^{s+1}-1, 7.2^s-2)$, 106. $(2^s-1, 2^{s+1}-1, 1, 7.2^s-2)$,
107. $(2^{s+1}-1, 1, 2^s-1, 7.2^s-2)$, 108. $(2^{s+1}-1, 1, 7.2^s-2, 2^s-1)$,
109. $(2^{s+1}-1, 2^s-1, 1, 7.2^s-2)$, 110. $(1, 2^{s+1}-1, 3.2^s-2, 5.2^s-1)$,
111. $(2^{s+1}-1, 1, 3.2^s-2, 5.2^s-1)$, 112. $(1, 2^{s+1}-1, 3.2^s-1, 5.2^s-2)$,
113. $(2^{s+1}-1, 1, 3.2^s-1, 5.2^s-2)$, 114. $(2^{s+1}-1, 3.2^s-1, 1, 5.2^s-2)$,
115. $(3, 2^{s+1}-3, 2^s-2, 7.2^s-1)$, 116. $(3, 2^{s+1}-3, 7.2^s-1, 2^s-2)$,
117. $(3, 2^{s+1}-1, 7.2^s-3, 2^s-2)$, 118. $(2^{s+1}-1, 3, 7.2^s-3, 2^s-2)$,
119. $(3, 2^{s+1}-3, 3.2^s-2, 5.2^s-1)$, 120. $(3, 2^{s+1}-3, 3.2^s-1, 5.2^s-2)$,
121. $(3, 2^{s+1}-1, 3.2^s-3, 5.2^s-2)$, 122. $(2^{s+1}-1, 3, 3.2^s-3, 5.2^s-2)$,
123. $(3, 2^s-1, 2^{s+3}-3, 2^s-2)$, 124. $(3, 2^{s+3}-3, 2^s-2, 2^s-1)$,
125. $(3, 2^{s+3}-3, 2^s-1, 2^s-2)$, 126. $(3, 2^s-1, 2^{s+1}-3, 7.2^s-2)$,
127. $(3, 2^{s+1}-3, 2^s-1, 7.2^s-2)$, 128. $(3, 2^{s+1}-3, 7.2^s-2, 2^s-1)$.

For $s = 2$,

129. $(3, 3, 3, 28)$, 130. $(3, 3, 28, 3)$, 131. $(3, 3, 4, 27)$,
132. $(3, 3, 7, 24)$, 133. $(3, 7, 3, 24)$, 134. $(7, 3, 3, 24)$,
135. $(7, 7, 9, 14)$.

For $s \geqslant 3$,

129. $(2^s-1, 3, 2^{s+1}-3, 7.2^s-2)$, 130. $(2^s-1, 3, 2^{s+3}-3, 2^s-2)$,
131. $(3, 2^s-3, 2^s-2, 2^{s+3}-1)$, 132. $(3, 2^s-3, 2^{s+3}-1, 2^s-2)$,
133. $(3, 2^{s+3}-1, 2^s-3, 2^s-2)$, 134. $(2^{s+3}-1, 3, 2^s-3, 2^s-2)$,
135. $(3, 2^s-3, 2^s-1, 2^{s+3}-2)$, 136. $(3, 2^s-3, 2^{s+3}-2, 2^s-1)$,
137. $(3, 2^s-1, 2^s-3, 2^{s+3}-2)$, 138. $(2^s-1, 3, 2^s-3, 2^{s+3}-2)$,
139. $(3, 2^s-3, 2^{s+1}-2, 7.2^s-1)$, 140. $(3, 2^s-3, 2^{s+1}-1, 7.2^s-2)$,
141. $(3, 2^{s+1}-1, 2^s-3, 7.2^s-2)$, 142. $(2^{s+1}-1, 3, 2^s-3, 7.2^s-2)$,
143. $(7, 2^{s+3}-5, 2^s-3, 2^s-2)$, 144. $(7, 2^{s+1}-5, 2^s-3, 7.2^s-2)$,
145. $(7, 2^{s+1}-5, 3.2^s-3, 5.2^s-2)$, 146. $(7, 2^{s+1}-5, 2^{s+2}-3, 2^{s+2}-2)$,
147. $(7, 2^{s+1}-5, 7.2^s-3, 2^s-2)$.

For $s = 3$,

148. $(7, 7, 7, 56)$, 149. $(7, 7, 9, 54)$, 150. $(7, 7, 57, 6)$.

For $s \geqslant 4$,

148. $(7, 2^s-5, 2^s-3, 2^{s+3}-2)$, 149. $(7, 2^s-5, 2^{s+1}-3, 7.2^s-2)$,
150. $(7, 2^s-5, 2^{s+3}-3, 2^s-2)$.

We begin the proof of this theorem with the following proposition.

Proposition 6.3.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+1}-3}$ is generated by the $\mu_3(s) - 40$ elements listed in Theorem 6.3.2.*

The proof of the proposition is based on some lemmas.

Lemma 6.3.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&0&1&1\\1&0&0&0\\0&1&0&0\\0&1&0&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&0&0\\0&1&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&0&0\\0&1&0&0\\0&0&1&0\end{pmatrix}$$

$$\begin{pmatrix}1&0&1&1\\1&0&1&1\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&1&1\\0&1&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&1\\0&1&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&0&1&0\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (3,12,1,1), (3,4,1,9), (3,4,9,1), (3,4,3,7), (3,4,7,3), (3,7,4,3), (7,3,4,3).

By a direct computation, we have

$$\begin{aligned}(3,12,1,1) &= Sq^1(3,11,1,1) + Sq^2\big((5,7,2,1)+(2,11,1,1)\\ &\quad + (5,7,1,2)\big) + Sq^4\big((3,7,2,1)+(3,7,1,2)\big) + (3,9,4,1)\\ &\quad + (3,9,1,4) + (2,13,1,1) \mod \mathcal{L}_4(3;1;1;1),\\ (3,4,9,1) &= Sq^1(3,1,11,1) + Sq^2\big((5,2,7,1)+(2,1,11,1)\\ &\quad + (5,1,7,2)\big) + Sq^4\big((3,2,7,1)+(3,1,7,2)\big) + (3,1,12,1)\\ &\quad + (2,1,13,1) + (3,1,9,4) \mod \mathcal{L}_4(3;1;1;1),\\ (3,4,1,9) &= \overline{\varphi}_3(3,4,9,1).\end{aligned}$$

If x is one of the four monomials (3,4,3,7), (3,4,7,3), (3,7,4,3), (7,3,4,3) then there is a homomorphism $\psi : P_4 \to P_4$ induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = \psi(3,4,3,7)$.

A direct computation shows

$$\begin{aligned}(3,4,3,7) &= Sq^1(3,3,3,7) + Sq^2(2,3,3,7) + (3,3,4,7)\\ &\quad + (2,5,3,7) + (2,3,5,7) \mod \mathcal{L}_4(3;3;2).\end{aligned}$$

So, the lemma is proved. □

Lemma 6.3.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&1&0&1\\1&1&0&1\\0&1&0&0\\0&0&1&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&0&0&0\\0&0&1&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&1\\1&0&0&0\\0&1&0&0\\0&0&1&0\end{pmatrix}$$

$$\begin{pmatrix}1&1&0&1\\1&1&0&1\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&0&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 0 & 1 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$\begin{aligned}&(3,7,24,3), (7,3,24,3), (7,11,17,2), (3,7,8,19), (7,3,8,19),\\ &(3,7,11,16), (7,3,11,16), (7,11,3,16), (7,11,16,3), (7,7,8,15),\\ &(7,7,15,8), (7,15,7,8), (15,7,7,8).\end{aligned}$$

By a direct computation, we have

$$\begin{aligned}(7,3,24,3) &= Sq^1(7,3,23,3) + Sq^2(7,2,23,3) + Sq^4\big((4,3,23,3)\\ &\quad + (5,2,23,3) + (11,2,15,5) + (11,3,15,4)\big) + Sq^8\big((7,3,15,4)\\ &\quad + (7,2,15,5)\big) + (4,3,27,3) + (5,2,27,3) + (7,2,25,3)\\ &\quad + (7,2,19,9) + (7,3,19,8) \mod \mathcal{L}_4(3;3;1;1;1),\\ (3,7,24,3) &= \overline{\varphi}_1(7,3,24,3),\end{aligned}$$

where $\overline{\varphi}_1$ is defined as in Section 2.

$$\begin{aligned}(7,11,17,2) &= Sq^1(7,7,19,3) + Sq^2(7,7,19,2) + Sq^4\big((11,7,12,3)\\ &\quad + (11,7,13,2) + (4,7,19,3) + (5,7,19,2)\big) + Sq^8\big((7,7,12,3)\\ &\quad + (7,7,13,2)\big) + (4,11,19,3) + (5,11,19,2) + (7,9,19,2)\\ &\quad + (7,8,19,3) + (7,11,16,3) \mod \mathcal{L}_4(3;3;1;1;1),\\ (7,3,8,19) &= Sq^1(7,3,3,23) + Sq^2(7,2,3,23) + Sq^4\big((11,3,4,15)\\ &\quad + (4,3,3,23) + (5,2,3,23) + (11,2,5,15)\big) + Sq^8\big((7,3,4,15)\\ &\quad + (7,2,5,15)\big) + (7,3,3,24) + (4,3,3,27) + (7,2,3,25)\\ &\quad + (5,2,3,27) + (7,2,9,19) \mod \mathcal{L}_4(3;3;1;1;1),\\ (3,7,8,19) &= \overline{\varphi}_1(7,3,8,19).\end{aligned}$$

If x is one of the monomials (3,7,11,16), (7,3,11,16), (7,11,3,16), then there is a homomorphism $\psi : P_4 \to P_4$ induced by a permutation of

$\{x_1, x_2, x_3, x_4\}$ such that $x = \psi(7, 11, 3, 6)$. We have

$$\begin{aligned}(7,11,3,16) &= Sq^1\big((7,7,3,19) + (7,7,1,21)\big) + Sq^2(7,7,2,19)\\ &\quad + Sq^4\big((11,7,3,12) + (4,7,3,19) + (5,7,2,19) + (11,7,1,14)\big)\\ &\quad + Sq^8\big((7,7,3,12) + (7,7,1,14)\big) + (4,11,3,19) + (5,11,2,19)\\ &\quad + (7,8,3,19) + (7,9,2,19) + (7,11,1,18) \mod \mathcal{L}_4(3;3;1;1;1),\\ (7,11,16,3) &= Sq^1(11,19,3,3) + Sq^2\big((10,19,3,3) + (7,22,3,3)\big)\\ &\quad + Sq^4\big((7,19,4,3) + (7,19,3,4) + (6,21,3,3) + (6,19,5,3)\\ &\quad + (6,19,3,5) + (7,20,3,3) + (5,22,3,3) + (11,14,5,3)\\ &\quad + (11,14,3,5)\big) + Sq^8\big((7,11,8,3) + (7,11,3,8) + (7,14,5,3)\\ &\quad + (7,14,3,5) + (7,10,9,3) + (7,10,3,9)\big) + (7,11,3,16)\\ &\quad + (6,25,3,3) + (6,19,9,3) + (6,19,3,9) + (5,26,3,3)\\ &\quad + (7,10,17,3) + (7,10,3,17) \mod \mathcal{L}_4(3;3;1;1;1).\end{aligned}$$

If x is one of the monomials (7,7,8,15), (7,7,15,8), (7,15,7,8), (15,7,7,8), then there is a homomorphism $\psi : P_4 \to P_4$ induced by a permutation of $\{x_1, x_2, x_3, x_4\}$ such that $x = \psi(7, 7, 8, 15)$. We have

$$\begin{aligned}(7,7,8,15) &= Sq^1(7,7,7,15) + Sq^2(7,6,7,15) + Sq^4\big((4,7,7,15)\\ &\quad + (5,6,7,15)\big) + (4,11,7,15) + (4,7,11,15) + (7,6,9,15)\\ &\quad + (5,10,7,15) + (5,6,11,15) \mod \mathcal{L}_4(3;3;3;2).\end{aligned}$$

The lemma follows. □

Combining Theorem 2.5, the lemmas in Sections 3, 5 and Lemmas 6.1.1, 6.2.4, 6.2.5, 6.3.4, 6.3.5, we obtain Proposition 6.3.3.

Now, we show that the elements $[a_{2,s,j}], 41 \leqslant j \leqslant \mu_3(s)$, are linearly independent.

Proposition 6.3.6. *The elements* $[a_{2,1,j}]$, $41 \leqslant j \leqslant 87$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{17}$.

Proof. Suppose there is a linear relation

$$\sum_{j=41}^{87} \gamma_j [a_{2,1,j}] = 0, \tag{6.3.6.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_i, i = 1, 2, 3$, to the relation (6.3.6.1), we get

$$\begin{aligned}
&\gamma_{68}[15,1,1] + \gamma_{70}[3,1,13] + \gamma_{71}[3,13,1]\\
&\quad + \gamma_{72}[3,5,9] + \gamma_{41}[3,7,7] + \gamma_{49}[7,3,7] + \gamma_{50}[7,7,3] = 0,\\
&\gamma_{\{67,81\}}[15,1,1] + \gamma_{\{44,53,57,69,83\}}[3,1,13] + \gamma_{\{79,87\}}[3,13,1]\\
&\quad + \gamma_{\{59,65,82\}}[3,5,9] + \gamma_{42}[3,7,7] + \gamma_{\{47,73\}}[7,3,7] + \gamma_{52}[7,7,3] = 0,\\
&\gamma_{\{66,80\}}[15,1,1] + a_1[3,1,13] + a_2[3,13,1] + a_3[3,5,9]\\
&\quad + \gamma_{43}[3,7,7] + \gamma_{\{48,74\}}[7,3,7] + \gamma_{\{51,75\}}[7,7,3] = 0,
\end{aligned}$$

where $a_1 = \gamma_{\{45,54,56,77,85\}},\ a_2 = \gamma_{\{46,55,58,78,86\}},\ a_3 = \gamma_{\{60,61,62,63,64,76,84\}}$.

Computing from the above equalities gives

$$\begin{cases}
a_1 = a_2 = a_3 = 0,\\
\gamma_j = 0,\ j = 41, 42, 43, 49, 50, 52, 68, 70, 71, 72,\\
\gamma_{\{67,81\}} = \gamma_{\{44,53,57,69,83\}} = \gamma_{\{79,87\}} = \gamma_{\{59,65,82\}} = 0,\\
\gamma_{\{47,73\}} = \gamma_{\{66,80\}} = \gamma_{\{48,74\}} = \gamma_{\{51,75\}} = 0.
\end{cases} \tag{6.3.6.2}$$

With the aid of (6.3.6.2), the homomorphisms f_4, f_5, f_6 send (6.3.6.1) to

$$\begin{aligned}
&\gamma_{\{67,79\}}[1,15,1] + \gamma_{\{47,69,82\}}[1,3,13] + \gamma_{\{81,87\}}[3,13,1]\\
&\quad + \gamma_{\{65,83\}}[3,5,9] + \gamma_{\{53,59,73\}}[3,7,7] + \gamma_{44}[7,3,7] + \gamma_{57}[7,7,3] = 0,\\
&\gamma_{\{66,78\}}[1,15,1] + \gamma_{\{48,51,77,84\}}[1,3,13] + a_4[3,13,1] + a_5[3,5,9]\\
&\quad + \gamma_{\{54,60,74\}}[3,7,7] + \gamma_{45}[7,3,7] + \gamma_{\{56,76\}}[7,7,3] = 0,\\
&\gamma_{\{66,67,69,77\}}[1,1,15] + a_6[1,3,13] + a_7[3,1,13] + a_8[3,5,9]\\
&\quad + \gamma_{\{55,61,75\}}[3,7,7] + \gamma_{\{58,62,76\}}[7,3,7] + \gamma_{46}[7,7,3] = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_4 = \gamma_{\{46,55,58,80,86\}},\quad a_5 = \gamma_{\{61,62,63,64,75,85\}},\\
&a_6 = \gamma_{\{47,48,51,78,79,82,84\}},\quad a_7 = \gamma_{\{44,45,53,54,56,57,80,81,83,85\}},\\
&a_8 = \gamma_{\{59,60,63,64,65,73,74,86,87\}}.
\end{aligned}$$

Computing directly from these equalities and using (6.3.6.2), we get

$$\begin{cases}
a_4 = a_5 = a_6 = a_7 = a_8 = 0,\\
\gamma_j = 0,\ j = 44, 45, 46, 57,\\
\gamma_{\{67,79\}} = \gamma_{\{47,69,82\}} = \gamma_{\{81,87\}} = \gamma_{\{65,83\}} = 0,\\
\gamma_{\{53,59,73\}} = \gamma_{\{66,78\}} = \gamma_{\{48,51,77,84\}} = \gamma_{\{54,60,74\}} = 0,\\
\gamma_{\{56,76\}} = \gamma_{\{66,67,69,77\}} = \gamma_{\{55,61,75\}} = \gamma_{\{58,62,76\}} = 0.
\end{cases} \tag{6.3.6.3}$$

With the aid of (6.3.6.2) and (6.3.6.3), the homomorphisms g_1, g_2 send (6.3.6.1) to

$$\begin{aligned}
&\gamma_{\{53,65,69\}}[1,3,13] + \gamma_{\{47,59,82\}}[3,5,9]\\
&\quad + \gamma_{53}[3,7,7] + \gamma_{59}[7,3,7] + \gamma_{65}[7,7,3] = 0,\\
&\gamma_{\{48,51,56,60,64,84,85\}}[3,5,9] + \gamma_{54}[3,7,7]\\
&\quad + \gamma_{\{54,56,77,85\}}[1,3,13] + \gamma_{60}[7,3,7] + \gamma_{64}[7,7,3] = 0.
\end{aligned}$$

From the above equalities, we get

$$\begin{cases}
\gamma_j = 0,\ j = 53, 54, 59, 60, 64, 65,\\
\gamma_{\{53,65,69\}} = \gamma_{\{54,56,77,85\}} = 0,\\
\gamma_{\{47,59,82\}} = \gamma_{\{48,51,56,60,64,84,85\}} = 0.
\end{cases} \tag{6.3.6.4}$$

With the aid of (6.3.6.2), (6.3.6.3) and (6.3.6.4), the homomorphisms g_3, g_4 send (6.3.6.1) respectively to

$$\begin{aligned}
&\gamma_{\{55,58,66,86\}}[1,3,13] + \gamma_{\{67,79\}}[3,1,13] + \gamma_{\{66,67,69,77\}}[3,13,1]\\
&\quad + a_9[3,5,9] + \gamma_{55}[3,7,7] + \gamma_{61}[7,3,7] + \gamma_{\{56,58,62,63\}}[7,7,3] = 0,\\
&\gamma_{\{55,58,66,86\}}[1,3,13] + \gamma_{\{67,79\}}[3,1,13] + \gamma_{\{66,67,69,77\}}[3,13,1]\\
&\quad + a_{10}[3,5,9] + \gamma_{58}[3,7,7] + \gamma_{62}[7,3,7] + a_{11}[7,7,3] = 0,
\end{aligned}$$

where

$$\begin{aligned}
a_9 &= \gamma_{\{47,48,51,56,61,62,63,79,82,84,86\}},\\
a_{10} &= \gamma_{\{47,48,51,56,61,62,63,79,85,86\}},\quad a_{11} = \gamma_{\{47,48,51,55,61,63\}}.
\end{aligned}$$

Hence, we get

$$\begin{cases}
a_9 = a_{10} = a_{11} = 0,\ \gamma_j = 0,\ j = 55, 58, 61, 62,\\
\gamma_{\{55,58,66,86\}} = \gamma_{\{67,79\}} = \gamma_{\{66,67,69,77\}} = \gamma_{\{56,58,62,63\}} = 0.
\end{cases} \tag{6.3.6.5}$$

Combining (6.3.6.2), (6.3.6.3), (6.3.6.4) and (6.3.6.5), we obtain

$$\begin{cases}
\gamma_j = 0,\ j \neq 66, 67, 78, 79, 80, 81, 86, 87,\\
\gamma_{66} = \gamma_{67} = \gamma_{78} = \gamma_{79} = \gamma_{80} = \gamma_{81} = \gamma_{86} = \gamma_{87}.
\end{cases} \tag{6.3.6.6}$$

Substituting (6.3.6.6) into the relation (6.3.6.1), we have

$$\gamma_{66}[\theta] = 0, \tag{6.3.6.7}$$

where $\theta = a_{2,1,66} + a_{2,1,67} + a_{2,1,78} + a_{2,1,79} + a_{2,1,80} + a_{2,1,81} + a_{2,1,86} + a_{2,1,87}$.

By a same argument as given in the proof of Proposition 6.2.7, we see that the polynomial θ is non-hit. Hence, we get $[\theta] \neq 0$ and the relation (6.3.6.7) implies $\gamma_{66} = 0$. The proposition is proved. □

Proposition 6.3.7. *The elements* $[a_{2,2,j}]$, $41 \leqslant j \leqslant 135$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{37}$.

Proof. Suppose there is a linear relation

$$\sum_{j=41}^{135} \gamma_j [a_{2,2,j}] = 0, \tag{6.3.7.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms $f_i, 1 \leqslant i \leqslant 6$, to the relation (6.3.7.1) and we get

$$\begin{aligned}
&\gamma_{80}[3,3,31] + \gamma_{81}[3,31,3] + \gamma_{68}[31,3,3] + \gamma_{95}[3,7,27]\\
&\quad + \gamma_{70}[7,3,27] + \gamma_{71}[7,27,3] + \gamma_{72}[7,11,19]\\
&\quad + \gamma_{41}[7,15,15] + \gamma_{49}[15,7,15] + \gamma_{50}[15,15,7] = 0,\\
&\gamma_{82}[3,3,31] + \gamma_{84}[3,31,3] + \gamma_{\{67,130\}}[31,3,3] + \gamma_{96}[3,7,27]\\
&\quad + \gamma_{\{69,131\}}[7,3,27] + \gamma_{104}[7,27,3] + \gamma_{110}[7,11,19]\\
&\quad + \gamma_{42}[7,15,15] + \gamma_{47}[15,7,15] + \gamma_{52}[15,15,7] = 0,\\
&\gamma_{83}[3,3,31] + \gamma_{85}[3,31,3] + \gamma_{\{66,129,134\}}[31,3,3] + \gamma_{97}[3,7,27]\\
&\quad + \gamma_{\{102,132\}}[7,3,27] + \gamma_{\{103,133\}}[7,27,3] + \gamma_{112}[7,11,19]\\
&\quad + \gamma_{43}[7,15,15] + \gamma_{48}[15,7,15] + \gamma_{51}[15,15,7] = 0,\\
&\gamma_{86}[3,3,31] + \gamma_{\{93,124,128,130\}}[3,31,3] + \gamma_{89}[31,3,3]\\
&\quad + a_1[3,7,27] + \gamma_{98}[7,3,27] + \gamma_{108}[7,27,3] + \gamma_{111}[7,11,19]\\
&\quad + \gamma_{53}[7,15,15] + \gamma_{44}[15,7,15] + \gamma_{57}[15,15,7] = 0,\\
&\gamma_{87}[3,3,31] + \gamma_{\{92,125,127,129,133\}}[3,31,3] + \gamma_{90}[31,3,3]\\
&\quad + a_2[3,7,27] + \gamma_{99}[7,3,27] + \gamma_{\{107,134\}}[7,27,3] + \gamma_{113}[7,11,19]\\
&\quad + \gamma_{54}[7,15,15] + \gamma_{45}[15,7,15] + \gamma_{56}[15,15,7] = 0,\\
&a_3[3,3,31] + \gamma_{88}[3,31,3] + \gamma_{91}[31,3,3] + a_4[3,7,27]\\
&\quad + a_5[7,3,27] + \gamma_{100}[7,27,3] + \gamma_{\{78,79,114,135\}}[7,11,19]\\
&\quad + \gamma_{55}[7,15,15] + \gamma_{58}[15,7,15] + \gamma_{46}[15,15,7] = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{59,65,74,101,115,119,131\}}, \quad a_2 = \gamma_{\{60,64,75,105,116,120,132\}},\\
&a_3 = \gamma_{\{94,123,126,129,130,131,132\}}, \quad a_4 = \gamma_{\{61,63,76,106,117,121,133\}},\\
&a_5 = \gamma_{\{62,73,77,109,118,122,134\}}.
\end{aligned}$$

Computing from the above equalities gives

$$\begin{cases}
\gamma_j = 0,\ j = 41,\ldots,57,58,68,70,71,72,80,\ldots,91,\\
95,\ldots,100,104,108,110,111,112,113,\\
\gamma_{\{67,130\}} = \gamma_{\{69,131\}} = \gamma_{\{66,129,134\}} = \gamma_{\{102,132\}} = 0,\\
\gamma_{\{103,133\}} = \gamma_{\{93,124,128,130\}} = \gamma_{\{92,125,127,129,133\}} = 0,\\
\gamma_{\{107,134\}} = \gamma_{\{78,79,114,135\}} = 0.
\end{cases} \tag{6.3.7.2}$$

With the aid of (6.3.7.2), the homomorphisms g_1, g_2 send (6.3.7.1) to

$$\begin{aligned}
&\gamma_{93}[3,31,3] + \gamma_{124}[31,3,3] + \gamma_{101}[3,7,27]\\
&\quad + \gamma_{115}[7,3,27] + \gamma_{128}[7,27,3] + \gamma_{119}[7,11,19]\\
&\quad + \gamma_{74}[7,15,15] + \gamma_{59}[15,7,15] + \gamma_{65}[15,15,7] = 0,\\
&\gamma_{\{66,92,129\}}[3,31,3] + \gamma_{125}[31,3,3] + \gamma_{105}[3,7,27]\\
&\quad + \gamma_{116}[7,3,27] + \gamma_{127}[7,27,3] + \gamma_{120}[7,11,19]\\
&\quad + \gamma_{75}[7,15,15] + \gamma_{60}[15,7,15] + \gamma_{64}[15,15,7] = 0.
\end{aligned}$$

From the above equalities, we get

$$\begin{cases}
\gamma_j = 0,\ j = 59, 60, 64, 65, 74, 75, 93, 101,\\
105, 115, 116, 119, 120, 124, 125, 127, 128,\\
\gamma_{\{66,92,129\}} = 0.
\end{cases} \tag{6.3.7.3}$$

With the aid of (6.3.7.2) and (6.3.7.3), the homomorphisms g_3, g_4 send (6.3.7.1) respectively to

$$\begin{aligned}
&\gamma_{\{66,94,107,109,129\}}[3,31,3] + \gamma_{\{118,123\}}[31,3,3] + \gamma_{\{106,114\}}[3,7,27]\\
&\quad + \gamma_{\{107,109,118,126\}}[7,27,3] + \gamma_{117}[7,3,27] + \gamma_{\{114,121,135\}}[7,11,19]\\
&\quad + \gamma_{76}[7,15,15] + \gamma_{61}[15,7,15] + \gamma_{\{63,135\}}[15,15,7] = 0,\\
&\gamma_{\{92,94,103,106,129\}}[3,31,3] + \gamma_{\{67,117,123\}}[31,3,3]\\
&\quad + \gamma_{118}[7,3,27] + \gamma_{\{69,102,103,106,117,126\}}[7,27,3]\\
&\quad + \gamma_{\{109,114\}}[3,7,27] + \gamma_{\{78,79,114,122,135\}}[7,11,19]\\
&\quad + \gamma_{77}[7,15,15] + \gamma_{62}[15,7,15] + \gamma_{\{73,78,79,135\}}[15,15,7] = 0.
\end{aligned}$$

Hence, we get

$$\begin{cases}
\gamma_j = 0,\ j = 59, 60, 64, 61, 62, 76, 77, 117, 118,\\
\gamma_{\{66,94,107,109,129\}} = \gamma_{\{118,123\}} = \gamma_{\{106,114\}} = 0,\\
\gamma_{\{114,121,135\}} = \gamma_{\{107,109,118,126\}} = \gamma_{\{63,135\}} = 0,\\
\gamma_{\{92,94,103,106,129\}} = \gamma_{\{109,114\}} = \gamma_{\{69,102,103,106,117,126\}} = 0,\\
\gamma_{\{67,117,123\}} = \gamma_{\{78,79,114,122,135\}} = \gamma_{\{73,78,79,135\}} = 0.
\end{cases} \tag{6.3.7.4}$$

With the aid of (6.3.7.2), (6.3.7.3), and (6.3.7.4), the homomorphism h sends (6.3.7.1) to

$$\begin{aligned}
&\gamma_{\{66,92,94,106,129\}}[3,3,31] + \gamma_{\{106,126\}}[3,7,27] + \gamma_{\{106,121,122\}}[7,11,19]\\
&\quad + \gamma_{73}[7,15,15] + \gamma_{78}[15,7,15] + \gamma_{79}[15,15,7] = 0.
\end{aligned}$$

From this equality, we get

$$\begin{cases}
\gamma_{73} = \gamma_{78} = \gamma_{79} = 0,\\
\gamma_{\{66,92,94,106,129\}} = \gamma_{\{106,126\}} = \gamma_{\{106,121,122\}} = 0.
\end{cases} \tag{6.3.7.5}$$

Combining (6.3.7.2), (6.3.7.3), (6.3.7.4) and (6.3.7.5), we obtain $\gamma_j = 0$ for any j. The proposition is proved. □

Proposition 6.3.8. *For any $s \geqslant 3$, the elements $[a_{2,s,j}]$, $41 \leqslant j \leqslant 150$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+1}-3}$.*

Proof. Suppose there is a linear relation

$$\sum_{j=41}^{150} \gamma_j [a_{2,s,j}] = 0, \tag{6.3.8.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_i, i = 1, 2, \ldots, 6$, to (6.3.8.1), we get

$$\begin{aligned}
&\gamma_{80}w_{2,s,1} + \gamma_{81}w_{2,s,2} + \gamma_{68}w_{2,s,3} + \gamma_{95}w_{2,s,4} + \gamma_{70}w_{2,s,5}\\
&\quad + \gamma_{71}w_{2,s,6} + \gamma_{72}w_{2,s,7} + \gamma_{41}w_{2,s,8} + \gamma_{49}w_{2,s,9} + \gamma_{50}w_{2,s,10} = 0,\\
&\gamma_{82}w_{2,s,1} + \gamma_{84}w_{2,s,2} + \gamma_{67}w_{2,s,3} + \gamma_{96}w_{2,s,4} + \gamma_{69}w_{2,s,5}\\
&\quad + \gamma_{104}w_{2,s,6} + \gamma_{110}w_{2,s,7} + \gamma_{42}w_{2,s,8} + \gamma_{47}w_{2,s,9} + \gamma_{52}w_{2,s,10} = 0,\\
&\gamma_{83}w_{2,s,1} + \gamma_{85}w_{2,s,2} + a_1w_{2,s,3} + \gamma_{97}w_{2,s,4} + \gamma_{102}w_{2,s,5}\\
&\quad + \gamma_{103}w_{2,s,6} + \gamma_{112}w_{2,s,7} + \gamma_{43}w_{2,s,8} + \gamma_{48}w_{2,s,9} + \gamma_{51}w_{2,s,10} = 0,\\
&\gamma_{86}w_{2,s,1} + \gamma_{93}w_{2,s,2} + \gamma_{89}w_{2,s,3} + \gamma_{101}w_{2,s,4} + \gamma_{98}w_{2,s,5}\\
&\quad + \gamma_{108}w_{2,s,6} + \gamma_{111}w_{2,s,7} + \gamma_{53}w_{2,s,8} + \gamma_{44}w_{2,s,9} + \gamma_{57}w_{2,s,10} = 0,\\
&\gamma_{87}w_{2,s,1} + a_2w_{2,s,2} + \gamma_{90}w_{2,s,3} + \gamma_{105}w_{2,s,4} + \gamma_{99}w_{2,s,5}\\
&\quad + \gamma_{107}w_{2,s,6} + \gamma_{113}w_{2,s,7} + \gamma_{54}w_{2,s,8} + \gamma_{45}w_{2,s,9} + \gamma_{56}w_{2,s,10} = 0,\\
&a_3w_{2,s,1} + \gamma_{88}w_{2,s,2} + \gamma_{91}w_{2,s,3} + \gamma_{106}w_{2,s,4} + \gamma_{109}w_{2,s,5}\\
&\quad + \gamma_{100}w_{2,s,6} + \gamma_{114}w_{2,s,7} + \gamma_{55}w_{2,s,8} + \gamma_{58}w_{2,s,9} + \gamma_{46}w_{2,s,10} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases} \gamma_{\{66,148\}}, & s = 3,\\ \gamma_{66}, & s \geqslant 4, \end{cases} \quad a_2 = \begin{cases} \gamma_{\{92,148\}}, & s = 3,\\ \gamma_{92}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{94,148,149,150\}}, & s = 3,\\ \gamma_{94}, & s \geqslant 4. \end{cases}$$

Computing from the above equalities, we obtain

$$\begin{cases} \gamma_j = 0,\ j = 29, \ldots, 34, 36, \ldots, 41, 43, 49, \ldots, 74,\\ a_1 = a_2 = a_3 = 0. \end{cases} \tag{6.3.8.2}$$

With the aid of (6.3.8.2), the homomorphisms g_1, g_2, g_3, g_4 send (6.3.8.1) to

$$\begin{aligned}
&\gamma_{131}w_{2,s,1} + \gamma_{136}w_{2,s,2} + \gamma_{124}w_{2,s,3} + \gamma_{139}w_{2,s,4} + \gamma_{115}w_{2,s,5}\\
&\quad + \gamma_{128}w_{2,s,6} + \gamma_{119}w_{2,s,7} + \gamma_{74}w_{2,s,8} + \gamma_{59}w_{2,s,9} + \gamma_{65}w_{2,s,10} = 0,\\
&\gamma_{132}w_{2,s,1} + a_4w_{2,s,2} + \gamma_{125}w_{2,s,3} + \gamma_{140}w_{2,s,4} + \gamma_{116}w_{2,s,5}\\
&\quad + \gamma_{127}w_{2,s,6} + \gamma_{120}w_{2,s,7} + \gamma_{75}w_{2,s,8} + \gamma_{60}w_{2,s,9} + \gamma_{64}w_{2,s,10} = 0,
\end{aligned}$$

$$\begin{aligned}&\gamma_{133}w_{2,s,1} + a_5w_{2,s,2} + a_6w_{2,s,3} + \gamma_{141}w_{2,s,4} + \gamma_{117}w_{2,s,5}\\ &\quad + a_7w_{2,s,6} + \gamma_{121}w_{2,s,7} + \gamma_{76}w_{2,s,8} + \gamma_{61}w_{2,s,9} + \gamma_{63}w_{2,s,10} = 0,\\ &\gamma_{134}w_{2,s,1} + a_8w_{2,s,2} + a_9w_{2,s,3} + \gamma_{142}w_{2,s,4} + \gamma_{118}w_{2,s,5}\\ &\quad + a_{10}w_{2,s,6} + \gamma_{122}w_{2,s,7} + \gamma_{77}w_{2,s,8} + \gamma_{62}w_{2,s,9} + \gamma_{73}w_{2,s,10} = 0,\end{aligned}$$

where

$$a_4 = \begin{cases}\gamma_{\{66,135\}}, & s = 3,\\ \gamma_{135}, & s \geqslant 4,\end{cases} \quad a_5 = \begin{cases}\gamma_{\{94,137\}}, & s = 3,\\ \gamma_{137}, & s \geqslant 4,\end{cases}$$

$$a_6 = \begin{cases}\gamma_{\{123,150\}}, & s = 3,\\ \gamma_{123}, & s \geqslant 4,\end{cases} \quad a_7 = \begin{cases}\gamma_{\{126,149\}}, & s = 3,\\ \gamma_{126}, & s \geqslant 4,\end{cases}$$

$$a_8 = \begin{cases}\gamma_{\{94,138,143,144\}}, & s = 3,\\ \gamma_{138}, & s \geqslant 4,\end{cases} \quad a_9 = \begin{cases}\gamma_{\{130,143,147,150\}}, & s = 3,\\ \gamma_{130}, & s \geqslant 4,\end{cases}$$

$$a_{10} = \begin{cases}\gamma_{\{129,144,147,149\}}, & s = 3,\\ \gamma_{129}, & s \geqslant 4.\end{cases}$$

These equalities imply

$$\text{(6.3.8.3)} \qquad \begin{cases}a_i = 0, i = 4, 5, \ldots, 10,\\ \gamma_j = 0,\ j = 59, \ldots, 65;\ 73, \ldots, 77;\ 115, \ldots, 122,\\ 124, 125, 127, 128, 131, \ldots, 134, 136, 139, \ldots, 142.\end{cases}$$

Using (6.3.8.2) and (6.3.8.3), we see that the image of (6.3.8.1) under the homomorphism h is

$$\begin{aligned}&a_{11}w_{2,s,1} + a_{12}w_{2,s,2} + \gamma_{143}w_{2,s,3} + a_{13}w_{2,s,4} + \gamma_{144}w_{2,s,5}\\ &\quad + \gamma_{147}w_{2,s,6} + \gamma_{145}w_{2,s,7} + \gamma_{146}w_{2,s,8} + \gamma_{78}w_{2,s,9} + \gamma_{79}w_{2,s,10} = 0,\end{aligned}$$

where

$$a_{11} = \begin{cases}\gamma_{138}, & s = 3,\\ \gamma_{148}, & s \geqslant 4,\end{cases} \quad a_{12} = \begin{cases}\gamma_{130}, & s = 3,\\ \gamma_{150}, & s \geqslant 4,\end{cases} \quad a_{13} = \begin{cases}\gamma_{129}, & s = 3,\\ \gamma_{149}, & s \geqslant 4.\end{cases}$$

From this equality, we obtain

$$\text{(6.3.8.4)} \qquad \begin{cases}\gamma_j = 0,\ j = 78, 79, 143, 144, 145, 146, 147,\\ a_{11} = a_{12} = a_{13} = 0.\end{cases}$$

Combining (6.3.8.2), (6.3.8.3) and (6.3.8.4), we obtain $\gamma_j = 0$ for all j. The proposition follows. □

Remark 6.3.9. The element $[\theta]$ defined in the relation (6.3.6.6) is an $GL_4(\mathbb{F}_2)$-invariant of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{17}$.

6.4. **The case $t = 3$.**

From the results in Kameko [8], we see that $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+4}+2^{s+1}-3} = 13$ with a basis consisting of all the classes $w_{3,s,j}, 1 \leqslant j \leqslant 13$, represented by the following monomials:

1. $(2^s - 1, 2^s - 1, 16.2^s - 1)$,
2. $(2^s - 1, 16.2^s - 1, 2^s - 1)$,
3. $(16.2^s - 1, 2^s - 1, 2^s - 1)$,
4. $(2^s - 1, 2.2^s - 1, 15.2^s - 1)$,
5. $(2.2^s - 1, 2^s - 1, 15.2^s - 1)$,
6. $(2.2^s - 1, 15.2^s - 1, 2^s - 1)$,
7. $(2.2^s - 1, 3.2^s - 1, 13.2^s - 1)$,
8. $(2.2^s - 1, 8.2^s - 1, 8.2^s - 1)$,
9. $(8.2^s - 1, 2.2^s - 1, 8.2^s - 1)$,
10. $(8.2^s - 1, 8.2^s - 1, 2.2^s - 1)$,
11. $(4.2^s - 1, 6.2^s - 1, 8.2^s - 1)$,
12. $(4.2^s - 1, 8.2^s - 1, 6.2^s - 1)$,
13. $(8.2^s - 1, 4.2^s - 1, 6.2^s - 1)$.

Hence, we easily obtain

Proposition 6.4.1. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+4}+2^{s+1}-3}$ is an $\mathbb{F}_2$-vector space of dimension 52 with a basis consisting of all the classes represented by the monomials $a_{3,s,j}, 1 \leqslant j \leqslant 52$, which are determined as follows:*

1. $(0, 2.2^s - 1, 8.2^s - 1, 8.2^s - 1)$,
2. $(0, 8.2^s - 1, 2.2^s - 1, 8.2^s - 1)$,
3. $(0, 8.2^s - 1, 8.2^s - 1, 2.2^s - 1)$,
4. $(2.2^s - 1, 0, 8.2^s - 1, 8.2^s - 1)$,
5. $(2.2^s - 1, 8.2^s - 1, 0, 8.2^s - 1)$,
6. $(2.2^s - 1, 8.2^s - 1, 8.2^s - 1, 0)$,
7. $(8.2^s - 1, 0, 2.2^s - 1, 8.2^s - 1)$,
8. $(8.2^s - 1, 0, 8.2^s - 1, 2.2^s - 1)$,
9. $(8.2^s - 1, 2.2^s - 1, 0, 8.2^s - 1)$,
10. $(8.2^s - 1, 2.2^s - 1, 8.2^s - 1, 0)$,
11. $(8.2^s - 1, 8.2^s - 1, 0, 2.2^s - 1)$,
12. $(8.2^s - 1, 8.2^s - 1, 2.2^s - 1, 0)$,
13. $(0, 4.2^s - 1, 6.2^s - 1, 8.2^s - 1)$,
14. $(0, 4.2^s - 1, 8.2^s - 1, 6.2^s - 1)$,
15. $(0, 8.2^s - 1, 4.2^s - 1, 6.2^s - 1)$,
16. $(4.2^s - 1, 0, 6.2^s - 1, 8.2^s - 1)$,
17. $(4.2^s - 1, 0, 8.2^s - 1, 6.2^s - 1)$,
18. $(4.2^s - 1, 6.2^s - 1, 0, 8.2^s - 1)$,
19. $(4.2^s - 1, 6.2^s - 1, 8.2^s - 1, 0)$,
20. $(4.2^s - 1, 8.2^s - 1, 0, 6.2^s - 1)$,
21. $(4.2^s - 1, 8.2^s - 1, 6.2^s - 1, 0)$,
22. $(8.2^s - 1, 0, 4.2^s - 1, 6.2^s - 1)$,
23. $(8.2^s - 1, 4.2^s - 1, 0, 6.2^s - 1)$,
24. $(8.2^s - 1, 4.2^s - 1, 6.2^s - 1, 0)$,
25. $(0, 2^s - 1, 2^s - 1, 16.2^s - 1)$,
26. $(0, 2^s - 1, 16.2^s - 1, 2^s - 1)$,
27. $(0, 16.2^s - 1, 2^s - 1, 2^s - 1)$,
28. $(2^s - 1, 0, 2^s - 1, 16.2^s - 1)$,
29. $(2^s - 1, 0, 16.2^s - 1, 2^s - 1)$,
30. $(2^s - 1, 2^s - 1, 0, 16.2^s - 1)$,
31. $(2^s - 1, 2^s - 1, 16.2^s - 1, 0)$,
32. $(2^s - 1, 16.2^s - 1, 0, 2^s - 1)$,
33. $(2^s - 1, 16.2^s - 1, 2^s - 1, 0)$,
34. $(16.2^s - 1, 0, 2^s - 1, 2^s - 1)$,
35. $(16.2^s - 1, 2^s - 1, 0, 2^s - 1)$,
36. $(16.2^s - 1, 2^s - 1, 2^s - 1, 0)$,
37. $(0, 2^s - 1, 2.2^s - 1, 15.2^s - 1)$,
38. $(0, 2.2^s - 1, 2^s - 1, 15.2^s - 1)$,
39. $(0, 2.2^s - 1, 15.2^s - 1, 2^s - 1)$,
40. $(2^s - 1, 0, 2.2^s - 1, 15.2^s - 1)$,
41. $(2^s - 1, 2.2^s - 1, 0, 15.2^s - 1)$,
42. $(2^s - 1, 2.2^s - 1, 15.2^s - 1, 0)$,
43. $(2.2^s - 1, 0, 2^s - 1, 15.2^s - 1)$,
44. $(2.2^s - 1, 0, 15.2^s - 1, 2^s - 1)$,
45. $(2.2^s - 1, 2^s - 1, 0, 15.2^s - 1)$,
46. $(2.2^s - 1, 2^s - 1, 15.2^s - 1, 0)$,
47. $(2.2^s - 1, 15.2^s - 1, 0, 2^s - 1)$,
48. $(2.2^s - 1, 15.2^s - 1, 2^s - 1, 0)$,
49. $(0, 2.2^s - 1, 3.2^s - 1, 13.2^s - 1)$,
50. $(2.2^s - 1, 0, 3.2^s - 1, 13.2^s - 1)$,
51. $(2.2^s - 1, 3.2^s - 1, 0, 13.2^s - 1)$,
52. $(2.2^s - 1, 3.2^s - 1, 13.2^s - 1, 0)$.

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+4}+2^{s+1}-3}$.

Set $\mu_4(1) = 136, \mu_4(2) = 180$ and $\mu_4(s) = 195$ for $s \geqslant 3$.

Theorem 6.4.2. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+4}+2^{s+1}-3}$ is an $\mathbb{F}_2$-vector space of dimension $\mu_3(s) - 52$ with a basis consisting of all the classes represented by the monomials $a_{3,s,j}, 53 \leqslant j \leqslant \mu_4(s)$, which are determined as follows:*

For $s \geqslant 1$,

53. $(1, 2.2^s - 2, 8.2^s - 1, 8.2^s - 1)$,
54. $(1, 8.2^s - 1, 2.2^s - 2, 8.2^s - 1)$,
55. $(1, 8.2^s - 1, 8.2^s - 1, 2.2^s - 2)$,
56. $(8.2^s - 1, 1, 2.2^s - 2, 8.2^s - 1)$,
57. $(8.2^s - 1, 1, 8.2^s - 1, 2.2^s - 2)$,
58. $(8.2^s - 1, 8.2^s - 1, 1, 2.2^s - 2)$,
59. $(1, 2.2^s - 1, 8.2^s - 2, 8.2^s - 1)$,
60. $(1, 2.2^s - 1, 8.2^s - 1, 8.2^s - 2)$,
61. $(1, 8.2^s - 2, 2.2^s - 1, 8.2^s - 1)$,
62. $(1, 8.2^s - 2, 8.2^s - 1, 2.2^s - 1)$,
63. $(1, 8.2^s - 1, 2.2^s - 1, 8.2^s - 2)$,
64. $(1, 8.2^s - 1, 8.2^s - 2, 2.2^s - 1)$,
65. $(2.2^s - 1, 1, 8.2^s - 2, 8.2^s - 1)$,
66. $(2.2^s - 1, 1, 8.2^s - 1, 8.2^s - 2)$,
67. $(2.2^s - 1, 8.2^s - 1, 1, 8.2^s - 2)$,
68. $(8.2^s - 1, 1, 2.2^s - 1, 8.2^s - 2)$,
69. $(8.2^s - 1, 1, 8.2^s - 2, 2.2^s - 1)$,
70. $(8.2^s - 1, 2.2^s - 1, 1, 8.2^s - 2)$,
71. $(1, 4.2^s - 2, 6.2^s - 1, 8.2^s - 1)$,
72. $(1, 4.2^s - 2, 8.2^s - 1, 6.2^s - 1)$,
73. $(1, 8.2^s - 1, 4.2^s - 2, 6.2^s - 1)$,
74. $(8.2^s - 1, 1, 4.2^s - 2, 6.2^s - 1)$,
75. $(1, 4.2^s - 1, 6.2^s - 2, 8.2^s - 1)$,
76. $(1, 4.2^s - 1, 8.2^s - 1, 6.2^s - 2)$,
77. $(1, 8.2^s - 1, 4.2^s - 1, 6.2^s - 2)$,
78. $(4.2^s - 1, 1, 6.2^s - 2, 8.2^s - 1)$,
79. $(4.2^s - 1, 1, 8.2^s - 1, 6.2^s - 2)$,
80. $(4.2^s - 1, 8.2^s - 1, 1, 6.2^s - 2)$,
81. $(8.2^s - 1, 1, 4.2^s - 1, 6.2^s - 2)$,
82. $(8.2^s - 1, 4.2^s - 1, 1, 6.2^s - 2)$,
83. $(1, 4.2^s - 1, 6.2^s - 1, 8.2^s - 2)$,
84. $(1, 4.2^s - 1, 8.2^s - 2, 6.2^s - 1)$,
85. $(1, 8.2^s - 2, 4.2^s - 1, 6.2^s - 1)$,
86. $(4.2^s - 1, 1, 6.2^s - 1, 8.2^s - 2)$,
87. $(4.2^s - 1, 1, 8.2^s - 2, 6.2^s - 1)$,
88. $(4.2^s - 1, 6.2^s - 1, 1, 8.2^s - 2)$,
89. $(3, 8.2^s - 3, 2.2^s - 2, 8.2^s - 1)$,
90. $(3, 8.2^s - 3, 8.2^s - 1, 2.2^s - 2)$,
91. $(3, 8.2^s - 1, 8.2^s - 3, 2.2^s - 2)$,
92. $(8.2^s - 1, 3, 8.2^s - 3, 2.2^s - 2)$,
93. $(3, 4.2^s - 3, 6.2^s - 2, 8.2^s - 1)$,
94. $(3, 4.2^s - 3, 8.2^s - 1, 6.2^s - 2)$,
95. $(3, 8.2^s - 1, 4.2^s - 3, 6.2^s - 2)$,
96. $(8.2^s - 1, 3, 4.2^s - 3, 6.2^s - 2)$,
97. $(3, 4.2^s - 3, 6.2^s - 1, 8.2^s - 2)$,
98. $(3, 4.2^s - 3, 8.2^s - 2, 6.2^s - 1)$,
99. $(3, 8.2^s - 3, 4.2^s - 2, 6.2^s - 1)$,
100. $(3, 4.2^s - 1, 6.2^s - 3, 8.2^s - 2)$,
101. $(4.2^s - 1, 3, 6.2^s - 3, 8.2^s - 2)$,
102. $(3, 4.2^s - 1, 8.2^s - 3, 6.2^s - 2)$,
103. $(3, 8.2^s - 3, 4.2^s - 1, 6.2^s - 2)$,
104. $(4.2^s - 1, 3, 8.2^s - 3, 6.2^s - 2)$,
105. $(7, 8.2^s - 5, 8.2^s - 3, 2.2^s - 2)$,
106. $(7, 8.2^s - 5, 4.2^s - 3, 6.2^s - 2)$,
107. $(1, 2.2^s - 2, 3.2^s - 1, 13.2^s - 1)$,
108. $(1, 2.2^s - 1, 3.2^s - 2, 13.2^s - 1)$,
109. $(2.2^s - 1, 1, 3.2^s - 2, 13.2^s - 1)$,
110. $(1, 2.2^s - 1, 3.2^s - 1, 13.2^s - 2)$,
111. $(2.2^s - 1, 1, 3.2^s - 1, 13.2^s - 2)$,
112. $(2.2^s - 1, 3.2^s - 1, 1, 13.2^s - 2)$,
113. $(3, 2.2^s - 1, 8.2^s - 3, 8.2^s - 2)$,
114. $(3, 8.2^s - 3, 2.2^s - 1, 8.2^s - 2)$,
115. $(3, 8.2^s - 3, 8.2^s - 2, 2.2^s - 1)$,
116. $(1, 2^s - 1, 2.2^s - 2, 15.2^s - 1)$,
117. $(1, 2.2^s - 2, 2^s - 1, 15.2^s - 1)$,
118. $(1, 2.2^s - 2, 15.2^s - 1, 2^s - 1)$,
119. $(1, 2^s - 1, 2^s - 1, 16.2^s - 2)$,
120. $(1, 2^s - 1, 16.2^s - 2, 2^s - 1)$,
121. $(1, 16.2^s - 2, 2^s - 1, 2^s - 1)$.

For $s = 1$,

122. $(3,3,12,15)$, 123. $(3,3,15,12)$, 124. $(3,15,3,12)$,
125. $(15,3,3,12)$, 126. $(3,7,11,12)$, 127. $(7,3,11,12)$,
128. $(7,11,3,12)$, 129. $(7,7,8,11)$, 130. $(7,7,11,8)$,
131. $(7,7,9,10)$, 132. $(1,1,3,28)$, 133. $(1,3,1,28)$,
134. $(1,3,28,1)$, 135. $(3,1,1,28)$, 136. $(3,1,28,1)$.

For $s \geqslant 2$,

122. $(2^s-1,1,2^s-1,16.2^s-2)$, 123. $(2^s-1,1,16.2^s-2,2^s-1)$,
124. $(2^s-1,2^s-1,1,16.2^s-2)$, 125. $(2.2^s-1,3,8.2^s-3,8.2^s-2)$,
126. $(2^s-1,1,2.2^s-2,15.2^s-1)$, 127. $(3,2.2^s-3,8.2^s-2,8.2^s-1)$,
128. $(3,2.2^s-3,8.2^s-1,8.2^s-2)$, 129. $(3,8.2^s-1,2.2^s-3,8.2^s-2)$,
130. $(8.2^s-1,3,2.2^s-3,8.2^s-2)$, 131. $(7,8.2^s-5,2.2^s-3,8.2^s-2)$,
132. $(7,2.2^s-5,8.2^s-3,8.2^s-2)$, 133. $(7,4.2^s-5,6.2^s-3,8.2^s-2)$,
134. $(7,4.2^s-5,8.2^s-3,6.2^s-2)$, 135. $(1,2^s-2,2^s-1,16.2^s-1)$,
136. $(1,2^s-2,16.2^s-1,2^s-1)$, 137. $(1,2^s-1,2^s-2,16.2^s-1)$,
138. $(1,2^s-1,16.2^s-1,2^s-2)$, 139. $(1,16.2^s-1,2^s-2,2^s-1)$,
140. $(1,16.2^s-1,2^s-1,2^s-2)$, 141. $(2^s-1,1,2^s-2,16.2^s-1)$,
142. $(2^s-1,1,16.2^s-1,2^s-2)$, 143. $(2^s-1,16.2^s-1,1,2^s-2)$,
144. $(16.2^s-1,1,2^s-2,2^s-1)$, 145. $(16.2^s-1,1,2^s-1,2^s-2)$,
146. $(16.2^s-1,2^s-1,1,2^s-2)$, 147. $(1,2^s-2,2.2^s-1,15.2^s-1)$,
148. $(1,2.2^s-1,2^s-2,15.2^s-1)$, 149. $(1,2.2^s-1,15.2^s-1,2^s-2)$,
150. $(2.2^s-1,1,2^s-2,15.2^s-1)$, 151. $(2.2^s-1,1,15.2^s-1,2^s-2)$,
152. $(2.2^s-1,15.2^s-1,1,2^s-2)$, 153. $(1,2^s-1,2.2^s-1,15.2^s-2)$,
154. $(1,2.2^s-1,2^s-1,15.2^s-2)$, 155. $(1,2.2^s-1,15.2^s-2,2^s-1)$,
156. $(2^s-1,1,2.2^s-1,15.2^s-2)$, 157. $(2^s-1,2.2^s-1,1,15.2^s-2)$,
158. $(2.2^s-1,1,2^s-1,15.2^s-2)$, 159. $(2.2^s-1,1,15.2^s-2,2^s-1)$,
160. $(2.2^s-1,2^s-1,1,15.2^s-2)$, 161. $(3,2.2^s-3,2^s-2,15.2^s-1)$,
162. $(3,2.2^s-3,15.2^s-1,2^s-2)$, 163. $(3,2.2^s-1,15.2^s-3,2^s-2)$,
164. $(2.2^s-1,3,15.2^s-3,2^s-2)$, 165. $(3,2.2^s-3,3.2^s-2,13.2^s-1)$,
166. $(3,2.2^s-3,3.2^s-1,13.2^s-2)$, 167. $(3,2.2^s-1,3.2^s-3,13.2^s-2)$,
168. $(2.2^s-1,3,3.2^s-3,13.2^s-2)$, 169. $(3,2^s-1,16.2^s-3,2^s-2)$,
170. $(3,16.2^s-3,2^s-2,2^s-1)$, 171. $(3,16.2^s-3,2^s-1,2^s-2)$,
172. $(3,2^s-1,2.2^s-3,15.2^s-2)$, 173. $(3,2.2^s-3,2^s-1,15.2^s-2)$,
174. $(3,2.2^s-3,15.2^s-2,2^s-1)$.

For $s = 2$,

175. $(3,3,3,60)$, 176. $(3,3,60,3)$, 177. $(3,3,4,59)$,
178. $(3,3,7,56)$, 179. $(3,7,3,56)$, 180. $(7,3,3,56)$.

For $s \geqslant 3$

175. $(2^s-1,3,2.2^s-3,15.2^s-2)$, 176. $(2^s-1,3,16.2^s-3,2^s-2)$,
177. $(3,2^s-3,2^s-2,16.2^s-1)$, 178. $(3,2^s-3,16.2^s-1,2^s-2)$,
179. $(3,16.2^s-1,2^s-3,2^s-2)$, 180. $(16.2^s-1,3,2^s-3,2^s-2)$,
181. $(3,2^s-3,2^s-1,16.2^s-2)$, 182. $(3,2^s-3,16.2^s-2,2^s-1)$,
183. $(3,2^s-1,2^s-3,16.2^s-2)$, 184. $(2^s-1,3,2^s-3,16.2^s-2)$,
185. $(3,2^s-3,2.2^s-2,15.2^s-1)$, 186. $(3,2^s-3,2.2^s-1,15.2^s-2)$,

187. $(3, 2.2^s-1, 2^s-3, 15.2^s-2)$, 188. $(2.2^s-1, 3, 2^s-3, 15.2^s-2)$,
189. $(7, 16.2^s-5, 2^s-3, 2^s-2)$, 190. $(7, 2.2^s-5, 2^s-3, 15.2^s-2)$,
191. $(7, 2.2^s-5, 15.2^s-3, 2^s-2)$, 192. $(7, 2.2^s-5, 3.2^s-3, 13.2^s-2)$.

For $s = 3$,

193. $(7, 7, 121, 6)$, 194. $(7, 7, 9, 118)$, 195. $(7, 7, 7, 120)$.

For $s \geqslant 4$,

193. $(7, 2^s-5, 16.2^s-3, 2^s-2)$, 194. $(7, 2^s-5, 2.2^s-3, 15.2^s-2)$,
195. $(7, 2^s-5, 2^s-3, 16.2^s-2)$.

We prove the theorem by proving the following propositions.

Proposition 6.4.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+4}+2^{s+1}-3}$ is generated by the $\mu_4(s) - 52$ elements listed in Theorem 6.4.2.*

We need the following for the proof of this proposition.

Lemma 6.4.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (3,5,24,1), (3,5,9,16), (3,5,8,17). By a direct computation we have

$$\begin{aligned}(3,5,24,1) &= Sq^1(3,3,25,1) + Sq^2\big((5,3,22,1) + (2,3,25,1)\\ &\quad + (5,3,21,2)\big) + Sq^4\big((3,3,22,1) + (3,3,21,2)\\ &\quad + (3,9,13,4)\big) + Sq^8(3,5,13,4) + (3,4,25,1)\\ &\quad + (2,5,25,1) + (3,5,17,8) \mod \mathcal{L}_4(3;1;1;1;1),\\ (3,5,9,16) &= Sq^1(3,3,9,17) + Sq^2\big((5,3,9,14) + (2,3,9,17)\\ &\quad + (5,3,6,17)\big) + Sq^4\big((3,3,9,14) + (3,3,6,17)\big)\\ &\quad + (3,4,9,17) + (2,5,9,17) + (3,5,8,17) \mod \mathcal{L}_4(3;1;1;1;1),\\ (3,5,8,17) &= Sq^1(5,9,1,17) + Sq^2\big((3,9,2,17) + (2,11,1,17)\\ &\quad + (6,7,1,17) + (5,10,1,15) + (3,10,1,17) + (5,6,1,19)\\ &\quad + (3,9,1,18)\big) + Sq^4\big((3,5,4,17) + (4,7,1,17) + (3,10,1,15)\\ &\quad + (3,6,1,19) + (3,4,1,21) + (3,5,1,20)\big) + (2,13,1,17)\\ &\quad + (3,4,1,25) + (3,5,1,24) \mod \mathcal{L}_4(3;1;1;1;1),\end{aligned}$$

The lemma is proved. □

Lemma 6.4.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&1&0&1\\1&0&1&1\\1&0&0&1\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&0&1&1\\1&0&1&0\\0&1&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&1\\1&1&0&0\\0&1&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&1\\1&1&0&0\\1&0&1&0\end{pmatrix}$$

$$\begin{pmatrix}1&1&0&1\\1&1&0&1\\0&1&0&1\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\0&1&1&0\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\0&1&1&0\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&0&0&1\\0&0&1&1\end{pmatrix}$$

$$\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&0&1&0\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&1&1\\1&0&0&1\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&1&1\\1&0&1&0\\0&1&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&0&0\\0&1&0&1\end{pmatrix}$$

$$\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&1&0&0\\0&1&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&0&1&0\\1&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&0&0\\1&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&1&0&0\\1&0&1&0\end{pmatrix}$$

$$\begin{pmatrix}1&1&1&0\\1&0&1&1\\1&0&0&1\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&0&1&1\\1&0&1&0\\0&1&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&1\\1&0&1&0\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&0&1&1\\1&0&1&0\\0&1&0&1\end{pmatrix}$$

$$\begin{pmatrix}1&0&1&1\\1&0&1&1\\1&0&1&0\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&0&1&0\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&0&1&1\\1&0&1&0\\0&1&0&1\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(7,9,2,15),(7,9,15,2),(7,15,9,2),(15,7,9,2),(3,7,8,15),(3,7,15,8),$$
$$(3,15,7,8),(7,3,8,15),(7,3,15,8),(7,8,3,15),(7,8,15,3),(7,15,3,8),$$
$$(7,15,8,3),(15,3,7,8),(15,7,3,8),(15,7,8,3),(7,9,3,14),(7,9,14,3),$$
$$(7,11,5,10),(7,9,6,11),(7,8,7,11),(7,11,7,8),(7,9,7,10).$$

We prove the lemma the matrices corresponding to the monomials

$$(7,9,2,15),(7,3,8,15),(7,9,3,14),(7,9,6,11),$$
$$(7,8,7,11),(7,11,7,8),(7,9,7,10).$$

The others can be obtained by a similar argument. We have

$$\begin{aligned}
(7,9,2,15) &= Sq^1(7,7,3,15) + Sq^2(7,7,2,15)\\
&\quad + Sq^4\big((5,7,2,15) + (4,7,3,15)\big) + (5,11,2,15)\\
&\quad + (4,11,3,15) + (7,8,3,15) \bmod \mathcal{L}_4(3;3;2;2),\\
(7,3,8,15) &= Sq^1(7,3,7,15) + Sq^2(7,2,7,15)\\
&\quad + Sq^4\big((4,3,7,15) + (5,2,7,15)\big) + (4,3,11,15)\\
&\quad + (7,2,9,15) + (5,2,11,15) \mod \mathcal{L}_4(3;3;2;2),\\
(7,9,3,14) &= Sq^1\big((7,7,11,7) + (7,7,5,13)\big) + Sq^2\big((7,7,10,7)\\
&\quad + (7,7,6,11) + (7,7,3,14)\big) + Sq^4\big((4,7,11,7) + (5,7,10,7)\\
&\quad + (11,5,6,7) + (5,7,6,11) + (5,7,3,14)\big) + Sq^8(7,5,6,7)\\
&\quad + (4,11,11,7) + (4,7,11,11) + (7,8,11,7) + (7,7,10,9) + (5,11,10,7)\\
&\quad + (5,7,10,11) + (7,5,10,11) + (7,7,8,11) + (7,7,11,8)\\
&\quad + (5,11,6,11) + (5,7,10,11) + (5,11,3,14) \mod \mathcal{L}_4(3;3;2;2),\\
(7,9,6,11) &= Sq^1(7,7,5,13) + Sq^2\big((7,7,6,11) + (7,7,3,14)\big)\\
&\quad + Sq^4\big((5,7,6,11) + (5,7,3,14)\big) + (5,11,6,11) + (5,7,10,11)\\
&\quad + (7,7,8,11) + (7,9,3,14) + (5,11,3,14) \mod \mathcal{L}_4(3;3;2;2),\\
(7,8,7,11) &= Sq^1(7,5,7,13) + Sq^2\big((7,6,7,11) + (7,3,7,14)\big)\\
&\quad + Sq^4\big((5,6,7,11) + (5,3,7,14)\big) + (5,6,11,11) + (5,10,7,11)\\
&\quad + (7,6,9,11) + (7,3,9,14) + (5,3,11,14) \mod \mathcal{L}_4(3;3;2;2),\\
(7,11,7,8) &= Sq^1(7,11,7,7) + Sq^2(7,10,7,7) + Sq^4\big((4,11,7,7)\\
&\quad + (5,10,7,7) + (11,7,4,7)\big) + Sq^8(7,7,4,7) + (4,11,11,7)\\
&\quad + (4,11,7,11) + (7,10,9,7) + (7,10,7,9) + (5,10,11,7)\\
&\quad + (5,10,7,11) + (7,11,4,11) + (7,7,8,11) \mod \mathcal{L}_4(3;3;2;2),\\
(7,9,7,10) &= Sq^1(7,7,7,11) + Sq^2(7,7,7,10) + Sq^4\big((4,7,7,11)\\
&\quad + (5,7,7,10)\big) + (4,11,7,11) + (4,7,11,11)\\
&\quad + (7,7,9,10) + (5,10,7,9) + (5,11,7,10) + (5,7,11,10)\\
&\quad + (7,7,8,11) + (7,8,7,11) \bmod \mathcal{L}_4(3;3;2;2).
\end{aligned}$$

The lemma is proved. □

Lemma 6.4.6. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1&1&0&1\\ 1&1&0&1\\ 1&1&0&1\\ 1&1&0&0\\ 0&0&1&1 \end{pmatrix} \quad \begin{pmatrix} 1&1&1&0\\ 1&1&1&0\\ 1&1&1&0\\ 1&1&0&0\\ 0&0&1&1 \end{pmatrix} \quad \begin{pmatrix} 1&1&1&0\\ 1&1&0&1\\ 1&1&0&1\\ 1&1&0&0\\ 0&0&1&1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (15,15,16,23), (15,15,23,16), (15,15,17,22). We prove the lemma for the monomials (15,15,16,23), (15,15,17,22). We have

$$\begin{aligned}
(15,15,16,23) &= Sq^1\big((15,15,13,25)+(15,15,11,27)+(15,15,9,29)\big)\\
&\quad + Sq^2\big((15,15,14,23)+(15,11,14,27)+(15,15,11,26)\\
&\quad + (15,15,10,27)+(15,15,7,30)+(15,11,11,30)\big)\\
&\quad + Sq^4\big((15,13,14,23)+(15,7,14,29)+(15,13,7,30)+(15,7,13,30)\big)\\
&\quad + Sq^8\big((9,15,14,23)+(11,13,14,23)+(11,7,14,29)\\
&\quad + (9,15,7,30)+(11,13,7,30)+(11,7,13,30)\big)+(9,23,14,23)\\
&\quad + (9,15,22,23)+(15,13,18,23)+(11,21,14,23)+(11,13,22,23)\\
&\quad + (11,7,22,29)+(9,23,7,30)+(11,21,7,30)+(15,7,17,30)\\
&\quad + (11,7,21,30)+(15,7,18,29) \mod \mathcal{L}_4(3;3;3;2;2),\\
(15,15,17,22) &= Sq^1\big((15,15,15,23)+(15,11,15,27)\big)\\
&\quad + Sq^2\big((15,15,15,22)+(15,11,15,26)\big)+Sq^4\big((15,13,15,22)\\
&\quad + (15,12,15,23)\big)+Sq^8\big((9,15,15,22)+(11,13,15,22)+(8,15,15,23)\\
&\quad + (11,12,15,23)\big)+(9,23,15,22)+(9,15,23,22)+(11,21,15,22)\\
&\quad + (11,13,23,22)+(15,13,19,22)+(8,23,15,23)\\
&\quad + (8,15,23,23)+(15,15,16,23)+(11,20,15,23)\\
&\quad + (11,12,23,23)+(15,12,19,23) \mod \mathcal{L}_4(3;3;3;2;2).
\end{aligned}$$

The lemma follows. □

Combining Theorem 2.5, the lemmas in Sections 3, 4, 5 and Lemmas 6.1.1, 6.2.4, 6.2.5, 6.3.4, 6.3.5, 6.4.4, 6.4.5, 6.4.6, we get Proposition 6.4.3.

Now, we show that the monomials $a_{3,s,j}, 53 \leqslant j \leqslant \mu_4(s)$ are admissible.

Proposition 6.4.7. *The elements* $[a_{3,1,j}]$, $53 \leqslant j \leqslant 136$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{33}$.

Proof. Suppose there is a linear relation

$$\sum_{j=53}^{136} \gamma_j [a_{3,1,j}] = 0, \tag{6.4.7.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Consider the homomorphisms $f_i,\ i = 1, 2, 3$. Under these homomorphisms, the images of the above relation respectively are

$$\begin{aligned}
&\gamma_{121}[31,1,1] + \gamma_{117}[3,1,29] + \gamma_{118}[3,29,1] + \gamma_{107}[3,5,25]\\
&\quad + \gamma_{53}[3,15,15] + \gamma_{61}[15,3,15] + \gamma_{62}[15,15,3]\\
&\quad + \gamma_{71}[7,11,15] + \gamma_{72}[7,15,11] + \gamma_{85}[15,7,11] = 0,\\
&\gamma_{\{120,136\}}[31,1,1] + a_1[3,1,29] + \gamma_{134}[3,29,1]\\
&\quad + a_2[3,5,25] + \gamma_{54}[3,15,15] + \gamma_{\{59,122\}}[15,3,15] + \gamma_{64}[15,15,3]\\
&\quad + \gamma_{75}[7,11,15] + \gamma_{73}[7,15,11] + \gamma_{\{84,129\}}[15,7,11] = 0,\\
&\gamma_{\{119,135\}}[31,1,1] + a_3[3,1,29] + a_4[3,29,1] + a_5[3,5,25]\\
&\quad + \gamma_{55}[3,15,15] + \gamma_{\{60,123\}}[15,3,15] + \gamma_{\{63,124\}}[15,15,3]\\
&\quad + \gamma_{76}[7,11,15] + \gamma_{77}[7,15,11] + \gamma_{\{83,126,130\}}[15,7,11] = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{56,65,69,74,78,87,109,116\}}, \quad a_2 = \gamma_{\{89,93,98,99,108,115\}},\\
&a_3 = \gamma_{\{57,66,68,79,81,86,111,132\}}, \quad a_4 = \gamma_{\{58,67,70,80,82,88,112,133\}},\\
&a_5 = \gamma_{\{90,91,92,94,95,96,97,100,101,102,103,104,105,106,110,113,114,125,127,128,131\}}.
\end{aligned}$$

From this, it implies

$$\text{(6.4.7.2)} \qquad \begin{cases}
a_1 = a_2 = a_3 = a_4 = a_5 = 0,\\
\gamma_j = 0,\ j = 53, 54, 55, 61, 62, 64, 71, 72,\\
73, 75, 76, 77, 85, 107, 117, 118, 121, 134,\\
\gamma_{\{120,136\}} = \gamma_{\{59,122\}} = \gamma_{\{84,129\}} = \gamma_{\{119,135\}} = 0,\\
\gamma_{\{60,123\}} = \gamma_{\{63,124\}} = \gamma_{\{83,126,130\}} = 0.
\end{cases}$$

With the aid of (6.4.7.2), the homomorphisms f_4, f_5, f_6 send (6.4.7.1) respectively to

$$\begin{aligned}
&\gamma_{120}[1,31,1] + \gamma_{\{59,84,108,116\}}[1,3,29] + \gamma_{136}[3,29,1]\\
&\quad + \gamma_{\{98,99,109,115\}}[3,5,25] + \gamma_{\{65,89,93,122\}}[3,15,15] + \gamma_{56}[15,3,15]\\
&\quad + \gamma_{69}[15,15,3] + \gamma_{78}[7,11,15] + \gamma_{\{87,129\}}[7,15,11] + \gamma_{74}[15,7,11] = 0,\\
&\gamma_{\{119,133\}}[1,31,1] + \gamma_{\{60,63,83,110,132\}}[1,3,29] + a_6[3,29,1] + a_7[3,5,25]\\
&\quad + \gamma_{\{66,90,94,123\}}[3,15,15] + \gamma_{57}[15,3,15] + \gamma_{\{68,125\}}[15,15,3]\\
&\quad + \gamma_{79}[7,11,15] + \gamma_{\{86,127,130\}}[7,15,11] + \gamma_{81}[15,7,11] = 0,\\
&\gamma_{\{116,119,120,132\}}[1,1,31] + a_8[1,3,29] + a_9[3,1,29] + a_{10}[3,5,25]\\
&\quad + \gamma_{\{67,91,95,124\}}[3,15,15] + \gamma_{\{70,92,96,125\}}[15,3,15] + \gamma_{58}[15,15,3]\\
&\quad + \gamma_{\{88,105,106,128\}}[7,11,15] + \gamma_{80}[7,15,11] + \gamma_{82}[15,7,11] = 0,
\end{aligned}$$

where

$$a_6 = \gamma_{\{58,67,70,80,82,88,112,135\}}, \quad a_8 = \gamma_{\{59,60,63,83,84,108,110,133\}},$$

$$a_7 = \gamma_{\{91,92,95,96,97,100,101,102,103,104,105,106,111,113,114,124,126,128,131\}},$$
$$a_9 = \gamma_{\{56,57,65,66,68,69,74,78,79,81,86,87,109,111,135,136\}},$$
$$a_{10} = \gamma_{\{89,90,93,94,97,98,99,100,101,102,103,104,112,113,114,115,122,123,126,127,129,130,131\}}.$$

From this, it implies

$$\text{(6.4.7.3)}\qquad \begin{cases} a_6 = a_7 = a_8 = a_9 = a_{10} = 0, \\ \gamma_j = 0,\ j = 56, 57, 58, 69, 74, 78, 79, 80, 81, 82, 120, 136, \\ \gamma_{\{59,84,108,116\}} = \gamma_{\{98,99,109,115\}} = \gamma_{\{65,89,93,122\}} = 0, \\ \gamma_{\{87,129\}} = \gamma_{\{119,133\}} = \gamma_{\{68,125\}} = \gamma_{\{60,63,83,110,132\}} = 0, \\ \gamma_{\{66,90,94,123\}} = \gamma_{\{86,127,130\}} = \gamma_{\{116,119,120,132\}} = 0, \\ \gamma_{\{67,91,95,124\}} = \gamma_{\{70,92,96,125\}} = \gamma_{\{88,105,106,128\}} = 0. \end{cases}$$

With the aid of (6.4.7.2) and (6.4.7.3), the homomorphisms g_1, g_2 send (6.4.7.1) respectively to

$$\begin{aligned} &\gamma_{\{65,84,109,116\}}[1,3,29] + a_{11}[3,5,25] + \gamma_{65}[3,15,15] + \gamma_{89}[15,3,15] \\ &\quad + \gamma_{115}[15,15,3] + \gamma_{93}[7,11,15] + \gamma_{\{84,98\}}[7,15,11] + \gamma_{99}[15,7,11] = 0, \\ &\gamma_{\{66,68,86,111,132\}}[1,3,29] + a_{12}[3,5,25] + \gamma_{66}[3,15,15] \\ &\quad + \gamma_{90}[15,3,15] + \gamma_{\{114,128\}}[15,15,3] + \gamma_{94}[7,11,15] \\ &\quad + \gamma_{\{83,86,97,126,127,130\}}[7,15,11] + \gamma_{103}[15,7,11] = 0, \end{aligned}$$

where

$$a_{11} = \gamma_{\{59,84,89,93,98,99,108,109,115\}},$$
$$a_{12} = \gamma_{\{60,63,68,90,94,97,103,110,111,114,126,127,128,130\}}.$$

From this, it implies

$$\text{(6.4.7.4)}\qquad \begin{cases} \gamma_j = 0,\ j = 65, 66, 89, 90, 93, 94, 99, 103, 115, \\ a_{11} = a_{12} = \gamma_{\{65,84,109,116\}} = \gamma_{\{84,98\}} = \gamma_{\{114,128\}} = 0, \\ \gamma_{\{66,68,86,111,132\}} = \gamma_{\{83,86,97,126,127,130\}} = 0. \end{cases}$$

With the aid of (6.4.7.2), (6.4.7.3) and (6.4.7.4), the homomorphisms g_3, g_4 send (6.4.7.1) respectively to

$$\begin{aligned} &\gamma_{\{67,70,88,112,119\}}[1,3,29] + \gamma_{\{116,119,132\}}[3,29,1] \\ &\quad + a_{13}[3,5,25] + \gamma_{67}[3,15,15] + \gamma_{91}[15,3,15] + a_{14}[15,15,3] \\ &\quad + \gamma_{95}[7,11,15] + a_{15}[7,15,11] + \gamma_{\{102,105,130,131\}}[15,7,11] = 0, \\ &\gamma_{\{67,70,88,112,119\}}[1,3,29] + \gamma_{\{116,119,132\}}[3,29,1] \\ &\quad + a_{16}[3,5,25] + \gamma_{70}[3,15,15] + \gamma_{92}[15,3,15] + a_{17}[15,15,3] \\ &\quad + \gamma_{96}[7,11,15] + a_{18}[7,15,11] + \gamma_{\{104,105,106,130,131\}}[15,7,11] = 0, \end{aligned}$$

where

$$a_{13} = \gamma_{\{59,60,63,68,91,92,95,96,100,101,102,104,105,106,108,110,112,113,126,127,128\}},$$
$$a_{14} = \gamma_{\{68,70,92,96,101,104,113,127\}},\quad a_{15} = \gamma_{\{83,84,88,100,106,126,128,130,131\}},$$

$a_{16} = \gamma_{\{59,60,63,68,91,92,95,96,97,98,100,101,102,104,105,106,109,111,112,113,114,126,127,128\}}$,

$a_{17} = \gamma_{\{59,60,63,67,91,95,97,98,100,102,113,114,126\}}$, $a_{18} = \gamma_{\{84,86,88,101,127,128,130,131\}}$.

From this, it implies

$$\begin{cases} a_{13} = a_{14} = a_{15} = a_{16} = a_{17} = a_{18} = 0, \\ \gamma_j = 0,\ j = 67, 91, 95, 70, 92, 96, \\ \gamma_{\{67,70,88,112,119\}} = \gamma_{\{116,119,132\}} = \gamma_{\{102,105,130,131\}} = 0, \\ \gamma_{\{116,119,132\}} = \gamma_{\{67,70,88,112,119\}} = \gamma_{\{104,105,106,130,131\}} = 0. \end{cases} \tag{6.4.7.5}$$

With the aid of (6.4.7.2), (6.4.7.3), (6.4.7.4) and (6.4.7.5), the homomorphisms h sends (6.4.7.1) to

$$\begin{aligned} &a_{19}[3,5,25] + a_{20}[3,15,15] + \gamma_{88}[15,3,15] + \gamma_{105}[15,15,3] \\ &\quad + a_{21}[7,11,15] + \gamma_{\{102,104,105,130,131\}}[7,15,11] + \gamma_{106}[15,7,11] = 0, \end{aligned}$$

where

$a_{19} = \gamma_{\{86,88,101,104,105,106,110,111,112,114,127,130,131\}}$,

$a_{20} = \gamma_{\{84,86,101,104,113,127\}}$, $a_{21} = \gamma_{\{83,84,86,88,97,100,101,106,114,126,127,130,131\}}$.

Hence, we obtain

$$\begin{cases} a_{19} = a_{20} = a_{21} = \gamma_{88} = \gamma_{105} = \gamma_{106} = 0, \\ \gamma_{\{102,104,105,130,131\}} = 0. \end{cases} \tag{6.4.7.6}$$

Combining (6.4.7.2), (6.4.7.3), (6.4.7.4), (6.4.7.5) and (6.4.7.6) gives

$$\begin{cases} \gamma_j = 0,\ j = 83, 84, 86, 87, 97, 98, 108, 109, \\ \qquad\qquad 112, 119, 129, 130, 131, 132, 133, 135. \\ \gamma_{83} = \gamma_{84} = \gamma_{86} = \gamma_{87} = \gamma_{97} = \gamma_{98} = \gamma_{108} = \gamma_{109} = \gamma_{112} \\ \qquad = \gamma_{119} = \gamma_{129} = \gamma_{130} = \gamma_{131} = \gamma_{132} = \gamma_{133} = \gamma_{135}. \end{cases} \tag{6.4.7.7}$$

Substituting (6.4.7.7) into the relation (6.4.7.1), we get

$$\gamma_{83}[\theta] = 0, \tag{6.4.7.8}$$

where $\theta = a_{3,1,83} + a_{3,1,84} + a_{3,1,86} + a_{3,1,87} + a_{3,1,97} + a_{3,1,98} + a_{3,1,108} + a_{3,1,109} + a_{3,1,112} + a_{3,1,119} + a_{3,1,129} + a_{3,1,130} + a_{3,1,131} + a_{3,1,132} + a_{3,1,133} + a_{3,1,135}$.

By a direct computation, we can show that the polynomial θ is non-hit. Hence, $[\theta] \neq 0$ and $\gamma_{83} = 0$. The proposition follows. □

Proposition 6.4.8. *The elements* $[a_{3,2,j}]$, $53 \leqslant j \leqslant 180$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{69}$.

Proof. Suppose there is a linear relation

$$\sum_{j=53}^{180} \gamma_j [a_{3,2,j}] = 0, \tag{6.4.8.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms $f_i, 1 \leqslant i \leqslant 6$, to (6.4.8.1) and we get

$$\begin{aligned}
&\gamma_{135}[3,3,63] + \gamma_{136}[3,63,3] + \gamma_{121}[63,3,3] + \gamma_{147}[3,7,59] + \gamma_{117}[7,3,59]\\
&\quad + \gamma_{118}[7,59,3] + \gamma_{107}[7,11,51] + \gamma_{53}[7,31,31] + \gamma_{61}[31,7,31]\\
&\quad + \gamma_{62}[31,31,7] + \gamma_{71}[15,23,31] + \gamma_{72}[15,31,23] + \gamma_{85}[31,15,23] = 0,\\
&\gamma_{137}[3,3,63] + \gamma_{139}[3,63,3] + \gamma_{\{120,176\}}[63,3,3]\\
&\quad + \gamma_{148}[3,7,59] + \gamma_{\{116,177\}}[7,3,59] + \gamma_{155}[7,59,3]\\
&\quad + \gamma_{108}[7,11,51] + \gamma_{54}[7,31,31] + \gamma_{59}[31,7,31] + \gamma_{64}[31,31,7]\\
&\quad + \gamma_{75}[15,23,31] + \gamma_{73}[15,31,23] + \gamma_{84}[31,15,23] = 0,\\
&\gamma_{138}[3,3,63] + \gamma_{140}[3,63,3] + \gamma_{\{119,175,180\}}[63,3,3]\\
&\quad + \gamma_{149}[3,7,59] + \gamma_{\{153,178\}}[7,3,59] + \gamma_{\{154,179\}}[7,59,3]\\
&\quad + \gamma_{110}[7,11,51] + \gamma_{55}[7,31,31] + \gamma_{60}[31,7,31] + \gamma_{63}[31,31,7]\\
&\quad + \gamma_{76}[15,23,31] + \gamma_{77}[15,31,23] + \gamma_{83}[31,15,23] = 0,\\
&\gamma_{141}[3,3,63] + \gamma_{\{123,170,174,176\}}[3,63,3] + \gamma_{144}[63,3,3] + a_1[3,7,59]\\
&\quad + \gamma_{150}[7,3,59] + \gamma_{159}[7,59,3] + \gamma_{109}[7,11,51]\\
&\quad + \gamma_{65}[7,31,31] + \gamma_{56}[31,7,31] + \gamma_{69}[31,31,7]\\
&\quad + \gamma_{78}[15,23,31] + \gamma_{87}[15,31,23] + \gamma_{74}[31,15,23] = 0,\\
&\gamma_{142}[3,3,63] + \gamma_{\{122,171,173,175,179\}}[3,63,3] + \gamma_{145}[63,3,3]\\
&\quad + a_2[3,7,59] + \gamma_{151}[7,3,59] + \gamma_{\{158,180\}}[7,59,3]\\
&\quad + \gamma_{111}[7,11,51] + \gamma_{66}[7,31,31] + \gamma_{57}[31,7,31] + \gamma_{68}[31,31,7]\\
&\quad + \gamma_{79}[15,23,31] + \gamma_{86}[15,31,23] + \gamma_{81}[31,15,23] = 0,\\
&a_3[3,3,63] + \gamma_{143}[3,63,3] + \gamma_{146}[63,3,3] + a_4[3,7,59] + a_5[7,3,59]\\
&\quad + \gamma_{152}[7,59,3] + a_6[7,11,51] + \gamma_{67}[7,31,31] + \gamma_{70}[31,7,31]\\
&\quad + \gamma_{58}[31,31,7] + \gamma_{88}[15,23,31] + \gamma_{80}[15,31,23] + \gamma_{82}[31,15,23] = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{89,93,98,99,115,126,127,161,165,177\}},\ a_2 = \gamma_{\{90,94,97,103,114,128,156,162,166,178\}},\\
&a_3 = \gamma_{\{124,169,172,175,176,177,178\}},\ a_4 = \gamma_{\{91,95,100,102,113,129,157,163,167,179\}},\\
&a_5 = \gamma_{\{92,96,101,104,125,130,160,164,168,180\}},\quad a_6 = \gamma_{\{105,106,112,131,132,133,134\}}.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
a_1 = a_2 = a_3 = a_4 = a_5 = a_6 = 0,\\
\gamma_j = 0,\ j = 53,\ldots,88;\ 107,\ldots,111,\\
117,118,121,135,\ldots,152,155,159,\\
\gamma_{\{120,176\}} = \gamma_{\{116,177\}} = \gamma_{\{119,175,180\}} = \gamma_{\{153,178\}} = 0,\\
\gamma_{\{154,179\}} = \gamma_{\{123,170,174,176\}} = 0,\\
\gamma_{\{122,171,173,175,179\}} = \gamma_{\{158,180\}} = 0.
\end{cases} \tag{6.4.8.2}$$

With the aid of (6.4.8.2), the homomorphisms g_1, g_2 send (6.4.8.1) respectively to

$$\begin{aligned}
&\gamma_{123}[3,63,3]+\gamma_{170}[63,3,3]+\gamma_{126}[3,7,59]+\gamma_{161}[7,3,59]\\
&\quad+\gamma_{174}[7,59,3]+\gamma_{165}[7,11,51]+\gamma_{127}[7,31,31]+\gamma_{89}[31,7,31]\\
&\quad+\gamma_{115}[31,31,7]+\gamma_{93}[15,23,31]+\gamma_{98}[15,31,23]+\gamma_{99}[31,15,23]=0,\\
&\gamma_{\{119,122,175\}}[3,63,3]+\gamma_{171}[63,3,3]+\gamma_{156}[3,7,59]+\gamma_{162}[7,3,59]\\
&\quad+\gamma_{173}[7,59,3]+\gamma_{166}[7,11,51]+\gamma_{128}[7,31,31]+\gamma_{90}[31,7,31]\\
&\quad+\gamma_{114}[31,31,7]+\gamma_{94}[15,23,31]+\gamma_{97}[15,31,23]+\gamma_{103}[31,15,23]=0.
\end{aligned}$$

From these equalities, we obtain

$$\text{(6.4.8.3)}\qquad \begin{cases}\gamma_j=0,\ j=89,90,93,94,97,98,99,103,114,115,123,\\ 126,127,128,156,161,162,165,166,170,171,173,174,\\ \gamma_{\{119,122,175\}}=0.\end{cases}$$

With the aid of (6.4.8.2) and (6.4.8.3), the homomorphisms g_3, g_4 send (6.4.8.1) respectively to

$$\begin{aligned}
&a_7[3,63,3]+\gamma_{\{164,169\}}[63,3,3]+\gamma_{\{112,157\}}[3,7,59]+\gamma_{163}[7,3,59]\\
&\quad+\gamma_{\{158,160,164,172\}}[7,59,3]+\gamma_{\{112,132,167\}}[7,11,51]\\
&\quad+\gamma_{129}[7,31,31]+\gamma_{91}[31,7,31]+\gamma_{\{113,132\}}[31,31,7]\\
&\quad+\gamma_{95}[15,23,31]+\gamma_{100}[15,31,23]+\gamma_{102}[31,15,23]=0,\\
&a_8[3,63,3]+\gamma_{\{120,163,169\}}[63,3,3]+\gamma_{\{112,160\}}[3,7,59]+\gamma_{164}[7,3,59]\\
&\quad+a_9[7,59,3]+a_{10}[7,11,51]+\gamma_{130}[7,31,31]+\gamma_{92}[31,7,31]\\
&\quad+a_{11}[31,31,7]+\gamma_{96}[15,23,31]+\gamma_{101}[15,31,23]+\gamma_{104}[31,15,23]=0,
\end{aligned}$$

where

$$\begin{aligned}
a_7&=\gamma_{\{119,124,158,160,175\}},\quad a_8=\gamma_{\{122,124,154,157,175\}},\\
a_9&=\gamma_{\{116,153,154,157,163,172\}},\quad a_{10}=\gamma_{\{105,106,112,131,132,133,134,168\}},\\
a_{11}&=\gamma_{\{105,106,125,131,132,133,134\}}.
\end{aligned}$$

Computing directly from these equalities we get

$$\text{(6.4.8.4)}\qquad \begin{cases}a_7=a_8=a_9=a_{10}=a_{11},\quad \gamma_j=0,\ j=91,92,95,96,\\ 100,101,102,104,129,130,163,164,\\ \gamma_{\{164,169\}}=\gamma_{\{112,157\}}=\gamma_{\{158,160,164,172\}}=0,\\ \gamma_{\{112,132,167\}}=\gamma_{\{113,132\}}=\gamma_{\{120,163,169\}}=\gamma_{\{112,160\}}=0.\end{cases}$$

With the aid of (6.4.8.2), (6.4.8.3) and (6.4.8.4), the homomorphism h sends (6.4.8.1) to

$$\begin{aligned}
&\gamma_{\{112,119,122,124,175\}}[3,3,63]+\gamma_{\{112,172\}}[3,7,59]+\gamma_{\{112,167,168\}}[7,11,51]\\
&\quad+\gamma_{125}[7,31,31]+\gamma_{131}[31,7,31]+\gamma_{105}[31,31,7]\\
&\quad+\gamma_{133}[15,23,31]+\gamma_{134}[15,31,23]+\gamma_{106}[31,15,23]=0.
\end{aligned}$$

From this, it implies

$$\begin{cases}\gamma_j = 0,\ j = 105, 106, 125, 131, 133, 134,\\ \gamma_{\{112,119,122,124,175\}} = \gamma_{\{112,172\}} = \gamma_{\{112,167,168\}} = 0.\end{cases} \tag{6.4.8.5}$$

Combining (6.4.8.2), (6.4.8.3), (6.4.8.4) and (6.4.8.5), we obtain $\gamma_j = 0$ for all j. The proposition is proved. □

Proposition 6.4.9. *For $s \geqslant 3$, the elements $[a_{3,s,j}]$, $53 \leqslant j \leqslant 195$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+4}+2^{s+1}-3}$.*

Proof. Suppose there is a linear relation

$$\sum_{j=53}^{195} \gamma_j [a_{3,s,j}] = 0, \tag{6.4.9.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_i, 1 \leqslant i \leqslant 6$, to the relation (6.4.9.1), we obtain

$$\begin{aligned}
&\gamma_{135}w_{3,s,1} + \gamma_{136}w_{3,s,2} + \gamma_{121}w_{3,s,3} + \gamma_{147}w_{3,s,4} + \gamma_{117}w_{3,s,5}\\
&\quad + \gamma_{118}w_{3,s,6} + \gamma_{107}w_{3,s,7} + \gamma_{53}w_{3,s,8} + \gamma_{61}w_{3,s,9}\\
&\quad + \gamma_{62}w_{3,s,10} + \gamma_{71}w_{3,s,11} + \gamma_{72}w_{3,s,12} + \gamma_{85}w_{3,s,13} = 0,\\
&\gamma_{137}w_{3,s,1} + \gamma_{139}w_{3,s,2} + \gamma_{120}w_{3,s,3} + \gamma_{148}w_{3,s,4} + \gamma_{116}w_{3,s,5}\\
&\quad + \gamma_{155}w_{3,s,6} + \gamma_{108}w_{3,s,7} + \gamma_{54}w_{3,s,8} + \gamma_{59}w_{3,s,9}\\
&\quad + \gamma_{64}w_{3,s,10} + \gamma_{75}w_{3,s,11} + \gamma_{73}w_{3,s,12} + \gamma_{84}w_{3,s,13} = 0,\\
&\gamma_{138}w_{3,s,1} + \gamma_{140}w_{3,s,2} + a_1w_{3,s,3} + \gamma_{149}w_{3,s,4} + \gamma_{153}w_{3,s,5}\\
&\quad + \gamma_{154}w_{3,s,6} + \gamma_{110}w_{3,s,7} + \gamma_{55}w_{3,s,8} + \gamma_{60}w_{3,s,9}\\
&\quad + \gamma_{63}w_{3,s,10} + \gamma_{76}w_{3,s,11} + \gamma_{77}w_{3,s,12} + \gamma_{83}w_{3,s,13} = 0,\\
&\gamma_{141}w_{3,s,1} + \gamma_{123}w_{3,s,2} + \gamma_{144}w_{3,s,3} + \gamma_{126}w_{3,s,4} + \gamma_{150}w_{3,s,5}\\
&\quad + \gamma_{159}w_{3,s,6} + \gamma_{109}w_{3,s,7} + \gamma_{65}w_{3,s,8} + \gamma_{56}w_{3,s,9}\\
&\quad + \gamma_{69}w_{3,s,10} + \gamma_{78}w_{3,s,11} + \gamma_{87}w_{3,s,12} + \gamma_{74}w_{3,s,13} = 0,\\
&\gamma_{142}w_{3,s,1} + a_2w_{3,s,2} + \gamma_{145}w_{3,s,3} + \gamma_{156}w_{3,s,4} + \gamma_{151}w_{3,s,5}\\
&\quad + \gamma_{158}w_{3,s,6} + \gamma_{111}w_{3,s,7} + \gamma_{66}w_{3,s,8} + \gamma_{57}w_{3,s,9}\\
&\quad + \gamma_{68}w_{3,s,10} + \gamma_{79}w_{3,s,11} + \gamma_{86}w_{3,s,12} + \gamma_{81}w_{3,s,13} = 0,\\
&a_3w_{3,s,1} + \gamma_{143}w_{3,s,2} + \gamma_{146}w_{3,s,3} + \gamma_{157}w_{3,s,4} + \gamma_{160}w_{3,s,5}\\
&\quad + \gamma_{152}w_{3,s,6} + \gamma_{112}w_{3,s,7} + \gamma_{67}w_{3,s,8} + \gamma_{70}w_{3,s,9}\\
&\quad + \gamma_{58}w_{3,s,10} + \gamma_{88}w_{3,s,11} + \gamma_{80}w_{3,s,12} + \gamma_{82}w_{3,s,13} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases} \gamma_{\{119,195\}}, & s = 3, \\ \gamma_{119}, & s \geqslant 4, \end{cases} \quad a_2 = \begin{cases} \gamma_{\{122,195\}}, & s = 3, \\ \gamma_{122}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{124,193,194,195\}}, & s = 3, \\ \gamma_{124}, & s \geqslant 4. \end{cases}$$

From these equalities, we get

$$\begin{cases} \gamma_j = 0,\ j = 53, \ldots, 88;\ 107, \ldots, 112;\ 116, \\ 117, 118,\ 120, 121, 123;\ 135, \ldots, 160, \\ a_1 = a_2 = a_3 = 0. \end{cases} \tag{6.4.9.2}$$

With the aid of (6.4.9.2) the homomorphisms $g_j, 1 \leqslant j \leqslant 4$, send (6.4.9.1) to

$$\begin{aligned} &\gamma_{177}w_{3,s,1} + \gamma_{182}w_{3,s,2} + \gamma_{170}w_{3,s,3} + \gamma_{185}w_{3,s,4} + \gamma_{161}w_{3,s,5} \\ &\quad + \gamma_{174}w_{3,s,6} + \gamma_{165}w_{3,s,7} + \gamma_{127}w_{3,s,8} + \gamma_{89}w_{3,s,9} \\ &\quad + \gamma_{115}w_{3,s,10} + \gamma_{93}w_{3,s,11} + \gamma_{98}w_{3,s,12} + \gamma_{99}w_{3,s,13} = 0, \\ &\gamma_{178}w_{3,s,1} + a_4w_{3,s,2} + \gamma_{171}w_{3,s,3} + \gamma_{186}w_{3,s,4} + \gamma_{162}w_{3,s,5} \\ &\quad + \gamma_{173}w_{3,s,6} + \gamma_{166}w_{3,s,7} + \gamma_{128}w_{3,s,8} + \gamma_{90}w_{3,s,9} \\ &\quad + \gamma_{114}w_{3,s,10} + \gamma_{94}w_{3,s,11} + \gamma_{97}w_{3,s,12} + \gamma_{103}w_{3,s,13} = 0, \\ &\gamma_{179}w_{3,s,1} + a_5w_{3,s,2} + a_6w_{3,s,3} + \gamma_{187}w_{3,s,4} + \gamma_{163}w_{3,s,5} \\ &\quad + a_7w_{3,s,6} + \gamma_{167}w_{3,s,7} + \gamma_{129}w_{3,s,8} + \gamma_{91}w_{3,s,9} \\ &\quad + \gamma_{113}w_{3,s,10} + \gamma_{95}w_{3,s,11} + \gamma_{100}w_{3,s,12} + \gamma_{102}w_{3,s,13} = 0, \\ &\gamma_{180}w_{3,s,1} + a_8w_{3,s,2} + a_9w_{3,s,3} + \gamma_{188}w_{3,s,4} + \gamma_{164}w_{3,s,5} \\ &\quad + a_{10}w_{3,s,6} + \gamma_{168}w_{3,s,7} + \gamma_{130}w_{3,s,8} + \gamma_{92}w_{3,s,9} \\ &\quad + \gamma_{125}w_{3,s,10} + \gamma_{96}w_{3,s,11} + \gamma_{101}w_{3,s,12} + \gamma_{104}w_{3,s,13} = 0, \end{aligned}$$

where

$$a_4 = \begin{cases} \gamma_{\{119,181\}}, & s = 3, \\ \gamma_{181}, & s \geqslant 4, \end{cases} \quad a_5 = \begin{cases} \gamma_{\{124,183\}}, & s = 3, \\ \gamma_{183}, & s \geqslant 4, \end{cases}$$

$$a_6 = \begin{cases} \gamma_{\{169,193\}}, & s = 3, \\ \gamma_{169}, & s \geqslant 4, \end{cases} \quad a_7 = \begin{cases} \gamma_{\{172,194\}}, & s = 3, \\ \gamma_{172}, & s \geqslant 4, \end{cases}$$

$$a_8 = \begin{cases} \gamma_{\{124,184,189,190\}}, & s = 3, \\ \gamma_{184}, & s \geqslant 4, \end{cases} \quad a_9 = \begin{cases} \gamma_{\{176,189,191,193\}}, & s = 3, \\ \gamma_{176}, & s \geqslant 4, \end{cases}$$

$$a_{10} = \begin{cases} \gamma_{\{175,190,191,194\}}, & s = 3, \\ \gamma_{175}, & s \geqslant 4. \end{cases}$$

Hence, we obtain

$$(6.4.9.3)\qquad \begin{cases} a_i = 0,\ i = 4, 5, \ldots, 10, \\ \gamma_j = 0, j = 89, \ldots, 104; 113, 114, 115, 125, 127, 128, \\ \qquad\qquad 129, 130, 161, \ldots, 168, 170, 171, 173, 174, \\ \qquad\qquad 177, 178, 179, 180, 182, 185, 186, 187, 188. \end{cases}$$

With the aid of (6.4.9.2) and (6.4.9.3), the homomorphism h sends (6.4.9.1) to

$$\begin{aligned} &a_{11}w_{3,s,1} + a_{12}w_{3,s,2} + \gamma_{189}w_{3,s,3} + a_{13}w_{3,s,4} + \gamma_{190}w_{3,s,5} \\ &\quad + \gamma_{191}w_{3,s,6} + \gamma_{192}w_{3,s,7} + \gamma_{132}w_{3,s,8} + \gamma_{131}w_{3,s,9} \\ &\quad + \gamma_{105}w_{3,s,10} + \gamma_{133}w_{3,s,11} + \gamma_{134}w_{3,s,12} + \gamma_{106}w_{3,s,13} = 0, \end{aligned}$$

where

$$a_{11} = \begin{cases} \gamma_{184}, & s = 3, \\ \gamma_{195}, & s \geqslant 4, \end{cases} \quad a_{12} = \begin{cases} \gamma_{176}, & s = 3, \\ \gamma_{193}, & s \geqslant 4, \end{cases} \quad a_{13} = \begin{cases} \gamma_{175}, & s = 3, \\ \gamma_{194}, & s \geqslant 4. \end{cases}$$

From this, it implies

$$(6.4.9.4)\qquad \begin{cases} a_{11} = a_{12} = a_{13} = 0, \\ \gamma_j = 0,\ j = 105, 106, 131, \ldots, 134, 189, \ldots, 192. \end{cases}$$

From (6.4.9.2), (6.4.9.3) and (6.4.9.4), we see that $\gamma_j = 0$ for all j. So, the proposition is proved. □

Remark 6.4.10. The element $[\theta]$ defined as in the proof of Proposition 6.4.7, is an $GL_4(\mathbb{F}_2)$-invariant in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{33}$.

6.5. **The case $t \geqslant 4$.**

According to Kameko [8], for $t \geqslant 4$, $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+t+1}+2^{s+1}-3} = 14$ with a basis given by the following classes:

$$\begin{aligned} w_{t,s,1} &= [2^s - 1, 2^s - 1, 2^{s+t+1} - 1], \\ w_{t,s,2} &= [2^s - 1, 2^{s+t+1} - 1, 2^s - 1], \\ w_{t,s,3} &= [2^{s+t+1} - 1, 2^s - 1, 2^s - 1], \\ w_{t,s,4} &= [2^s - 1, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 1], \\ w_{t,s,5} &= [2^{s+1} - 1, 2^s - 1, 2^{s+t+1} - 2^s - 1], \\ w_{t,s,6} &= [2^{s+1} - 1, 2^{s+t+1} - 2^s - 1, 2^s - 1], \\ w_{t,s,7} &= [2^{s+1} - 1, 3.2^s - 1, 2^{s+t+1} - 3.2^s - 1], \\ w_{t,s,8} &= [2^{s+1} - 1, 2^{s+t} - 1, 2^{s+t} - 1], \\ w_{t,s,9} &= [2^{s+t} - 1, 2^{s+1} - 1, 2^{s+t} - 1], \\ w_{t,s,10} &= [2^{s+t} - 1, 2^{s+t} - 1, 2^{s+1} - 1], \\ w_{t,s,11} &= [2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1, 2^{s+t} - 1], \\ w_{t,s,12} &= [2^{s+2} - 1, 2^{s+t} - 1, 2^{s+t} - 2^{s+1} - 1], \\ w_{t,s,13} &= [2^{s+t} - 1, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1], \\ w_{t,s,14} &= [2^{s+3} - 1, 2^{s+t} - 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1]. \end{aligned}$$

So we easily obtain

Proposition 6.5.1. *For any $t \geqslant 4$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+t+1}+2^{s+1}-3}$ is an $\mathbb{F}_2$-vector space of dimension 56 with a basis consisting of all the classes represented by the following monomials:*

$$\begin{aligned}
a_{t,s,1} &= (0, 2^{s+1}-1, 2^{s+t}-1, 2^{s+t}-1),\\
a_{t,s,2} &= (0, 2^{s+t}-1, 2^{s+1}-1, 2^{s+t}-1),\\
a_{t,s,3} &= (0, 2^{s+t}-1, 2^{s+t}-1, 2^{s+1}-1),\\
a_{t,s,4} &= (2^{s+1}-1, 0, 2^{s+t}-1, 2^{s+t}-1),\\
a_{t,s,5} &= (2^{s+1}-1, 2^{s+t}-1, 0, 2^{s+t}-1),\\
a_{t,s,6} &= (2^{s+1}-1, 2^{s+t}-1, 2^{s+t}-1, 0),\\
a_{t,s,7} &= (2^{s+t}-1, 0, 2^{s+1}-1, 2^{s+t}-1),\\
a_{t,s,8} &= (2^{s+t}-1, 0, 2^{s+t}-1, 2^{s+1}-1),\\
a_{t,s,9} &= (2^{s+t}-1, 2^{s+1}-1, 0, 2^{s+t}-1),\\
a_{t,s,10} &= (2^{s+t}-1, 2^{s+1}-1, 2^{s+t}-1, 0),\\
a_{t,s,11} &= (2^{s+t}-1, 2^{s+t}-1, 0, 2^{s+1}-1),\\
a_{t,s,12} &= (2^{s+t}-1, 2^{s+t}-1, 2^{s+1}-1, 0),\\
a_{t,s,13} &= (0, 2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 2^{s+t}-1),\\
a_{t,s,14} &= (0, 2^{s+2}-1, 2^{s+t}-1, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,15} &= (0, 2^{s+t}-1, 2^{s+2}-1, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,16} &= (2^{s+2}-1, 0, 2^{s+t}-2^{s+1}-1, 2^{s+t}-1),\\
a_{t,s,17} &= (2^{s+2}-1, 0, 2^{s+t}-1, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,18} &= (2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 0, 2^{s+t}-1),\\
a_{t,s,19} &= (2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 2^{s+t}-1, 0),\\
a_{t,s,20} &= (2^{s+2}-1, 2^{s+t}-1, 0, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,21} &= (2^{s+2}-1, 2^{s+t}-1, 2^{s+t}-2^{s+1}-1, 0),\\
a_{t,s,22} &= (2^{s+t}-1, 0, 2^{s+2}-1, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,23} &= (2^{s+t}-1, 2^{s+2}-1, 0, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,24} &= (2^{s+t}-1, 2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 0),\\
a_{t,s,25} &= (0, 2^{s+3}-1, 2^{s+t}-2^{s+2}-1, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,26} &= (2^{s+3}-1, 0, 2^{s+t}-2^{s+2}-1, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,27} &= (2^{s+3}-1, 2^{s+t}-2^{s+2}-1, 0, 2^{s+t}-2^{s+1}-1),\\
a_{t,s,28} &= (2^{s+3}-1, 2^{s+t}-2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 0),\\
a_{t,s,29} &= (0, 2^{s}-1, 2^{s}-1, 2^{s+t+1}-1),\\
a_{t,s,30} &= (0, 2^{s}-1, 2^{s+t+1}-1, 2^{s}-1),\\
a_{t,s,31} &= (0, 2^{s+t+1}-1, 2^{s}-1, 2^{s}-1),\\
a_{t,s,32} &= (2^{s}-1, 0, 2^{s}-1, 2^{s+t+1}-1),\\
a_{t,s,33} &= (2^{s}-1, 0, 2^{s+t+1}-1, 2^{s}-1),\\
a_{t,s,34} &= (2^{s}-1, 2^{s}-1, 0, 2^{s+t+1}-1),\\
a_{t,s,35} &= (2^{s}-1, 2^{s}-1, 2^{s+t+1}-1, 0),\\
a_{t,s,36} &= (2^{s}-1, 2^{s+t+1}-1, 0, 2^{s}-1),\\
a_{t,s,37} &= (2^{s}-1, 2^{s+t+1}-1, 2^{s}-1, 0),\\
a_{t,s,38} &= (2^{s+t+1}-1, 0, 2^{s}-1, 2^{s}-1),\\
a_{t,s,39} &= (2^{s+t+1}-1, 2^{s}-1, 0, 2^{s}-1),\\
a_{t,s,40} &= (2^{s+t+1}-1, 2^{s}-1, 2^{s}-1, 0),
\end{aligned}$$

$$\begin{aligned}
a_{t,s,41} &= (0, 2^s - 1, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,42} &= (0, 2^{s+1} - 1, 2^s - 1, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,43} &= (0, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 1, 2^s - 1),\\
a_{t,s,44} &= (2^s - 1, 0, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,45} &= (2^s - 1, 2^{s+1} - 1, 0, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,46} &= (2^s - 1, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 1, 0),\\
a_{t,s,47} &= (2^{s+1} - 1, 0, 2^s - 1, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,48} &= (2^{s+1} - 1, 0, 2^{s+t+1} - 2^s - 1, 2^s - 1),\\
a_{t,s,49} &= (2^{s+1} - 1, 2^s - 1, 0, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,50} &= (2^{s+1} - 1, 2^s - 1, 2^{s+t+1} - 2^s - 1, 0),\\
a_{t,s,51} &= (2^{s+1} - 1, 2^{s+t+1} - 2^s - 1, 0, 2^s - 1),\\
a_{t,s,52} &= (2^{s+1} - 1, 2^{s+t+1} - 2^s - 1, 2^s - 1, 0),\\
a_{t,s,53} &= (0, 2^{s+1} - 1, 3.2^s - 1, 2^{s+t+1} - 3.2^s - 1),\\
a_{t,s,54} &= (2^{s+1} - 1, 0, 3.2^s - 1, 2^{s+t+1} - 3.2^s - 1),\\
a_{t,s,55} &= (2^{s+1} - 1, 3.2^s - 1, 0, 2^{s+t+1} - 3.2^s - 1),\\
a_{t,s,56} &= (2^{s+1} - 1, 3.2^s - 1, 2^{s+t+1} - 3.2^s - 1, 0).
\end{aligned}$$

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+1}-3}$.

Set $\mu_5(1) = 150, \mu_5(2) = 195$ and $\mu_5(s) = 210$ for $s \geqslant 3$. The main result of this subsection is

Theorem 6.5.2. *For any $t \geqslant 4$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+1}-3}$ is an $\mathbb{F}_2$-vector space of dimension $\mu_5(s) - 56$ with a basis consisting of all the classes represented by the following monomials:*

For $s \geqslant 1$,

$$\begin{aligned}
a_{t,s,57} &= (1, 2^{s+1} - 2, 2^{s+t} - 1, 2^{s+t} - 1),\\
a_{t,s,58} &= (1, 2^{s+t} - 1, 2^{s+1} - 2, 2^{s+t} - 1),\\
a_{t,s,59} &= (1, 2^{s+t} - 1, 2^{s+t} - 1, 2^{s+1} - 2),\\
a_{t,s,60} &= (2^{s+t} - 1, 1, 2^{s+1} - 2, 2^{s+t} - 1),\\
a_{t,s,61} &= (2^{s+t} - 1, 1, 2^{s+t} - 1, 2^{s+1} - 2),\\
a_{t,s,62} &= (2^{s+t} - 1, 2^{s+t} - 1, 1, 2^{s+1} - 2),\\
a_{t,s,63} &= (1, 2^{s+1} - 1, 2^{s+t} - 2, 2^{s+t} - 1),\\
a_{t,s,64} &= (1, 2^{s+1} - 1, 2^{s+t} - 1, 2^{s+t} - 2),\\
a_{t,s,65} &= (1, 2^{s+t} - 2, 2^{s+1} - 1, 2^{s+t} - 1),\\
a_{t,s,66} &= (1, 2^{s+t} - 2, 2^{s+t} - 1, 2^{s+1} - 1),\\
a_{t,s,67} &= (1, 2^{s+t} - 1, 2^{s+1} - 1, 2^{s+t} - 2),\\
a_{t,s,68} &= (1, 2^{s+t} - 1, 2^{s+t} - 2, 2^{s+1} - 1),\\
a_{t,s,69} &= (2^{s+1} - 1, 1, 2^{s+t} - 2, 2^{s+t} - 1),\\
a_{t,s,70} &= (2^{s+1} - 1, 1, 2^{s+t} - 1, 2^{s+t} - 2),\\
a_{t,s,71} &= (2^{s+1} - 1, 2^{s+t} - 1, 1, 2^{s+t} - 2),\\
a_{t,s,72} &= (2^{s+t} - 1, 1, 2^{s+1} - 1, 2^{s+t} - 2),\\
a_{t,s,73} &= (2^{s+t} - 1, 1, 2^{s+t} - 2, 2^{s+1} - 1),\\
a_{t,s,74} &= (2^{s+t} - 1, 2^{s+1} - 1, 1, 2^{s+t} - 2),\\
a_{t,s,75} &= (1, 2^{s+2} - 2, 2^{s+t} - 2^{s+1} - 1, 2^{s+t} - 1),\\
a_{t,s,76} &= (1, 2^{s+2} - 2, 2^{s+t} - 1, 2^{s+t} - 2^{s+1} - 1),
\end{aligned}$$

$$a_{t,s,77} = (1, 2^{s+t} - 1, 2^{s+2} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,78} = (2^{s+t} - 1, 1, 2^{s+2} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,79} = (1, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 2, 2^{s+t} - 1),$$
$$a_{t,s,80} = (1, 2^{s+2} - 1, 2^{s+t} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,81} = (1, 2^{s+t} - 1, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,82} = (2^{s+2} - 1, 1, 2^{s+t} - 2^{s+1} - 2, 2^{s+t} - 1),$$
$$a_{t,s,83} = (2^{s+2} - 1, 1, 2^{s+t} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,84} = (2^{s+2} - 1, 2^{s+t} - 1, 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,85} = (2^{s+t} - 1, 1, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,86} = (2^{s+t} - 1, 2^{s+2} - 1, 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,87} = (1, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1, 2^{s+t} - 2),$$
$$a_{t,s,88} = (1, 2^{s+2} - 1, 2^{s+t} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,89} = (1, 2^{s+t} - 2, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,90} = (2^{s+2} - 1, 1, 2^{s+t} - 2^{s+1} - 1, 2^{s+t} - 2),$$
$$a_{t,s,91} = (2^{s+2} - 1, 1, 2^{s+t} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,92} = (2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1, 1, 2^{s+t} - 2),$$
$$a_{t,s,93} = (1, 2^{s+3} - 2, 2^{s+t} - 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,94} = (1, 2^{s+3} - 1, 2^{s+t} - 2^{s+2} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,95} = (2^{s+3} - 1, 1, 2^{s+t} - 2^{s+2} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,96} = (1, 2^{s+3} - 1, 2^{s+t} - 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,97} = (2^{s+3} - 1, 1, 2^{s+t} - 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,98} = (2^{s+3} - 1, 2^{s+t} - 2^{s+2} - 1, 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,99} = (3, 2^{s+t} - 3, 2^{s+1} - 2, 2^{s+t} - 1),$$
$$a_{t,s,100} = (3, 2^{s+t} - 3, 2^{s+t} - 1, 2^{s+1} - 2),$$
$$a_{t,s,101} = (3, 2^{s+t} - 1, 2^{s+t} - 3, 2^{s+1} - 2),$$
$$a_{t,s,102} = (2^{s+t} - 1, 3, 2^{s+t} - 3, 2^{s+1} - 2),$$
$$a_{t,s,103} = (3, 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 2, 2^{s+t} - 1),$$
$$a_{t,s,104} = (3, 2^{s+2} - 3, 2^{s+t} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,105} = (3, 2^{s+t} - 1, 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,106} = (2^{s+t} - 1, 3, 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,107} = (3, 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 1, 2^{s+t} - 2),$$
$$a_{t,s,108} = (3, 2^{s+2} - 3, 2^{s+t} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,109} = (3, 2^{s+t} - 3, 2^{s+2} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,110} = (3, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 3, 2^{s+t} - 2),$$
$$a_{t,s,111} = (2^{s+2} - 1, 3, 2^{s+t} - 2^{s+1} - 3, 2^{s+t} - 2),$$
$$a_{t,s,112} = (3, 2^{s+2} - 1, 2^{s+t} - 3, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,113} = (3, 2^{s+t} - 3, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,114} = (2^{s+2} - 1, 3, 2^{s+t} - 3, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,115} = (3, 2^{s+3} - 3, 2^{s+t} - 2^{s+2} - 2, 2^{s+t} - 2^{s+1} - 1),$$
$$a_{t,s,116} = (3, 2^{s+3} - 3, 2^{s+t} - 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,117} = (3, 2^{s+3} - 1, 2^{s+t} - 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,118} = (2^{s+3} - 1, 3, 2^{s+t} - 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,119} = (7, 2^{s+t} - 5, 2^{s+t} - 3, 2^{s+1} - 2),$$
$$a_{t,s,120} = (7, 2^{s+t} - 5, 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 2),$$
$$a_{t,s,121} = (7, 2^{s+3} - 5, 2^{s+t} - 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 2),$$

$$\begin{aligned}
a_{t,s,122} &= (1, 2^s - 1, 2^s - 1, 2^{s+t+1} - 2),\\
a_{t,s,123} &= (1, 2^s - 1, 2^{s+t+1} - 2, 2^s - 1),\\
a_{t,s,124} &= (1, 2^{s+t+1} - 2, 2^s - 1, 2^s - 1),\\
a_{t,s,125} &= (1, 2^s - 1, 2^{s+1} - 2, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,126} &= (1, 2^{s+1} - 2, 2^s - 1, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,127} &= (1, 2^{s+1} - 2, 2^{s+t+1} - 2^s - 1, 2^s - 1),\\
a_{t,s,128} &= (1, 2^s - 1, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 2),\\
a_{t,s,129} &= (1, 2^{s+1} - 1, 2^s - 1, 2^{s+t+1} - 2^s - 2),\\
a_{t,s,130} &= (1, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 2, 2^s - 1),\\
a_{t,s,131} &= (2^{s+1} - 1, 1, 2^s - 1, 2^{s+t+1} - 2^s - 2),\\
a_{t,s,132} &= (2^{s+1} - 1, 1, 2^{s+t+1} - 2^s - 2, 2^s - 1),\\
a_{t,s,133} &= (1, 2^{s+1} - 2, 3.2^s - 1, 2^{s+t+1} - 3.2^s - 1),\\
a_{t,s,134} &= (1, 2^{s+1} - 1, 3.2^s - 2, 2^{s+t+1} - 3.2^s - 1),\\
a_{t,s,135} &= (2^{s+1} - 1, 1, 3.2^s - 2, 2^{s+t+1} - 3.2^s - 1),\\
a_{t,s,136} &= (1, 2^{s+1} - 1, 3.2^s - 1, 2^{s+t+1} - 3.2^s - 2),\\
a_{t,s,137} &= (2^{s+1} - 1, 1, 3.2^s - 1, 2^{s+t+1} - 3.2^s - 2),\\
a_{t,s,138} &= (2^{s+1} - 1, 3.2^s - 1, 1, 2^{s+t+1} - 3.2^s - 2),\\
a_{t,s,139} &= (3, 2^{s+1} - 1, 2^{s+t} - 3, 2^{s+t} - 2),\\
a_{t,s,140} &= (3, 2^{s+t} - 3, 2^{s+1} - 1, 2^{s+t} - 2),\\
a_{t,s,141} &= (3, 2^{s+t} - 3, 2^{s+t} - 2, 2^{s+1} - 1).
\end{aligned}$$

For $s = 1$,

$$\begin{aligned}
a_{t,1,142} &= (3, 3, 2^{t+1} - 4, 2^{t+1} - 1), \quad a_{t,1,143} = (3, 3, 2^{t+1} - 1, 2^{t+1} - 4),\\
a_{t,1,144} &= (3, 2^{t+1} - 1, 3, 2^{t+1} - 4), \quad a_{t,1,145} = (2^{t+1} - 1, 3, 3, 2^{t+1} - 4),\\
a_{t,1,146} &= (3, 7, 2^{t+1} - 5, 2^{t+1} - 4), \quad a_{t,1,147} = (7, 2^{t+1} - 5, 3, 2^{t+1} - 4),\\
a_{t,1,148} &= (7, 3, 2^{t+1} - 5, 2^{t+1} - 4), \quad a_{t,1,149} = (7, 7, 2^{t+1} - 8, 2^{t+1} - 5),\\
a_{t,1,150} &= (7, 7, 2^{t+1} - 7, 2^{t+1} - 6).
\end{aligned}$$

For $s \geqslant 2$,

$$\begin{aligned}
a_{t,s,142} &= (2^{s+1} - 1, 3, 2^{s+t} - 3, 2^{s+t} - 2),\\
a_{t,s,143} &= (3, 2^{s+1} - 3, 2^{s+t} - 2, 2^{s+t} - 1),\\
a_{t,s,144} &= (3, 2^{s+1} - 3, 2^{s+t} - 1, 2^{s+t} - 2),\\
a_{t,s,145} &= (3, 2^{s+t} - 1, 2^{s+1} - 3, 2^{s+t} - 2),\\
a_{t,s,146} &= (2^{s+t} - 1, 3, 2^{s+1} - 3, 2^{s+t} - 2),\\
a_{t,s,147} &= (7, 2^{s+t} - 5, 2^{s+1} - 3, 2^{s+t} - 2),\\
a_{t,s,148} &= (7, 2^{s+2} - 5, 2^{s+t} - 2^{s+1} - 3, 2^{s+t} - 2),\\
a_{t,s,149} &= (7, 2^{s+2} - 5, 2^{s+t} - 3, 2^{s+t} - 2^{s+1} - 2),\\
a_{t,s,150} &= (1, 2^s - 2, 2^s - 1, 2^{s+t+1} - 1),\\
a_{t,s,151} &= (1, 2^s - 2, 2^{s+t+1} - 1, 2^s - 1),\\
a_{t,s,152} &= (1, 2^s - 1, 2^s - 2, 2^{s+t+1} - 1),\\
a_{t,s,153} &= (1, 2^s - 1, 2^{s+t+1} - 1, 2^s - 2),\\
a_{t,s,154} &= (1, 2^{s+t+1} - 1, 2^s - 2, 2^s - 1),\\
a_{t,s,155} &= (1, 2^{s+t+1} - 1, 2^s - 1, 2^s - 2),\\
a_{t,s,156} &= (2^s - 1, 1, 2^s - 2, 2^{s+t+1} - 1),\\
a_{t,s,157} &= (2^s - 1, 1, 2^{s+t+1} - 1, 2^s - 2),
\end{aligned}$$

$$
\begin{aligned}
a_{t,s,158} &= (2^s-1, 2^{s+t+1}-1, 1, 2^s-2),\\
a_{t,s,159} &= (2^{s+t+1}-1, 1, 2^s-2, 2^s-1),\\
a_{t,s,160} &= (2^{s+t+1}-1, 1, 2^s-1, 2^s-2),\\
a_{t,s,161} &= (2^{s+t+1}-1, 2^s-1, 1, 2^s-2),\\
a_{t,s,162} &= (2^s-1, 1, 2^s-1, 2^{s+t+1}-2),\\
a_{t,s,163} &= (2^s-1, 1, 2^{s+t+1}-2, 2^s-1),\\
a_{t,s,164} &= (2^s-1, 2^s-1, 1, 2^{s+t+1}-2),\\
a_{t,s,165} &= (1, 2^s-2, 2^{s+1}-1, 2^{s+t+1}-2^s-1),\\
a_{t,s,166} &= (1, 2^{s+1}-1, 2^s-2, 2^{s+t+1}-2^s-1),\\
a_{t,s,167} &= (1, 2^{s+1}-1, 2^{s+t+1}-2^s-1, 2^s-2),\\
a_{t,s,168} &= (2^{s+1}-1, 1, 2^s-2, 2^{s+t+1}-2^s-1),\\
a_{t,s,169} &= (2^{s+1}-1, 1, 2^{s+t+1}-2^s-1, 2^s-2),\\
a_{t,s,170} &= (2^{s+1}-1, 2^{s+t+1}-2^s-1, 1, 2^s-2),\\
a_{t,s,171} &= (2^s-1, 1, 2^{s+1}-2, 2^{s+t+1}-2^s-1),\\
a_{t,s,172} &= (2^s-1, 1, 2^{s+1}-1, 2^{s+t+1}-2^s-2),\\
a_{t,s,173} &= (2^s-1, 2^{s+1}-1, 1, 2^{s+t+1}-2^s-2),\\
a_{t,s,174} &= (2^{s+1}-1, 2^s-1, 1, 2^{s+t+1}-2^s-2),\\
a_{t,s,175} &= (3, 2^{s+1}-3, 2^s-2, 2^{s+t+1}-2^s-1),\\
a_{t,s,176} &= (3, 2^{s+1}-3, 2^{s+t+1}-2^s-1, 2^s-2),\\
a_{t,s,177} &= (3, 2^{s+1}-1, 2^{s+t+1}-2^s-3, 2^s-2),\\
a_{t,s,178} &= (2^{s+1}-1, 3, 2^{s+t+1}-2^s-3, 2^s-2),\\
a_{t,s,179} &= (3, 2^{s+1}-3, 3.2^s-2, 2^{s+t+1}-3.2^s-1),\\
a_{t,s,180} &= (3, 2^{s+1}-3, 3.2^s-1, 2^{s+t+1}-3.2^s-2),\\
a_{t,s,181} &= (3, 2^{s+1}-1, 3.2^s-3, 2^{s+t+1}-3.2^s-2),\\
a_{t,s,182} &= (2^{s+1}-1, 3, 3.2^s-3, 2^{s+t+1}-3.2^s-2),\\
a_{t,s,183} &= (3, 2^s-1, 2^{s+t+1}-3, 2^s-2),\\
a_{t,s,184} &= (3, 2^{s+t+1}-3, 2^s-2, 2^s-1),\\
a_{t,s,185} &= (3, 2^{s+t+1}-3, 2^s-1, 2^s-2),\\
a_{t,s,186} &= (3, 2^s-1, 2^{s+1}-3, 2^{s+t+1}-2^s-2),\\
a_{t,s,187} &= (3, 2^{s+1}-3, 2^s-1, 2^{s+t+1}-2^s-2),\\
a_{t,s,188} &= (3, 2^{s+1}-3, 2^{s+t+1}-2^s-2, 2^s-1).
\end{aligned}
$$

For $s = 2$,

$$
\begin{aligned}
a_{t,2,189} &= (7, 7, 2^{t+2}-7, 2^{t+2}-2), & a_{t,2,190} &= (3, 3, 3, 2^{t+3}-4),\\
a_{t,2,191} &= (3, 3, 2^{t+3}-4, 3), & a_{t,2,192} &= (3, 3, 4, 2^{t+3}-5),\\
a_{t,2,193} &= (3, 3, 7, 2^{t+3}-8), & a_{t,2,194} &= (3, 7, 3, 2^{t+3}-8),\\
a_{t,2,195} &= (7, 3, 3, 2^{t+3}-8).
\end{aligned}
$$

For $s \geqslant 3$,

$$
\begin{aligned}
a_{t,s,189} &= (7, 2^{s+1}-5, 2^{s+t}-3, 2^{s+t}-2),\\
a_{t,s,190} &= (3, 2^s-3, 2^s-2, 2^{s+t+1}-1),\\
a_{t,s,191} &= (3, 2^s-3, 2^{s+t+1}-1, 2^s-2),\\
a_{t,s,192} &= (3, 2^{s+t+1}-1, 2^s-3, 2^s-2),\\
a_{t,s,193} &= (2^{s+t+1}-1, 3, 2^s-3, 2^s-2),\\
a_{t,s,194} &= (3, 2^s-3, 2^s-1, 2^{s+t+1}-2),
\end{aligned}
$$

$$\begin{aligned}
a_{t,s,195} &= (3, 2^s - 3, 2^{s+t+1} - 2, 2^s - 1),\\
a_{t,s,196} &= (3, 2^s - 1, 2^s - 3, 2^{s+t+1} - 2),\\
a_{t,s,197} &= (2^s - 1, 3, 2^s - 3, 2^{s+t+1} - 2),\\
a_{t,s,198} &= (2^s - 1, 3, 2^{s+t+1} - 3, 2^s - 2),\\
a_{t,s,199} &= (3, 2^s - 3, 2^{s+1} - 2, 2^{s+t+1} - 2^s - 1),\\
a_{t,s,200} &= (3, 2^s - 3, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 2),\\
a_{t,s,201} &= (3, 2^{s+1} - 1, 2^s - 3, 2^{s+t+1} - 2^s - 2),\\
a_{t,s,202} &= (2^{s+1} - 1, 3, 2^s - 3, 2^{s+t+1} - 2^s - 2),\\
a_{t,s,203} &= (2^s - 1, 3, 2^{s+1} - 3, 2^{s+t+1} - 2^s - 2),\\
a_{t,s,204} &= (7, 2^{s+t+1} - 5, 2^s - 3, 2^s - 2),\\
a_{t,s,205} &= (7, 2^{s+1} - 5, 2^s - 3, 2^{s+t+1} - 2^s - 2),\\
a_{t,s,206} &= (7, 2^{s+1} - 5, 2^{s+t+1} - 2^s - 3, 2^s - 2),\\
a_{t,s,207} &= (7, 2^{s+1} - 5, 3.2^s - 3, 2^{s+t+1} - 3.2^s - 2).
\end{aligned}$$

For $s = 3$,

$$\begin{aligned}
&a_{t,3,208} = (7, 7, 2^{t+4} - 7, 6), \quad a_{t,3,209} = (7, 7, 7, 2^{t+4} - 8),\\
&a_{t,3,210} = (7, 7, 9, 2^{t+4} - 10).
\end{aligned}$$

For $s \geqslant 4$,

$$\begin{aligned}
a_{t,s,208} &= (7, 2^s - 5, 2^s - 3, 2^{s+t+1} - 2),\\
a_{t,s,209} &= (7, 2^s - 5, 2^{s+t+1} - 3, 2^s - 2),\\
a_{t,s,210} &= (7, 2^s - 5, 2^{s+1} - 3, 2^{s+t+1} - 2^s - 2).
\end{aligned}$$

The theorem is proved by combining the following propositions.

Proposition 6.5.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+1}-3}$ is generated by the $\mu_5(s) - 56$ elements listed in Theorem 6.5.2.*

The proof of this proposition is based on Theorem 2.5 and the following lemma.

Lemma 6.5.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 1\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 1\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 1\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$\begin{aligned}
&(7, 7, 27, 24), (7, 15, 16, 27), (15, 7, 16, 27), (7, 15, 17, 26),\\
&(15, 7, 17, 26), (15, 15, 16, 19), (15, 15, 19, 16), (15, 15, 17, 18).
\end{aligned}$$

We proved the lemma for the matrices associated with

$$(7,7,27,24),(15,7,16,27),(15,7,17,26),$$
$$(15,15,16,19),(15,15,19,16),(15,15,17,18).$$

The others can be obtained by a similar computation. By a direct computation, we have

$$\begin{aligned}
(7,7,27,24) &= Sq^1\big((7,7,27,23)+(7,5,29,23)+(7,7,21,29)\big)\\
&\quad + Sq^2\big((7,6,27,23)+(7,3,30,23)+(7,7,26,23)+(7,7,22,27)\\
&\quad + (7,7,19,30)+(7,3,23,30)+(7,3,15,38)\big)+Sq^4\big((4,7,27,23)\\
&\quad + (5,6,27,23)+(5,3,30,23)+(5,7,26,23)+(11,5,22,23)\\
&\quad + (5,7,22,27)+(5,7,19,30)+(11,5,15,30)+(5,3,23,30)\big)\\
&\quad + Sq^8\big((7,5,22,23)+(7,5,15,30)\big)+(4,11,27,23)+(4,7,27,27)\\
&\quad + (7,6,27,25)+(5,10,27,23)+(5,6,27,27)+(7,3,30,25)\\
&\quad + (5,3,30,27)+(7,7,26,25)+(5,11,26,23)+(7,5,26,27)\\
&\quad + (7,7,24,27)+(5,11,22,27)+(5,11,19,30)\\
&\quad + (7,3,25,30)+(5,3,27,30) \mod \mathcal{L}_4(3;3;2;2;2),\\
(15,7,16,27) &= Sq^1(15,7,13,29)+Sq^2\big((15,7,11,30)\\
&\quad + (15,7,14,27)\big)+Sq^4\big((15,5,14,27)+(15,5,11,30)\big)\\
&\quad + Sq^8\big((9,7,14,27)+(11,5,14,27)+(9,7,11,30)\\
&\quad + (11,5,11,30)\big)+(15,5,18,27)+(9,7,22,27)+(11,5,22,27)\\
&\quad + (9,7,19,30)+(11,5,19,30) \mod \mathcal{L}_4(3;3;2;2;2),\\
(15,7,17,26) &= Sq^1(15,7,15,27)+Sq^2(15,7,15,26)\\
&\quad + Sq^4\big((15,5,15,26)+(15,4,15,27)\big)+Sq^8\big((9,7,15,26)\\
&\quad + (11,5,15,26)+(8,7,15,27)+(11,4,15,27)\big)+(15,5,19,26)\\
&\quad + (11,5,23,26)+(15,7,16,27)+(8,7,23,27)+(15,4,19,27)\\
&\quad + (11,4,23,27)+(9,7,23,26) \mod \mathcal{L}_4(3;3;2;2;2),\\
(15,15,16,19) &= Sq^1\big((15,15,13,21)+(15,15,11,23)+(15,15,9,25)\\
&\quad + (15,15,7,27)+(15,15,5,29)\big)+Sq^2\big((15,15,14,19)+(15,15,11,22)\\
&\quad + (15,15,10,23)+(15,15,7,26)+(15,15,6,27)+(15,15,3,30)\big)\\
&\quad + Sq^4\big((15,13,14,19)+(15,13,11,22)+(15,12,11,23)\\
&\quad + (15,13,10,23)+(15,13,7,26)+(15,12,7,27)+(15,13,6,27)\\
&\quad + (15,13,3,30)\big)+Sq^8\big((9,15,14,19)+(11,13,14,19)+(9,15,11,22)\\
&\quad + (11,13,11,22)+(8,15,11,23)+(11,12,11,23)+(9,15,10,23)\\
&\quad + (11,13,10,23)+(9,15,7,26)+(11,13,7,26)+(8,15,7,27)
\end{aligned}$$

$$
\begin{aligned}
&+ (11,12,7,27) + (9,15,6,27) + (11,13,6,27) + (9,15,3,30)\\
&+ (11,13,3,30)\big) + (9,23,14,19) + (9,15,22,19) + (15,13,18,19)\\
&+ (11,21,14,19) + (11,13,22,19) + (9,23,11,22) + (9,15,19,22)\\
&+ (11,21,11,22) + (11,13,19,22) + (8,23,11,23) + (8,15,19,23)\\
&+ (11,20,11,23) + (11,12,19,23) + (9,23,10,23) + (9,15,18,23)\\
&+ (11,21,10,23) + (11,13,18,23) + (9,23,7,26) + (11,21,7,26)\\
&+ (8,23,7,27) + (11,20,7,27) + (9,23,6,27) + (11,21,6,27)\\
&+ (9,23,3,30) + (11,21,3,30) \mod \mathcal{L}_4(3;3;2;2;2),\\
(15,15,19,16) &= Sq^1(15,15,19,15) + Sq^2(15,15,18,15)\\
&+ Sq^4\big((15,12,19,15) + (15,13,18,15)\big) + Sq^8\big((8,15,19,15)\\
&+ (11,12,19,15) + (9,15,18,15) + (11,13,18,15)\big) + (15,15,18,17)\\
&+ (15,12,19,19) + (15,13,18,19) + (8,23,19,15) + (8,15,19,23)\\
&+ (11,20,19,15) + (11,12,19,23) + (9,23,18,15) + (9,15,18,23)\\
&+ (11,21,18,15) + (11,13,18,23) \mod \mathcal{L}_4(3;3;2;2;2),\\
(15,15,17,18) &= Sq^1(15,15,15,19) + Sq^2(15,15,15,18)\\
&+ Sq^4\big((15,13,15,18) + (15,12,15,19)\big) + Sq^8\big((9,15,15,18)\\
&+ (11,13,15,18) + (8,15,15,19) + (11,12,15,19)\big) + (15,15,16,19)\\
&+ (9,23,15,18) + (9,15,23,18) + (15,13,19,18) + (11,21,15,18)\\
&+ (11,13,23,18) + (8,23,15,19) + (8,15,23,19) + (15,12,19,19)\\
&+ (11,20,15,19) + (11,12,23,19) \mod \mathcal{L}_4(3;3;2;2;2).
\end{aligned}
$$

The lemma is proved. □

Combining Theorem 2.5 the lemmas in Sections 3, 4, 5 and Lemmas 6.1.1, 6.2.4, 6.2.5, 6.3.4, 6.3.5, 6.4.4, 6.4.5, 6.4.6, 6.5.4, we obtain Proposition 6.5.3.

Now, we show that the elements $[a_{t,s,j}], 57 \leqslant j \leqslant \mu_5(s)$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+1}-3}$.

Proposition 6.5.5. *For $t \geqslant 4$, the elements $[a_{t,1,j}]$, $57 \leqslant j \leqslant 150$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+2}+1}$.*

Proof. Suppose there is a linear relation

$$\sum_{j=57}^{150} \gamma_j [a_{t,1,j}] = 0, \tag{6.5.5.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Consider the homomorphisms $f_i,\ i = 1, 2, 3$. Under these homomorphisms, the images of the relation (6.5.5.1) respectively are

$$\begin{aligned}
&\gamma_{124}w_{t,1,3} + \gamma_{126}w_{t,1,5} + \gamma_{127}w_{t,1,6} + \gamma_{133}w_{t,1,7} + \gamma_{57}w_{t,1,8} + \gamma_{65}w_{t,1,9}\\
&\quad + \gamma_{66}w_{t,1,10} + \gamma_{75}w_{t,1,11} + \gamma_{76}w_{t,1,12} + \gamma_{89}w_{t,1,13} + \gamma_{93}w_{t,1,14} = 0,\\
&\gamma_{\{123,132\}}w_{t,1,3} + a_1w_{t,1,5} + \gamma_{130}w_{t,1,6} + a_2w_{t,1,7} + \gamma_{58}w_{t,1,8} + \gamma_{\{63,142\}}w_{t,1,9}\\
&\quad + \gamma_{68}w_{t,1,10} + \gamma_{79}w_{t,1,11} + \gamma_{77}w_{t,1,12} + \gamma_{\{88,149\}}w_{t,1,13} + \gamma_{94}w_{t,1,14} = 0,\\
&\gamma_{\{122,131\}}w_{t,1,3} + a_3w_{t,1,5} + a_4w_{t,1,6} + a_5w_{t,1,7} + \gamma_{59}w_{t,1,8} + \gamma_{\{64,143\}}w_{t,1,9}\\
&\quad + \gamma_{\{67,144\}}w_{t,1,10} + \gamma_{80}w_{t,1,11} + \gamma_{81}w_{t,1,12} + \gamma_{\{87,146\}}w_{t,1,13} + \gamma_{96}w_{t,1,14} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{60,69,73,78,82,91,95,125,135\}},\ a_2 = \gamma_{\{99,103,108,109,115,134,141\}},\\
&a_3 = \gamma_{\{61,70,72,83,85,90,97,128,137\}},\ a_4 = \gamma_{\{62,71,74,84,86,92,98,129,138\}},\\
&a_5 = \gamma_{\{100,101,102,104,105,106,107,110,\ldots,114,116,117,\ldots,121,136,139,140,145,147,148,150\}}.
\end{aligned}$$

From the above equalities, we get

$$\text{(6.5.5.2)}\qquad \begin{cases}
a_1 = a_2 = a_3 = a_4 = a_5 = 0,\\
\gamma_j = 0,\ j = 57, 58, 59, 65, 66, 68, 75, 76, 77,\\
79, 80, 81, 89, 93, 94, 96, 124, 126, 127, 130, 133,\\
\gamma_{\{123,132\}} = \gamma_{\{63,142\}} = \gamma_{\{88,149\}} = 0,\\
\gamma_{\{122,131\}} = \gamma_{\{64,143\}} = \gamma_{\{67,144\}} = \gamma_{\{87,146\}} = 0.
\end{cases}$$

With the aid of (6.5.5.2), the homomorphisms f_4, f_5, f_6 send (6.5.5.1) respectively to

$$\begin{aligned}
&\gamma_{123}w_{t,1,2} + \gamma_{\{63,88,125,134\}}w_{t,1,4} + \gamma_{\{108,109,115,135,141\}}w_{t,1,7}\\
&\quad + \gamma_{132}w_{t,1,6} + \gamma_{60}w_{t,1,9} + \gamma_{73}w_{t,1,10} + \gamma_{\{69,99,103,142\}}w_{t,1,8}\\
&\quad + \gamma_{82}w_{t,1,11} + \gamma_{\{91,149\}}w_{t,1,12} + \gamma_{78}w_{t,1,13} + \gamma_{95}w_{t,1,14} = 0,\\
&\gamma_{\{122,129\}}w_{t,1,2} + \gamma_{\{64,67,87,128,136\}}w_{t,1,4} + a_6w_{t,1,6} + a_7w_{t,1,7}\\
&\quad + \gamma_{\{70,100,104,143\}}w_{t,1,8} + \gamma_{61}w_{t,1,9} + \gamma_{\{72,145\}}w_{t,1,10} + \gamma_{83}w_{t,1,11}\\
&\quad + \gamma_{\{90,148\}}w_{t,1,12} + \gamma_{85}w_{t,1,13} + \gamma_{97}w_{t,1,14} = 0,\\
&\gamma_{\{122,123,125,128\}}w_{t,1,1} + a_8w_{t,1,4} + a_9w_{t,1,5} + a_{10}w_{t,1,7}\\
&\quad + \gamma_{\{71,101,105,144\}}w_{t,1,8} + \gamma_{\{74,102,106,145\}}w_{t,1,9} + \gamma_{62}w_{t,1,10}\\
&\quad + \gamma_{\{92,119,120,147\}}w_{t,1,11} + \gamma_{84}w_{t,1,12} + \gamma_{86}w_{t,1,13} + \gamma_{98}w_{t,1,14} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_6 = \gamma_{\{62,71,74,84,86,92,98,131,138\}},\quad a_8 = \gamma_{\{63,64,67,87,88,129,134,136\}},\\
&a_7 = \gamma_{\{101,102,105,106,107,110,111,112,113,114,116,\ldots,121,137,139,140,144,146,147,150\}},\\
&a_9 = \gamma_{\{60,61,69,70,72,73,78,82,83,85,90,91,95,97,131,132,135,137\}},\\
&a_{10} = \gamma_{\{99,100,103,104,107,\ldots,118,121,138,139,140,141,142,143,146,148,149,150\}}.
\end{aligned}$$

From theses equalities, we get

$$(6.5.5.3)\quad \begin{cases} a_6 = a_7 = a_8 = a_9 = a_{10} = 0,\ \gamma_j = 0,\ j = 60, 61, 62, \\ 73, 78, 82, 83, 84, 85, 86, 95, 97, 98, 123, 132 = 0, \\ \gamma_{\{63,88,125,134\}} = \gamma_{\{69,99,103,142\}} = \gamma_{\{91,149\}} = 0, \\ \gamma_{\{108,109,115,135,141\}} = \gamma_{\{122,129\}} = \gamma_{\{64,67,87,128,136\}} = 0, \\ \gamma_{\{70,100,104,143\}} = \gamma_{\{90,148\}} = \gamma_{\{72,145\}} = \gamma_{\{122,123,125,128\}} = 0, \\ \gamma_{\{71,101,105,144\}} = \gamma_{\{74,102,106,145\}} = \gamma_{\{92,119,120,147\}} = 0. \end{cases}$$

With the aid of (6.5.5.2) and (6.5.5.3), the homomorphisms g_1, g_2 send (6.5.5.1) respectively to

$$\begin{aligned} &\gamma_{\{69,88,125,135\}} w_{t,1,4} + a_{11} w_{t,1,7} + \gamma_{69} w_{t,1,8} + \gamma_{99} w_{t,1,9} + \gamma_{141} w_{t,1,10} \\ &\quad + \gamma_{103} w_{t,1,11} + \gamma_{\{88,108\}} w_{t,1,12} + \gamma_{109} w_{t,1,13} + \gamma_{115} w_{t,1,14} = 0, \\ &\gamma_{\{70,72,90,128,137\}} w_{t,1,4} + a_{12} w_{t,1,7} + \gamma_{70} w_{t,1,8} + \gamma_{100} w_{t,1,9} + \gamma_{\{140,147\}} w_{t,1,10} \\ &\quad + \gamma_{104} w_{t,1,11} + \gamma_{107} w_{t,1,12} + \gamma_{113} w_{t,1,13} + \gamma_{116} w_{t,1,14} = 0, \end{aligned}$$

where

$$\begin{aligned} a_{11} &= \gamma_{\{63,88,99,103,108,109,115,134,135,141\}}, \\ a_{12} &= \gamma_{\{64,67,72,87,90,100,104,107,113,116,136,137,140,147\}}. \end{aligned}$$

From this, we obtain

$$(6.5.5.4)\quad \begin{cases} a_{11} = a_{12} = 0, \quad \gamma_j = 0,\ j = 69, 70, 99, \\ 100, 103, 104, 107, 109, 113, 115, 116, 141, \\ \gamma_{\{69,88,125,135\}} = \gamma_{\{88,108\}} = 0, \\ \gamma_{\{70,72,90,128,137\}} = \gamma_{\{140,147\}} = 0. \end{cases}$$

With the aid of (6.5.5.2), (6.5.5.3) and (6.5.5.4), the homomorphisms g_3, g_4 send (6.5.5.1) respectively to

$$\begin{aligned} &\gamma_{\{71,74,92,122,138\}} w_{t,1,4} + \gamma_{\{122,125,128\}} w_{t,1,6} + a_{13} w_{t,1,7} \\ &\quad + \gamma_{71} w_{t,1,8} + \gamma_{101} w_{t,1,9} + a_{14} w_{t,1,10} + \gamma_{105} w_{t,1,11} \\ &\quad + a_{15} w_{t,1,12} + \gamma_{\{112,119,150\}} w_{t,1,13} + \gamma_{117} w_{t,1,14} = 0, \\ &\gamma_{\{71,74,92,122,138\}} w_{t,1,4} + \gamma_{\{122,125,128\}} w_{t,1,6} + a_{16} w_{t,1,7} + \gamma_{74} w_{t,1,8} \\ &\quad + \gamma_{102} w_{t,1,9} + a_{17} w_{t,1,10} + \gamma_{106} w_{t,1,11} + \gamma_{\{88,92,111,147,150\}} w_{t,1,12} \\ &\quad + \gamma_{\{114,119,120,121,150\}} w_{t,1,13} + \gamma_{118} w_{t,1,14} = 0, \end{aligned}$$

where

$$\begin{aligned} a_{13} &= \gamma_{\{63,64,67,72,87,90,101,102,105,106,110,111,112,114,117,\ldots,121,134,136,138,139,147\}}, \\ a_{14} &= \gamma_{\{72,74,90,102,106,111,114,118,139\}}, \quad a_{15} = \gamma_{\{88,92,110,120,121,147,150\}}, \\ a_{16} &= \gamma_{\{63,64,67,72,87,90,101,102,105,106,108,110,111,112,114,117,\ldots,121,135,137,\ldots,140,147\}}, \\ a_{17} &= \gamma_{\{63,64,67,71,87,101,105,108,110,112,117,139,140\}}. \end{aligned}$$

From these equalities, we get

$$\begin{cases} a_{13} = a_{14} = a_{15} = a_{16} = a_{17} = 0, \\ \gamma_j = 0,\ j = 71, 74, 101, 102, 105, 106, 117, 118, \\ \gamma_{\{71,74,92,122,138\}} = \gamma_{\{122,125,128\}} = 0, \\ \gamma_{\{112,119,150\}} = \gamma_{\{71,74,92,122,138\}} = 0, \\ \gamma_{\{122,125,128\}} = \gamma_{\{88,92,111,147,150\}} = \gamma_{\{114,119,120,121,150\}} = 0. \end{cases} \tag{6.5.5.5}$$

With the aid of (6.5.5.2), (6.5.5.3), (6.5.5.4) and (6.5.5.5) the homomorphism h sends (6.5.5.1) to

$$\begin{aligned} &a_{18}w_{t,1,7} + \gamma_{\{88,111,114,139\}}w_{t,1,8} + \gamma_{92}w_{t,1,9} + \gamma_{119}w_{t,1,10} \\ &\quad + a_{19}w_{t,1,11} + \gamma_{\{112,114,119,150\}}w_{t,1,12} + \gamma_{120}w_{t,1,13} + \gamma_{121}w_{t,1,14} = 0, \end{aligned}$$

where

$$\begin{aligned} a_{18} &= \gamma_{\{92,111,114,119,120,121,136,137,138,140,150\}}, \\ a_{19} &= \gamma_{\{88,92,110,111,120,121,140,150\}}. \end{aligned}$$

This equality implies

$$\begin{cases} a_{18} = a_{19} = \gamma_{92} = \gamma_{119} = \gamma_{120} = \gamma_{121} = 0, \\ \gamma_{\{88,111,114,139\}} = \gamma_{\{112,114,119,150\}} = 0. \end{cases} \tag{6.5.5.6}$$

Combining (6.5.5.2) - (6.5.5.6), we get $\gamma_j = 0$ for all j. The proposition is proved. □

Proposition 6.5.6. *For any $t \geqslant 4$, the elements $[a_{t,2,j}]$, $57 \leqslant j \leqslant 195$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+3}+5}$.*

Proof. Suppose there is a linear relation

$$\sum_{j=57}^{195} \gamma_j [a_{t,2,j}] = 0, \tag{6.5.6.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_i, 1 \leqslant i \leqslant 6$, to the relation (6.5.6.1), we get

$$\begin{aligned} &\gamma_{150}w_{t,2,1} + \gamma_{151}w_{t,2,2} + \gamma_{124}w_{t,2,3} + \gamma_{165}w_{t,2,4} + \gamma_{126}w_{t,2,5} \\ &\quad + \gamma_{127}w_{t,2,6} + \gamma_{133}w_{t,2,7} + \gamma_{57}w_{t,2,8} + \gamma_{65}w_{t,2,9} + \gamma_{66}w_{t,2,10} \\ &\quad + \gamma_{75}w_{t,2,11} + \gamma_{76}w_{t,2,12} + \gamma_{89}w_{t,2,13} + \gamma_{93}w_{t,2,14} = 0, \\ &\gamma_{152}w_{t,2,1} + \gamma_{154}w_{t,2,2} + \gamma_{\{123,191\}}w_{t,2,3} + \gamma_{166}w_{t,2,4} + \gamma_{\{125,192\}}w_{t,2,5} \\ &\quad + \gamma_{130}w_{t,2,6} + \gamma_{134}w_{t,2,7} + \gamma_{58}w_{t,2,8} + \gamma_{63}w_{t,2,9} + \gamma_{68}w_{t,2,10} \\ &\quad + \gamma_{79}w_{t,2,11} + \gamma_{77}w_{t,2,12} + \gamma_{88}w_{t,2,13} + \gamma_{94}w_{t,2,14} = 0, \\ &\gamma_{153}w_{t,2,1} + \gamma_{155}w_{t,2,2} + \gamma_{\{122,190,195\}}w_{t,2,3} + \gamma_{167}w_{t,2,4} + \gamma_{\{128,193\}}w_{t,2,5} \\ &\quad + \gamma_{\{129,194\}}w_{t,2,6} + \gamma_{136}w_{t,2,7} + \gamma_{59}w_{t,2,8} + \gamma_{64}w_{t,2,9} + \gamma_{67}w_{t,2,10} \\ &\quad + \gamma_{80}w_{t,2,11} + \gamma_{81}w_{t,2,12} + \gamma_{87}w_{t,2,13} + \gamma_{96}w_{t,2,14} = 0, \end{aligned}$$

$$\begin{aligned}
&\gamma_{156}w_{t,2,1} + \gamma_{\{163,184,188,191\}}w_{t,2,2} + \gamma_{159}w_{t,2,3} + a_1w_{t,2,4} + \gamma_{168}w_{t,2,5}\\
&\quad + \gamma_{132}w_{t,2,6} + \gamma_{135}w_{t,2,7} + \gamma_{69}w_{t,2,8} + \gamma_{60}w_{t,2,9} + \gamma_{73}w_{t,2,10}\\
&\quad + \gamma_{82}w_{t,2,11} + \gamma_{91}w_{t,2,12} + \gamma_{78}w_{t,2,13} + \gamma_{95}w_{t,2,14} = 0,\\
&\gamma_{157}w_{t,2,1} + a_2w_{t,2,2} + \gamma_{160}w_{t,2,3} + a_3w_{t,2,4} + \gamma_{169}w_{t,2,5}\\
&\quad + \gamma_{\{131,195\}}w_{t,2,6} + \gamma_{137}w_{t,2,7} + \gamma_{70}w_{t,2,8} + \gamma_{61}w_{t,2,9} + \gamma_{72}w_{t,2,10}\\
&\quad + \gamma_{83}w_{t,2,11} + \gamma_{90}w_{t,2,12} + \gamma_{85}w_{t,2,13} + \gamma_{97}w_{t,2,14} = 0,\\
&a_4w_{t,2,1} + \gamma_{158}w_{t,2,2} + \gamma_{161}w_{t,2,3} + a_5w_{t,2,4} + a_6w_{t,2,5}\\
&\quad + \gamma_{170}w_{t,2,6} + a_7w_{t,2,7} + \gamma_{71}w_{t,2,8} + \gamma_{74}w_{t,2,9} + \gamma_{62}w_{t,2,10}\\
&\quad + \gamma_{92}w_{t,2,11} + \gamma_{84}w_{t,2,12} + \gamma_{86}w_{t,2,13} + \gamma_{98}w_{t,2,14} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{99,103,108,109,115,141,143,171,175,179,192\}}, \quad a_2 = \gamma_{\{162,185,187,190,194\}},\\
&a_3 = \gamma_{\{100,104,107,113,116,140,144,172,176,180,193\}}, \quad a_4 = \gamma_{\{164,183,186,190,191,192,193\}},\\
&a_5 = \gamma_{\{101,105,110,112,117,139,145,173,177,181,194\}},\\
&a_6 = \gamma_{\{102,106,111,114,118,142,146,174,178,182,195\}},\\
&a_7 = \gamma_{\{119,120,121,138,147,148,149,189\}}.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
a_1 = a_2 = a_3 = a_4 = a_5 = a_6 = a_7 = 0,\\
\gamma_j = 0, \ j = 57, \ldots, 98, 124, 126, 127,\\
130, 132 \ldots, 137, 150, \ldots, 161, 165, \ldots, 170,\\
\gamma_{\{123,191\}} = \gamma_{\{125,192\}} = \gamma_{\{122,190,195\}} = \gamma_{\{128,193\}} = 0,\\
\gamma_{\{129,194\}} = \gamma_{\{131,195\}} = \gamma_{\{163,184,188,191\}} = 0.
\end{cases} \tag{6.5.6.2}$$

With the aid of (6.5.6.2), the homomorphisms g_1, g_2 send (6.5.6.1) respectively to

$$\begin{aligned}
&\gamma_{163}w_{t,2,2} + \gamma_{184}w_{t,2,3} + \gamma_{171}w_{t,2,4} + \gamma_{175}w_{t,2,5} + \gamma_{188}w_{t,2,6}\\
&\quad + \gamma_{179}w_{t,2,7} + \gamma_{143}w_{t,2,8} + \gamma_{99}w_{t,2,9} + \gamma_{141}w_{t,2,10}\\
&\quad + \gamma_{103}w_{t,2,11} + \gamma_{108}w_{t,2,12} + \gamma_{109}w_{t,2,13} + \gamma_{115}w_{t,2,14} = 0,\\
&\gamma_{\{122,162,190\}}w_{t,2,2} + \gamma_{185}w_{t,2,3} + \gamma_{172}w_{t,2,4} + \gamma_{176}w_{t,2,5}\\
&\quad + \gamma_{187}w_{t,2,6} + \gamma_{180}w_{t,2,7} + \gamma_{144}w_{t,2,8} + \gamma_{100}w_{t,2,9} + \gamma_{140}w_{t,2,10}\\
&\quad + \gamma_{104}w_{t,2,11} + \gamma_{107}w_{t,2,12} + \gamma_{113}w_{t,2,13} + \gamma_{116}w_{t,2,14} = 0.
\end{aligned}$$

From these equalities, we obtain

$$\begin{cases}
\gamma_j = 0, \ j = 99, 100, 103, 104, 107, 108, 109,\\
113, 115, 116, 140, 141, 143, 144, 163, 171,\\
172, 175, 176, 179, 180, 184, 185, 187, 188,\\
\gamma_{\{122,162,190\}} = 0.
\end{cases} \tag{6.5.6.3}$$

With the aid of (6.5.6.2) and (6.5.6.3), the homomorphisms g_3, g_4 send (6.5.6.1) respectively to

$$\begin{aligned}
&a_8w_{t,2,2} + \gamma_{\{178,183\}}w_{t,2,3} + \gamma_{\{138,173\}}w_{t,2,4} + \gamma_{\{131,174,178,186\}}w_{t,2,6}\\
&\quad + \gamma_{177}w_{t,2,5} + \gamma_{\{138,181,189\}}w_{t,2,7} + \gamma_{145}w_{t,2,8} + \gamma_{\{139,189\}}w_{t,2,10}\\
&\quad + \gamma_{101}w_{t,2,9} + \gamma_{105}w_{t,2,11} + \gamma_{110}w_{t,2,12} + \gamma_{112}w_{t,2,13} + \gamma_{117}w_{t,2,14} = 0,\\
&a_9w_{t,2,2} + \gamma_{\{123,177,183\}}w_{t,2,3} + \gamma_{\{138,174\}}w_{t,2,4} + \gamma_{178}w_{t,2,5}\\
&\quad + a_{10}w_{t,2,6} + a_{11}w_{t,2,7} + \gamma_{146}w_{t,2,8} + \gamma_{102}w_{t,2,9} + a_{12}w_{t,2,10}\\
&\quad + \gamma_{106}w_{t,2,11} + \gamma_{111}w_{t,2,12} + \gamma_{114}w_{t,2,13} + \gamma_{118}w_{t,2,14} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_8 = \gamma_{\{122,131,164,174,190\}}, \quad a_9 = \gamma_{\{129,162,164,173,190\}},\\
&a_{10} = \gamma_{\{125,128,129,173,177,186\}}, \quad a_{11} = \gamma_{\{119,120,121,138,147,148,149,182,189\}},\\
&a_{12} = \gamma_{\{119,120,121,142,147,148,149,189\}}.
\end{aligned}$$

Computing directly from these equalities, we get

$$\begin{cases}
a_8 = a_9 = a_{10} = a_{11} = a_{12} = 0,\\
\gamma_j = 0,\ j = 101, 102, 105, 106, 110, 111,\\
112, 114, 117, 118, 145, 146, 177, 178,\\
\gamma_{\{178,183\}} = \gamma_{\{138,173\}} = \gamma_{\{131,174,178,186\}} = \gamma_{\{139,189\}} = 0,\\
\gamma_{\{138,181,189\}} = \gamma_{\{123,177,183\}} = \gamma_{\{138,174\}} = 0.
\end{cases} \tag{6.5.6.4}$$

With the aid of (6.5.6.2), (6.5.6.3) and (6.5.6.4), the homomorphism h sends (6.5.6.1) to

$$\begin{aligned}
&\gamma_{\{122,138,162,164,190\}}w_{t,2,1} + \gamma_{\{138,186\}}w_{t,2,4} + \gamma_{\{138,181,182\}}w_{t,2,7}\\
&\quad + \gamma_{142}w_{t,2,8} + \gamma_{147}w_{t,2,9} + \gamma_{119}w_{t,2,10} + \gamma_{148}w_{t,2,11}\\
&\quad + \gamma_{149}w_{t,2,12} + \gamma_{120}w_{t,2,13} + \gamma_{121}w_{t,2,14} = 0.
\end{aligned}$$

From this, it implies

$$\begin{cases}
\gamma_{119} = \gamma_{120} = \gamma_{121} = \gamma_{142} = \gamma_{147} = \gamma_{148} = \gamma_{149} = 0,\\
\gamma_{\{122,138,162,164,190\}} = \gamma_{\{138,186\}} = \gamma_{\{138,181,182\}} = 0.
\end{cases} \tag{6.5.6.5}$$

Combining (6.5.6.2), (6.5.6.3), (6.5.6.4) and (6.5.6.5), we obtain $\gamma_j = 0$ for all j. The proposition is proved. □

Proposition 6.5.7. *For any $t \geqslant 4$ and $s \geqslant 3$, the elements $[a_{t,s,j}]$, $57 \leqslant j \leqslant 210$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+1}-3}$.*

Proof. Suppose there is a linear relation

$$\sum_{j=57}^{210} \gamma_j[a_{t,s,j}] = 0, \tag{6.5.7.1}$$

where $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms $f_i, 1 \leqslant i \leqslant 6$ to the relation (6.5.7.1) and we obtain

$$\begin{aligned}
&\gamma_{150}w_{t,s,1} + \gamma_{151}w_{t,s,2} + \gamma_{124}w_{t,s,3} + \gamma_{165}w_{t,s,4} + \gamma_{126}w_{t,s,5}\\
&\quad + \gamma_{127}w_{t,s,6} + \gamma_{133}w_{t,s,7} + \gamma_{57}w_{t,s,8} + \gamma_{65}w_{t,s,9} + \gamma_{66}w_{t,s,10}\\
&\quad + \gamma_{75}w_{t,s,11} + \gamma_{76}w_{t,s,12} + \gamma_{89}w_{t,s,13} + \gamma_{93}w_{t,s,14} = 0,\\
&\gamma_{152}w_{t,s,1} + \gamma_{154}w_{t,s,2} + \gamma_{123}w_{t,s,3} + \gamma_{166}w_{t,s,4} + \gamma_{125}w_{t,s,5}\\
&\quad + \gamma_{130}w_{t,s,6} + \gamma_{134}w_{t,s,7} + \gamma_{58}w_{t,s,8} + \gamma_{63}w_{t,s,9} + \gamma_{68}w_{t,s,10}\\
&\quad + \gamma_{79}w_{t,s,11} + \gamma_{77}w_{t,s,12} + \gamma_{88}w_{t,s,13} + \gamma_{94}w_{t,s,14} = 0,\\
&\gamma_{153}w_{t,s,1} + \gamma_{155}w_{t,s,2} + a_1w_{t,s,3} + \gamma_{167}w_{t,s,4} + \gamma_{128}w_{t,s,5}\\
&\quad + \gamma_{129}w_{t,s,6} + \gamma_{136}w_{t,s,7} + \gamma_{59}w_{t,s,8} + \gamma_{64}w_{t,s,9} + \gamma_{67}w_{t,s,10}\\
&\quad + \gamma_{80}w_{t,s,11} + \gamma_{81}w_{t,s,12} + \gamma_{87}w_{t,s,13} + \gamma_{96}w_{t,s,14} = 0,\\
&\gamma_{156}w_{t,s,1} + \gamma_{163}w_{t,s,2} + \gamma_{159}w_{t,s,3} + \gamma_{171}w_{t,s,4} + \gamma_{168}w_{t,s,5}\\
&\quad + \gamma_{132}w_{t,s,6} + \gamma_{135}w_{t,s,7} + \gamma_{69}w_{t,s,8} + \gamma_{60}w_{t,s,9} + \gamma_{73}w_{t,s,10}\\
&\quad + \gamma_{82}w_{t,s,11} + \gamma_{91}w_{t,s,12} + \gamma_{78}w_{t,s,13} + \gamma_{95}w_{t,s,14} = 0,\\
&\gamma_{157}w_{t,s,1} + a_2w_{t,s,2} + \gamma_{160}w_{t,s,3} + \gamma_{172}w_{t,s,4} + \gamma_{169}w_{t,s,5}\\
&\quad + \gamma_{131}w_{t,s,6} + \gamma_{137}w_{t,s,7} + \gamma_{70}w_{t,s,8} + \gamma_{61}w_{t,s,9} + \gamma_{72}w_{t,s,10}\\
&\quad + \gamma_{83}w_{t,s,11} + \gamma_{90}w_{t,s,12} + \gamma_{85}w_{t,s,13} + \gamma_{97}w_{t,s,14} = 0,\\
&a_3w_{t,s,1} + \gamma_{158}w_{t,s,2} + \gamma_{161}w_{t,s,3} + \gamma_{173}w_{t,s,4} + \gamma_{174}w_{t,s,5}\\
&\quad + \gamma_{170}w_{t,s,6} + \gamma_{138}w_{t,s,7} + \gamma_{71}w_{t,s,8} + \gamma_{74}w_{t,s,9} + \gamma_{62}w_{t,s,10}\\
&\quad + \gamma_{92}w_{t,s,11} + \gamma_{84}w_{t,s,12} + \gamma_{86}w_{t,s,13} + \gamma_{98}w_{t,s,14} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases} \gamma_{\{122,209\}}, & s = 3,\\ \gamma_{122}, & s \geqslant 4, \end{cases} \quad a_2 = \begin{cases} \gamma_{\{162,209\}}, & s = 3,\\ \gamma_{162}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{164,208,209,210\}}, & s = 3,\\ \gamma_{164}, & s \geqslant 4. \end{cases}$$

From this, it implies

$$\begin{cases} a_1 = a_2 = a_3 = 0,\\ \gamma_j = 0,\ j = 57, \ldots, 98, 123, \ldots, 138,\\ 150, \ldots, 161, 163, 165, \ldots, 174. \end{cases} \tag{6.5.7.2}$$

With the aid of (6.5.7.2), the homomorphisms $g_i,\ i = 1, 2, 3, 4$, send (6.5.7.1) respectively to

$$\begin{aligned}
&\gamma_{190}w_{t,s,1} + \gamma_{195}w_{t,s,2} + \gamma_{184}w_{t,s,3} + \gamma_{199}w_{t,s,4} + \gamma_{175}w_{t,s,5}\\
&\quad + \gamma_{188}w_{t,s,6} + \gamma_{179}w_{t,s,7} + \gamma_{143}w_{t,s,8} + \gamma_{99}w_{t,s,9} + \gamma_{141}w_{t,s,10}\\
&\quad + \gamma_{103}w_{t,s,11} + \gamma_{108}w_{t,s,12} + \gamma_{109}w_{t,s,13} + \gamma_{115}w_{t,s,14} = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_{191}w_{t,s,1} + a_4w_{t,s,2} + \gamma_{185}w_{t,s,3} + \gamma_{200}w_{t,s,4} + \gamma_{176}w_{t,s,5}\\
&\quad + \gamma_{187}w_{t,s,6} + \gamma_{180}w_{t,s,7} + \gamma_{144}w_{t,s,8} + \gamma_{100}w_{t,s,9} + \gamma_{140}w_{t,s,10}\\
&\quad + \gamma_{104}w_{t,s,11} + \gamma_{107}w_{t,s,12} + \gamma_{113}w_{t,s,13} + \gamma_{116}w_{t,s,14} = 0,\\
&\gamma_{192}w_{t,s,1} + a_5w_{t,s,2} + a_6w_{t,s,3} + \gamma_{201}w_{t,s,4} + \gamma_{177}w_{t,s,5}\\
&\quad + a_7w_{t,s,6} + \gamma_{181}w_{t,s,7} + \gamma_{145}w_{t,s,8} + \gamma_{101}w_{t,s,9} + \gamma_{139}w_{t,s,10}\\
&\quad + \gamma_{105}w_{t,s,11} + \gamma_{110}w_{t,s,12} + \gamma_{112}w_{t,s,13} + \gamma_{117}w_{t,s,14} = 0,\\
&\gamma_{193}w_{t,s,1} + a_8w_{t,s,2} + a_9w_{t,s,3} + \gamma_{202}w_{t,s,4} + \gamma_{178}w_{t,s,5}\\
&\quad + a_{10}w_{t,s,6} + \gamma_{182}w_{t,s,7} + \gamma_{146}w_{t,s,8} + \gamma_{102}w_{t,s,9} + \gamma_{142}w_{t,s,10}\\
&\quad + \gamma_{106}w_{t,s,11} + \gamma_{111}w_{t,s,12} + \gamma_{114}w_{t,s,13} + \gamma_{118}w_{t,s,14} = 0,
\end{aligned}$$

where

$$a_4 = \begin{cases} \gamma_{\{122,194\}}, & s = 3,\\ \gamma_{194}, & s \geqslant 4, \end{cases} \quad a_5 = \begin{cases} \gamma_{\{164,196\}}, & s = 3,\\ \gamma_{196}, & s \geqslant 4, \end{cases}$$

$$a_6 = \begin{cases} \gamma_{\{183,208\}}, & s = 3,\\ \gamma_{183}, & s \geqslant 4, \end{cases} \quad a_7 = \begin{cases} \gamma_{\{186,210\}}, & s = 3,\\ \gamma_{186}, & s \geqslant 4, \end{cases}$$

$$a_8 = \begin{cases} \gamma_{\{164,197,204,205\}}, & s = 3,\\ \gamma_{197}, & s \geqslant 4, \end{cases} \quad a_9 = \begin{cases} \gamma_{\{198,204,206,208\}}, & s = 3,\\ \gamma_{198}, & s \geqslant 4, \end{cases}$$

$$a_{10} = \begin{cases} \gamma_{\{203,205,206,210\}}, & s = 3,\\ \gamma_{203}, & s \geqslant 4. \end{cases}$$

Hence, we obtain

$$\text{(6.5.7.3)} \qquad \begin{cases} a_i = 0, i = 4, 5, \ldots, 10,\\ \gamma_j = 0,\ j = 99, \ldots, 118, 139, \ldots, 146, 175, \ldots, 182,\\ 184, 185, 187, 188, 190, \ldots, 193, 195, 199, \ldots, 202. \end{cases}$$

With the aid of (6.5.7.2) and (6.5.7.3), the homomorphism h sends (6.5.7.1) to

$$\begin{aligned}
&a_{11}w_{t,s,1} + a_{12}w_{t,s,2} + \gamma_{204}w_{t,s,3} + a_{13}w_{t,s,4} + \gamma_{205}w_{t,s,5}\\
&\quad + \gamma_{206}w_{t,s,6} + \gamma_{207}w_{t,s,7} + \gamma_{189}w_{t,s,8} + \gamma_{147}w_{t,s,9} + \gamma_{119}w_{t,s,10}\\
&\quad + \gamma_{148}w_{t,s,11} + \gamma_{149}w_{t,s,12} + \gamma_{120}w_{t,s,13} + \gamma_{121}w_{t,s,14} = 0,
\end{aligned}$$

where

$$a_{11} = \begin{cases} \gamma_{197}, & s = 3,\\ \gamma_{208}, & s \geqslant 4, \end{cases} \quad a_{12} = \begin{cases} \gamma_{198}, & s = 3,\\ \gamma_{209}, & s \geqslant 4, \end{cases} \quad a_{13} = \begin{cases} \gamma_{203}, & s = 3,\\ \gamma_{210}, & s \geqslant 4. \end{cases}$$

From these equalities, we get

$$\text{(6.5.7.4)} \qquad \begin{cases} a_{11} = a_{12} = a_{13} = 0,\ \gamma_j = 0,\ j = 119, 120,\\ 121, 147, 148, 149, 189, 204, 205, 206, 207. \end{cases}$$

Combining (6.5.7.2), (6.5.7.3) and (6.5.7.4), we obtain $\gamma_j = 0$ for all j. The proposition is proved. □

7. The indecomposables of P_4 in degree $2^{s+t} + 2^s - 2$

7.1. The τ-sequence of the admissible monomials.

In this subsection, we prove the following.

Lemma 7.1.1. *Let x be an admissible monomial of degree $2^{s+t} + 2^s - 2$ in P_4. We have*

1. *If $s = 1$ and $t = 1$ then either $\tau(x) = (2;1)$ or $\tau(x) = (4;0)$.*
2. *If $s = 1$ and $t = 2$ then either $\tau(x) = (2;1;1)$ or $\tau(x) = (2;3)$ or $\tau(x) = (4;2)$.*
3. *If $s = 1$ and $t = 3$ then either $\tau(x) = (2;1;1;1)$ or $\tau(x) = (2;3;2)$ or $\tau(x) = (4;2;2)$ or $\tau(x) = (4;4;1)$.*
4. *If $s = 1$ and $t \geqslant 4$ then $\tau(x)$ is one of the following sequences*

$$(2;\underbrace{1;1;\ldots;1}_{t\ times}),\quad (4;\underbrace{2;2;\ldots;2}_{t-1\ times}),\quad (4;4;\underbrace{3;3;\ldots;3}_{t-3\ times};1).$$

5. *If $s \geqslant 2$ then $\tau(x)$ is one of the following sequences*

$$(\underbrace{2;2;\ldots;2}_{s\ times};\underbrace{1;1;\ldots;1}_{t\ times}),\quad (4;\underbrace{3;3;\ldots;3}_{s-2\ times};\underbrace{1;1;\ldots;1}_{t+1\ times}),$$
$$(4;\underbrace{3;3;\ldots;3}_{s-1\ times};\underbrace{2;2;\ldots;2}_{t-1\ times}).$$

We need the following for the proof of this lemma. From Lemmas 4.4, 2.6, Theorem 2.5 and the results in Section 6, we get

Lemma 7.1.2. *Let x be a monomial of degree $2^{s+1} + 2^s - 3$ in P_4 with $s \geqslant 2$. If x is inadmissible then there is a strictly inadmissible matrix Δ such that $\Delta \triangleright x$.*

Lemma 7.1.3. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}0&1&0&1\\1&0&1&1\end{pmatrix}\quad \begin{pmatrix}0&1&1&0\\1&0&1&1\end{pmatrix}\quad \begin{pmatrix}0&0&1&1\\1&1&0&1\end{pmatrix}\quad \begin{pmatrix}0&0&1&1\\1&1&1&0\end{pmatrix}$$
$$\begin{pmatrix}0&1&1&0\\1&1&0&1\end{pmatrix}\quad \begin{pmatrix}0&1&0&1\\1&1&1&0\end{pmatrix}\quad \begin{pmatrix}1&0&1&0\\1&1&0&1\end{pmatrix}\quad \begin{pmatrix}1&0&0&1\\1&1&1&0\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (2,1,2,3), (2,1,3,2), (2,2,1,3), (2,2,3,1), (2,3,1,2), (2,3,2,1), (3,2,1,2), (3,2,2,1). We prove the lemma for the matrix associated with the monomial (3,2,1,2). The others can be obtained by a similar argument. Obviously, we have

$$(3,2,1,2) = Sq^1(3,1,1,2) + (4,1,1,2) + (3,1,2,2).$$

The lemma is proved. □

Lemma 7.1.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&0&0&1\\0&1&1&1\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\0&1&1&1\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&0&0&1\\0&1&1&1\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\0&1&1&1\\0&1&1&0\end{pmatrix}$$
$$\begin{pmatrix}1&1&0&0\\0&1&1&1\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\0&1&1&1\\0&1&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&1\\1&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&1\\1&0&1&0\end{pmatrix}$$
$$\begin{pmatrix}1&0&1&0\\0&1&1&1\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&0&0&1\\0&1&1&1\\0&1&1&0\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (1,2,6,7), (1,2,7,6), (1,6,2,7), (1,6,7,2), (1,7,2,6), (1,7,6,2), (7,1,2,6), (7,1,6,2), (1,6,3,6), (1,6,6,3). It suffices to prove the lemma for the matrices associated with the monomials (1,2,6,7), (1,6,3,6). A simple computation shows

$$\begin{aligned}(1,2,6,7) &= Sq^1(2,1,5,7) + Sq^2\big((1,1,5,7)+(1,1,3,9)\big) \bmod \mathcal{L}_4(2;3;2),\\ (1,6,3,6) &= Sq^1\big((2,5,3,5)+(2,3,5,5)\big) + Sq^2\big((1,5,3,5)\\ &\quad + (1,3,5,5)\big) + (1,3,6,6) \bmod \mathcal{L}_4(2;3;2).\end{aligned}$$

The lemma follows. □

Lemma 7.1.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&0&0&1\\1&0&0&1\\0&1&1&1\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\1&0&1&0\\0&1&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&1&0&0\\0&1&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&1&0&0\\1&0&1&1\end{pmatrix}$$
$$\begin{pmatrix}1&0&1&0\\1&0&0&1\\0&1&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&0&1\\0&1&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&0\\0&1&1&1\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (3,4,4,7), (3,4,7,4), (3,7,4,4), (7,3,4,4), (3,4,5,6), (3,5,4,6), (3,5,6,4). It suffices to prove the lemma for the matrices associated with the monomials (3,4,4,7), (3,4,5,6). By a direct computation, we have

$$\begin{aligned}(3,4,4,7) &= Sq^1(3,1,2,11) + Sq^2(5,2,2,7)\\ &\quad + Sq^4(3,2,2,7) \bmod \mathcal{L}_4(2;2;3),\\ (3,4,5,6) &= Sq^1(3,1,3,10) + Sq^2(5,2,3,6)\\ &\quad + Sq^4(3,2,3,6) \bmod \mathcal{L}_4(2;2;3).\end{aligned}$$

The lemma follows. □

Lemma 7.1.6. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&1&0&0\\0&1&1&1\\0&0&1&1\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&1\\0&0&1&1\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&1\\0&1&0&1\\0&1&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&1\\0&1&1&0\\0&1&1&0\end{pmatrix}$$
$$\begin{pmatrix}1&1&0&0\\1&0&1&1\\0&1&0&1\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&1\\0&1&1&0\\0&0&1&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&1\\0&1&1&0\\0&1&0&1\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(1,3,14,14),(3,1,14,14),(3,13,2,14),(3,13,14,2),$$
$$(3,5,10,14),(3,5,14,10),(3,13,6,10).$$

It suffices to prove the lemma for the matrices associated with the monomials (3,1,14,14), (3,13,2,14), (3,5,10,14), (3,13,6,10). By a direct computation, we have

$$\begin{aligned}(3,1,14,14) &= Sq^1\big((3,1,13,14)+(3,1,14,13)\big)+Sq^2\big((5,1,11,13)\\ &\quad+(5,1,7,17)\big)+Sq^4\big((3,1,11,13)+(3,1,7,17)\big) \mod \mathcal{L}_4(2;3;2;2).\\ (3,13,2,14) &= Sq^1\big((3,13,2,13)+(3,13,1,14)\big)+Sq^2\big((5,11,1,13)\\ &\quad+(5,7,1,17)\big)+Sq^4\big((3,11,1,13)+(3,7,1,17)\big) \mod \mathcal{L}_4(2;3;2;2).\\ (3,5,10,14) &= Sq^1\big((3,5,10,13)+(3,5,9,14)\big)+Sq^2\big((5,3,9,13)\\ &\quad+(5,3,5,17)\big)+Sq^4\big((3,3,9,13)+(3,3,5,17)\big) \mod \mathcal{L}_4(2;3;2;2).\\ (3,13,6,10) &= Sq^1\big((3,13,6,9)+(3,13,5,10)\big)+Sq^2\big((5,11,5,9)\\ &\quad+(5,7,9,9)\big)+Sq^4\big((3,11,5,9)+(3,7,9,9)\big) \mod \mathcal{L}_4(2;3;2;2).\end{aligned}$$

The lemma follows. □

Proof of Lemma 7.1.1. Observe that $z=(2^{s+t}-1,2^s-1,0,0)$ is the minimal spike of degree $2^{s+t}+2^s-2$ and $\tau(z)=(\underbrace{2;2;\ldots;2}_{s\text{ times}};\underbrace{1;1;\ldots;1}_{t\text{ times}})$. Since $2^{s+t}+2^s-2$ is even and x is admissible, using Theorem 2.13, we get either $\tau_1(x)=2$ or $\tau_1(x)=4$. Let $M=(\varepsilon_{ij}(x)), i\geqslant 1, 1\leqslant j\leqslant 4$, be the matrix associated with x. We set $M'=(\varepsilon_{ij}(x)), i\geqslant 2, 1\leqslant j\leqslant 4$ and denote by x' the monomial corresponding to M'.

Suppose $s=1$ and $t=1$. Then $\deg x=4$. If $\tau_1(x)=2$ then $\deg x'=1$. Hence $\tau(x')=(\tau_2(x))=(1;0)$. So $\tau(x)=(2;1)$. If $\tau_1(x)=4$ then $\deg x'=0$. Hence $\tau(x)=(4;0)$. The first part of the lemma holds.

Suppose $s=1$ and $t=2$. Then we have $\deg x=8$. If $\tau_1(x)=2$ then $\deg x'=3$ and x' is admissible. According to Lemma 5.4, either $\tau(x')=(\tau_2(x);\tau_3(x))=(1;1)$, or $\tau(x')=(3;0)$. So, either $\tau(x)=(2;1;1)$

or $\tau(x) = (2;3)$. If $\tau_1(x) = 4$ then $\deg x' = 2$. Hence $\tau(x') = (2;0)$ and $\tau(x) = (4;2)$. The second part of the lemma holds.

Let $s = 1$ and $t = 3$. Then $\deg x = 16$. If $\tau_1(x) = 2$ then $\deg x' = 7$. Since x is admissible, by Lemma 6.1.2 and Theorem 2.5, x' is admissible. According to Lemma 5.4, either $\tau(x') = (\tau_2(x); \tau_3(x)) = (1;1;1)$, or $\tau(x') = (3;2)$, so, either $\tau(x) = (2;1;1;1)$ or $\tau(x) = (2;3;2)$.

If $\tau_1(x) = 4$ then $\deg x' = 6$. Since x is admissible, by Lemma 5.5 and Theorem 2.5, x' is admissible. Using Lemma 4.6, we get either $\tau(x') = (2;2)$ or $\tau(x') = (4;1)$. So, we obtain either $\tau(x) = (4;2;2)$ or $\tau(x) = (4;4;1)$. The lemma holds.

Let $s = 1$ and $t \geqslant 4$. Then $\deg x = 2^{t+1}$. If $\tau_1(x) = 2$ then $\deg x' = 2^t - 1$. Since x is admissible, by Lemma 6.1.2 and Theorem 2.5, x' is admissible. According to Lemma 5.4, either $\tau(x') = (\tau_2(x); \tau_3(x); \ldots) = (\underbrace{1;1;\ldots;1}_{t \text{ times}})$ or $\tau(x') = (3; \underbrace{2;2;\ldots;2}_{t-2 \text{ times}})$.

If $\tau(x') = (3; \underbrace{2;2;\ldots;2}_{t-2 \text{ times}})$ then $\tau(x) = (2;3; \underbrace{2;2;\ldots;2}_{t-2 \text{ times}})$. From Lemmas 4.7, 4.8, 4.4, 5.10, 7.1.3, 7.1.4, 7.1.6, we see that x is inadmissible and we have a contradiction. Hence, we get $\tau(x) = (2; \underbrace{1;1;\ldots;1}_{t \text{ times}})$.

If $\tau_1(x) = 4$ then $\deg x' = 2^t - 2$. By Lemma 5.5 and Theorem 2.5, x' is admissible. Using Lemma 4.6, we get either $\tau(x') = (\underbrace{2;2;\ldots;2}_{t-1 \text{ times}})$ or $\tau(x') = (4; \underbrace{3;3;\ldots;3}_{t-3 \text{ times}}; 1)$. The lemma holds.

Suppose that $s \geqslant 2$. If $\tau_1(x) = 2$ then $\deg x' = 2^{s+t-1} + 2^{s-1} - 2$. Using Lemmas 4.7, 4.8, 4.4, 5.10, 7.1.3, 7.1.5, Proposition 2.7 and by an inductive argument we see that x' is admissible and $\tau(x') = (\underbrace{2;2;\ldots;2}_{s-1 \text{ times}}; \underbrace{1;1;\ldots;1}_{t \text{ times}})$. Hence, we get

$$\tau(x) = (\underbrace{2;2;\ldots;2}_{s \text{ times}}; \underbrace{1;1;\ldots;1}_{t \text{ times}}).$$

If $\tau_1(x) = 4$ then $\deg x' = 2^{s+t-1} + 2^{s-1} - 3$. Since x is admissible, by Lemma 7.1.2 and Theorem 2.5, x' is admissible. According to Lemma 5.4 and Lemma 6.1.1, $\tau(x')$ is one of the following sequences

$$(\underbrace{3;3;\ldots;3}_{s-2 \text{ times}}; \underbrace{1;1;\ldots;1}_{t+1 \text{ times}}), \quad (\underbrace{3;3;\ldots;3}_{s-1 \text{ times}}; \underbrace{2;2;\ldots;2}_{t-1 \text{ times}}).$$

The lemma is completely proved. □

7.2. **The case $s = 1$.**

According to Kameko [8], $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{t+1}}$ is an $\mathbb{F}_2$-vector space with basis of the consisting of all the following classes:

For $t \geqslant 1$,

$$\begin{aligned} v_{t,1,1} &= [1, 1, 2^{t+1} - 2], & v_{t,1,2} &= [1, 2^{t+1} - 2, 1],\\ v_{t,1,3} &= [0, 1, 2^{t+1} - 1], & v_{t,1,4} &= [0, 2^{t+1} - 1, 1],\\ v_{t,1,5} &= [1, 0, 2^{t+1} - 1], & v_{t,1,6} &= [1, 2^{t+1} - 1, 0],\\ v_{t,1,7} &= [2^{t+1} - 1, 0, 1], & v_{t,1,8} &= [2^{t+1} - 1, 1, 0]. \end{aligned}$$

For $t \geqslant 2$

$$\begin{aligned} v_{t,1,9} &= [0, 3, 2^{t+1} - 3], & v_{t,1,10} &= [3, 0, 2^{t+1} - 3],\\ v_{t,1,11} &= [3, 2^{t+1} - 3, 0], & v_{t,1,12} &= [1, 2, 2^{t+1} - 3],\\ v_{t,1,13} &= [1, 3, 2^{t+1} - 4], & v_{t,1,14} &= [3, 1, 2^{t+1} - 4]. \end{aligned}$$

For $t = 2$, $v_{2,1,15} = [3, 4, 1]$.

So, we have

Proposition 7.2.1. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{t+1}}$ is an $\mathbb{F}_2$-vector space with a basis consisting of all the classes represented by the following monomials:*

For $t \geqslant 1$,

$$\begin{aligned} b_{t,1,1} &= (0, 1, 1, 2^{t+1} - 2), & b_{t,1,2} &= (0, 1, 2^{t+1} - 2, 1),\\ b_{t,1,3} &= (1, 0, 1, 2^{t+1} - 2), & b_{t,1,4} &= (1, 0, 2^{t+1} - 2, 1),\\ b_{t,1,5} &= (1, 1, 0, 2^{t+1} - 2), & b_{t,1,6} &= (1, 1, 2^{t+1} - 2, 0),\\ b_{t,1,7} &= (1, 2^{t+1} - 2, 0, 1), & b_{t,1,8} &= (1, 2^{t+1} - 2, 1, 0),\\ b_{t,1,9} &= (0, 0, 1, 2^{t+1} - 1), & b_{t,1,10} &= (0, 0, 2^{t+1} - 1, 1),\\ b_{t,1,11} &= (0, 1, 0, 2^{t+1} - 1), & b_{t,1,12} &= (0, 1, 2^{t+1} - 1, 0),\\ b_{t,1,13} &= (0, 2^{t+1} - 1, 0, 1), & b_{t,1,14} &= (0, 2^{t+1} - 1, 1, 0),\\ b_{t,1,15} &= (1, 0, 0, 2^{t+1} - 1), & b_{t,1,16} &= (1, 0, 2^{t+1} - 1, 0),\\ b_{t,1,17} &= (1, 2^{t+1} - 1, 0, 0), & b_{t,1,18} &= (2^{t+1} - 1, 0, 0, 1),\\ b_{t,1,19} &= (2^{t+1} - 1, 0, 1, 0), & b_{t,1,20} &= (2^{t+1} - 1, 1, 0, 0), \end{aligned}$$

For $t \geqslant 2$,

$$\begin{aligned} b_{t,1,21} &= (0, 0, 3, 2^{t+1} - 3), & b_{t,1,22} &= (0, 3, 0, 2^{t+1} - 3).\\ b_{t,1,23} &= (0, 3, 2^{t+1} - 3, 0), & b_{t,1,24} &= (3, 0, 0, 2^{t+1} - 3),\\ b_{t,1,25} &= (3, 0, 2^{t+1} - 3, 0), & b_{t,1,26} &= (3, 2^{t+1} - 3, 0, 0),\\ b_{t,1,27} &= (0, 1, 2, 2^{t+1} - 3), & b_{t,1,28} &= (1, 0, 2, 2^{t+1} - 3),\\ b_{t,1,29} &= (1, 2, 0, 2^{t+1} - 3), & b_{t,1,30} &= (1, 2, 2^{t+1} - 3, 0),\\ b_{t,1,31} &= (0, 1, 3, 2^{t+1} - 4), & b_{t,1,32} &= (0, 3, 1, 2^{t+1} - 4),\\ b_{t,1,33} &= (1, 0, 3, 2^{t+1} - 4), & b_{t,1,34} &= (1, 3, 0, 2^{t+1} - 4),\\ b_{t,1,35} &= (1, 3, 2^{t+1} - 4, 0), & b_{t,1,36} &= (3, 0, 1, 2^{t+1} - 4),\\ b_{t,1,37} &= (3, 1, 0, 2^{t+1} - 4), & b_{t,1,38} &= (3, 1, 2^{t+1} - 4, 0). \end{aligned}$$

For $t = 2$,

$$\begin{aligned} b_{2,1,39} &= (0, 3, 4, 1), & b_{2,1,40} &= (3, 0, 4, 1),\\ b_{2,1,41} &= (3, 4, 0, 1), & b_{2,1,42} &= (3, 4, 1, 0). \end{aligned}$$

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+1}}$. We have

Theorem 7.2.2. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+1}}$ is an $\mathbb{F}_2$-vector space with a basis consisting of all the classes represented by the following monomials:*

1. $\phi(b_{1,j}), 1 \leqslant j \leqslant 6$ *for* $t = 2$.

2. $\phi(b_{t-1,j}), 1 \leqslant j \leqslant \mu_6(t-1)$, $\phi^2(a_{t-2,j}), 1 \leqslant j \leqslant \mu_1(t-2)$ *for* $t \geqslant 3$. *Here* $b_{t-1,j}$, *is defined as in Section 4,* $a_{t-3,j}, \mu_1(t-2)$ *are defined as in Section 3, the homomorphism* ϕ *is defined as in Definition 2.9 and* $\mu_6(1) = 6, \mu_6(2) = 24, \mu_6(t-1) = 35$ *for any* $t \geqslant 4$.

3. For $t = 1$, $c_{1,1,1} = (1,1,1,1)$.

For $t \geqslant 2$,

$$c_{t,1,1} = (1,1,2,2^{t+1}-4), \quad c_{t,1,2} = (1,2,1,2^{t+1}-4),$$
$$c_{t,1,3} = (1,2,2^{t+1}-4,1).$$

For $t = 2$,

$$c_{2,1,4} = (1,2,2,3), \quad c_{2,1,5} = (1,2,3,2),$$
$$c_{2,1,6} = (1,3,2,2), \quad c_{2,1,7} = (3,1,2,2).$$

For $t \geqslant 3$,

$$c_{t,1,4} = (1,2,4,2^{t+1}-7), \quad c_{t,1,5} = (1,2,5,2^{t+1}-8),$$
$$c_{t,1,6} = (1,3,4,2^{t+1}-8), \quad c_{t,1,7} = (3,1,4,2^{t+1}-8),$$

For $t = 3$,

$$c_{3,1,8} = (1,3,6,6), \quad c_{3,1,9} = (3,1,6,6),$$
$$c_{3,1,10} = (3,5,2,6), \quad c_{3,1,11} = (3,5,6,2).$$

We prove this theorem by proving some propositions.

Proposition 7.2.3. *The* $\mathbb{F}_2$*-vector space* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+1}}$ *is generated by the elements listed in Theorem 7.2.2.*

The proof of the proposition is based on the following lemmas.

Lemma 7.2.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 0&0&1&1\\ 0&1&0&0 \end{pmatrix} \quad \begin{pmatrix} 0&0&1&1\\ 1&0&0&0 \end{pmatrix} \quad \begin{pmatrix} 0&1&0&1\\ 1&0&0&0 \end{pmatrix} \quad \begin{pmatrix} 0&1&1&0\\ 1&0&0&0 \end{pmatrix}.$$

Proof. See [8] □

Lemma 7.2.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 0&1&0&1\\ 0&1&0&0\\ 0&0&1&0\\ 0&0&1&0 \end{pmatrix} \quad \begin{pmatrix} 1&0&0&1\\ 1&0&0&0\\ 0&0&1&0\\ 0&0&1&0 \end{pmatrix} \quad \begin{pmatrix} 1&0&0&1\\ 1&0&0&0\\ 0&1&0&0\\ 0&1&0&0 \end{pmatrix} \quad \begin{pmatrix} 1&0&1&0\\ 1&0&0&0\\ 0&1&0&0\\ 0&1&0&0 \end{pmatrix}$$

$$\begin{pmatrix} 0&1&0&1\\ 0&1&0&0\\ 0&0&1&0\\ 0&0&0&1 \end{pmatrix} \quad \begin{pmatrix} 1&0&0&1\\ 1&0&0&0\\ 0&0&1&0\\ 0&0&0&1 \end{pmatrix} \quad \begin{pmatrix} 1&0&0&1\\ 1&0&0&0\\ 0&1&0&0\\ 0&0&0&1 \end{pmatrix} \quad \begin{pmatrix} 1&0&1&0\\ 1&0&0&0\\ 0&1&0&0\\ 0&0&1&0 \end{pmatrix}$$

$$\begin{pmatrix} 0&1&1&0\\ 0&1&0&0\\ 0&0&1&0\\ 0&0&0&1 \end{pmatrix} \quad \begin{pmatrix} 1&0&1&0\\ 1&0&0&0\\ 0&0&1&0\\ 0&0&0&1 \end{pmatrix} \quad \begin{pmatrix} 1&1&0&0\\ 1&0&0&0\\ 0&1&0&0\\ 0&0&0&1 \end{pmatrix} \quad \begin{pmatrix} 1&1&0&0\\ 1&0&0&0\\ 0&1&0&0\\ 0&0&1&0 \end{pmatrix}.$$

Proof. See [8] □

Lemma 7.2.6. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (3,4,1,8), (3,4,8,1). We have

$$\begin{aligned}(3,4,1,8) &= Sq^1(3,1,1,10) + Sq^2\big((5,2,1,6) + (5,1,2,6)\\ &\quad + (2,1,1,10)\big) + Sq^4\big((3,2,1,6) + (3,1,2,6)\big)\\ &\quad + (2,1,1,12) + (3,1,4,8) \mod \mathcal{L}_4(2;1;1;1),\\ (3,4,8,1) &= Sq^1(3,1,10,1) + Sq^2\big((5,2,6,1) + (5,1,6,2)\\ &\quad + (2,1,10,1)\big) + Sq^4\big((3,2,6,1) + (3,1,6,2)\big)\\ &\quad + (2,1,12,1) + (3,1,8,4) \mod \mathcal{L}_4(2;1;1;1).\end{aligned}$$

The lemma follows. □

Using the results in Section 4, Section 5, Lemmas 7.1.3-7.1.1, 7.2.4, 7.2.5, 7.2.6 and Theorem 2.5, we get Proposition 7.2.3.

Now, we show that the elements listed in Theorem 7.2.2 are linearly independent.

The case $t = 1$ is trivial. We begin with the case $t = 2$.

Proposition 7.2.7. *The elements* $[c_{2,1,i}], 1 \leqslant i \leqslant 7$ *and* $[\phi(b_{1,j})], 1 \leqslant j \leqslant 6$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_8$.

Proof. Suppose that there is a linear relation

$$\sum_{1\leqslant i\leqslant 7} \gamma_i[c_{2,1,i}] + \sum_{1\leqslant j\leqslant 6} \eta_j[\phi(b_{1,j})] = 0, \tag{7.2.7.1}$$

with $\gamma_i, \eta_j \in \mathbb{F}_2$.

Apply the squaring operation Sq^0_* to (7.2.7.1) and we get

$$\sum_{1\leqslant j\leqslant 6} \eta_j[b_{1,j}] = 0.$$

Since $\{[b_{1,j}]; 1 \leqslant j \leqslant 6\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_2$, we obtain $\eta_j = 0, 1 \leqslant j \leqslant 6$. Hence, (7.2.7.1) becomes

$$\sum_{1\leqslant i\leqslant 7} \gamma_i[c_{2,1,i}] = 0. \tag{7.2.7.2}$$

Now, we prove $\gamma_i = 0, 1 \leqslant i \leqslant 7$.

Applying the homomorphisms f_1, f_2, f_4, f_6 to the relation (7.2.7.2), we obtain

$$\begin{aligned}
&\gamma_4[1,1,6] + \gamma_5[1,6,1] + \gamma_4[1,2,5]\\
&\quad + \gamma_5[1,3,4] + \gamma_{\{2,4\}}[3,1,4] + \gamma_{\{3,5\}}[3,4,1] = 0,\\
&\gamma_4[1,1,6] + \gamma_6[1,6,1] + \gamma_4[1,2,5] + \gamma_6[1,3,4]\\
&\quad + \gamma_{\{1,4,7\}}[3,1,4] + \gamma_{\{3,6\}}[3,4,1] = 0,\\
&\gamma_{\{3,7\}}[1,6,1] + \gamma_4[1,2,5] + \gamma_{\{1,2,5,6,7\}}[1,3,4] + \gamma_7[3,4,1] = 0,\\
&\gamma_1[1,1,6] + \gamma_{\{2,3,4,5\}}[1,2,5] + \gamma_6[1,3,4] + \gamma_7[3,1,4] = 0.
\end{aligned}$$

From these equalities, we get $\gamma_i = 0,\ 1 \leqslant i \leqslant 7$. The proposition is proved. □

Proposition 7.2.8. *The elements* $[c_{3,1,i}], 1 \leqslant i \leqslant 11$, $[\phi(b_{2,j})], 1 \leqslant j \leqslant 24$, *and* $[\phi^2(a_{1,\ell})], 1 \leqslant \ell \leqslant 4$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{16}$.

Proof. Observe that

$$Sq^0_*([c_{3,1,i}]) = 0, Sq^0_*([\phi(b_{2,j})]) = [b_{2,j}], Sq^0_*([\phi^2(a_{1,\ell})]) = [\phi(a_{1,\ell})],$$

and $\{[b_{2,j}], [\phi(a_{1,\ell})]; 1 \leqslant j \leqslant 20, 1 \leqslant \ell \leqslant 4\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_6$. Hence, it suffices to show that the elements $[c_{3,1,i}], 1 \leqslant i \leqslant 11$, are linearly independent.

Suppose that there is a linear relation

$$\sum_{1 \leqslant i \leqslant 11} \gamma_i [c_{3,1,i}] = 0, \tag{7.2.8.1}$$

with $\gamma_i \in \mathbb{F}_2, 1 \leqslant i \leqslant 11$. Applying the homomorphisms f_1, f_2, f_3, f_4 to the relation (7.2.8.1), we obtain

$$\begin{aligned}
&\gamma_{\{3,4,5\}}[1,1,14] + \gamma_3[1,14,1] + \gamma_4[1,2,13]\\
&\quad + \gamma_5[1,3,12] + \gamma_{\{2,3,4,5\}}[3,1,12] = 0,\\
&\gamma_{\{3,4,6,8,10,11\}}[1,1,14] + \gamma_3[1,14,1] + \gamma_4[1,2,13]\\
&\quad + \gamma_{\{6,8,10,11\}}[1,3,12] + \gamma_{\{1,3,4,6,7,8,9,10,11\}}[3,1,12] = 0,\\
&\gamma_{\{2,5,6,8,10,11\}}[1,1,14] + \gamma_2[1,14,1] + \gamma_5[1,2,13]\\
&\quad + \gamma_{\{6,8,10,11\}}[1,3,12] + \gamma_{\{1,2,5,6,7,8,9,10,11\}}[3,1,12] = 0,\\
&\gamma_{\{7,9,10,11\}}[1,1,14] + \gamma_3[1,14,1] + \gamma_4[1,2,13]\\
&\quad + \gamma_{\{1,2,5,6,7,8,9,10,11\}}[1,3,12] + \gamma_{\{7,9,10,11\}}[3,1,12] = 0.
\end{aligned}$$

Computing directly from the above equalities gives

$$\begin{cases}
\gamma_2 = \gamma_3 = \gamma_4 = \gamma_5 = 0,\\
\gamma_{\{1,10,11\}} = \gamma_{\{1,7,9\}} = \gamma_{\{1,6,8\}} = 0.
\end{cases} \tag{7.2.8.2}$$

Substituting (7.2.8.2) into the relation (7.2.8.1), we get

$$\gamma_1[\theta_1] + \gamma_6[\theta_2] + \gamma_7[\theta_3] + \gamma_{11}[\theta_4] = 0, \tag{7.2.8.3}$$

where $\theta_1 = b_{2,1,1} + b_{2,1,8} + b_{2,1,9} + b_{2,1,10}, \theta_2 = b_{2,1,6} + b_{2,1,8}, \theta_3 = b_{2,1,7} + b_{2,1,9}, \theta_4 = b_{2,1,10} + b_{2,1,11}$.

We prove $\gamma_1 = \gamma_6 = \gamma_7 = \gamma_{11} = 0$. The proof is divided into 4 steps.

Step 1. The homomorphism φ_3 sends (7.2.8.3) to

$$\gamma_1([\theta_1] + [\theta_4]) + \gamma_6[\theta_2] + \gamma_7[\theta_3] + \gamma_{11}[\theta_4] = 0. \tag{7.2.8.4}$$

Combining (7.2.8.3) and (7.2.8.4), we obtain

$$\gamma_1[\theta_4] = 0. \tag{7.2.8.5}$$

Now we prove $[\theta_4] \neq 0$. Suppose the contrary, that $[\theta_4] = 0$. Then, the polynomial θ_4 is hit. Hence, we have

$$\theta_4 = Sq^1(A) + Sq^2(B) + Sq^4(C) + Sq^8(D),$$

for some polynomials $A \in (R_4)_{15}, B \in (R_4)_{14}, C \in (R_4)_{12}, D \in (R_4)_8$. Let $(Sq^2)^3$ act on the both sides of this equality. Since $Sq^8(D) = D^2$, $(Sq^2)^3Sq^1 = 0$ and $(Sq^2)^3Sq^2 = 0$, we obtain

$$(Sq^2)^3(\theta_4) = (Sq^2)^3Sq^4(C).$$

On the other hand, by a direct calculation, we see that the monomial (6,6,6,4) is a term of $(Sq^2)^3(\theta_4)$. This monomial is not a term of $(Sq^2)^3Sq^4(y)$ for all monomial $y \in (R_4)_{12}$. So, $(Sq^2)^3(\theta_4) \neq (Sq^2)^3Sq^4(C)$, for all $C \in (R_4)_{12}$ and we have a contradiction. Hence, $[\theta_4] \neq 0$ and $\gamma_1 = 0$.

Step 2. Since $\gamma_1 = 0$, the relation (7.2.8.3) becomes

$$\gamma_6[\theta_2] + \gamma_7[\theta_3] + \gamma_{11}[\theta_4] = 0. \tag{7.2.8.6}$$

The homomorphism φ_2 sends (7.2.8.6) to

$$(\gamma_7 + \gamma_{11})[\theta_1] + (\gamma_6 + \gamma_7 + \gamma_{11})[\theta_2] + \gamma_7[\theta_3] + \gamma_{11}[\theta_4] = 0. \tag{7.2.8.7}$$

Using the relation (7.2.8.7) and by an analogous argument as given in Step 1, we get

$$\gamma_7 = \gamma_{11}.$$

Step 3. The homomorphism φ_1 sends (7.2.8.6) to

$$\gamma_7[\theta_2] + \gamma_6[\theta_3] + \gamma_{11}[\theta_4] = 0. \tag{7.2.8.8}$$

Using the relation (7.2.8.8) and by a same argument as given in Step 2, we obtain

$$\gamma_6 = \gamma_{11}.$$

Step 4. The homomorphism φ_4 sends (7.2.8.6) to

$$\gamma_6[\theta_2] + \gamma_7([\theta_2] + [\theta_3]) + \gamma_{11}[\theta_4] = 0. \tag{7.2.8.9}$$

Combining (7.2.8.8) and (7.2.8.9), we obtain

$$\gamma_7[\theta_2] = 0. \tag{7.2.8.10}$$

Using the relation (7.2.8.10) and by an analogous argument as given in Step 3, we get $\gamma_7 = 0$. Hence, the proposition is proved. □

Proposition 7.2.9. *For $t \geqslant 4$, the elements $[c_{t,1,i}], 1 \leqslant i \leqslant 7$, and $[\phi(b_{t-1,j})], 1 \leqslant j \leqslant \mu_6(t-1)$, and $[\phi^2(a_{t-2,\ell})], 1 \leqslant \ell \leqslant \mu_1(t-2)$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+1}}$.*

Proof. By an analogous argument as given in the proof of Proposition 7.2.8, it suffices to show that the elements $[c_{t,1,j}], 1 \leqslant j \leqslant 7$, are linearly independent. Suppose that there is a linear relation

$$\sum_{i=1}^{7} \gamma_i [c_{t,1,i}] = 0,$$

with $\gamma_i \in \mathbb{F}_2$ for all i.

Applying the homomorphisms f_1, f_2, f_4 to this relation, we obtain

$$\begin{aligned}
&\gamma_{\{3,4,5\}} v_{t,1,1} + \gamma_3 v_{t,1,2} + \gamma_4 v_{t,1,12} + \gamma_5 v_{t,1,13} + \gamma_{\{2,3,4,5\}} v_{t,1,14} = 0,\\
&\gamma_{\{3,4,6\}} v_{t,1,1} + \gamma_3 v_{t,1,2} + \gamma_4 v_{t,1,12} + \gamma_6 v_{t,1,13} + \gamma_{\{1,3,4,6,7\}} v_{t,1,14} = 0,\\
&\gamma_7 v_{t,1,1} + \gamma_3 v_{t,1,2} + \gamma_4 v_{t,1,12} + \gamma_{\{1,2,5,6,7\}} v_{t,1,13} + \gamma_7 v_{t,1,14} = 0.
\end{aligned}$$

From these equalities, we obtain $\gamma_i = 0$ for $i = 1, 2, \ldots, 7$. The proposition follows. □

Remark 7.2.10. The $\mathbb{F}_2$-subspace of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{16}$ generated by $[\theta_1], [\theta_2]$, $[\theta_3]$, $[\theta_4]$, which are defined as in the proof of Proposition 7.2.8, is an $GL_4(\mathbb{F}_2)$-submodule of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{16}$.

7.3. The case $s \geqslant 2$ and $t = 1$.

According to Kameko [8], for $s \geqslant 2, \dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+1}+2^s-2} = 14$ with a basis given by the following classes:

$$\begin{aligned}
&v_{1,s,1} = [1, 2^s - 2, 2^{s+1} - 1], && v_{1,s,2} = [1, 2^{s+1} - 1, 2^s - 2],\\
&v_{1,s,3} = [2^{s+1} - 1, 1, 2^s - 2], && v_{1,s,4} = [1, 2^s - 1, 2^{s+1} - 2],\\
&v_{1,s,5} = [1, 2^{s+1} - 2, 2^s - 1], && v_{1,s,6} = [2^s - 1, 1, 2^{s+1} - 2],\\
&v_{1,s,7} = [0, 2^s - 1, 2^{s+1} - 1], && v_{1,s,8} = [0, 2^{s+1} - 1, 2^s - 1],\\
&v_{1,s,9} = [2^s - 1, 0, 2^{s+1} - 1], && v_{1,s,10} = [2^s - 1, 2^{s+1} - 1, 0],\\
&v_{1,s,11} = [2^{s+1} - 1, 0, 2^s - 1], && v_{1,s,12} = [2^{s+1} - 1, 2^s - 1, 0],\\
&v_{1,s,13} = [3, 2^{s+1} - 3, 2^s - 2].
\end{aligned}$$

For $s = 2$, $v_{1,2,14} = [3, 3, 4]$.
For $s \geqslant 3$, $v_{1,s,14} = [3, 2^s - 3, 2^{s+1} - 2]$.

So, we easily obtain

Proposition 7.3.1. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+1}+2^s-2}$ is an $\mathbb{F}_2$-vector space of dimension 44 with a basis consisting of all the classes represented by the following monomials:*

For $s \geqslant 2$,

$b_{1,s,1} = (0, 1, 2^s - 2, 2^{s+1} - 1)$, $b_{1,s,2} = (0, 1, 2^{s+1} - 1, 2^s - 2)$,
$b_{1,s,3} = (0, 2^{s+1} - 1, 1, 2^s - 2)$, $b_{1,s,4} = (1, 0, 2^s - 2, 2^{s+1} - 1)$,
$b_{1,s,5} = (1, 0, 2^{s+1} - 1, 2^s - 2)$, $b_{1,s,6} = (1, 2^s - 2, 0, 2^{s+1} - 1)$,
$b_{1,s,7} = (1, 2^s - 2, 2^{s+1} - 1, 0)$, $b_{1,s,8} = (1, 2^{s+1} - 1, 0, 2^s - 2)$,
$b_{1,s,9} = (1, 2^{s+1} - 1, 2^s - 2, 0)$, $b_{1,s,10} = (2^{s+1} - 1, 0, 1, 2^s - 2)$,
$b_{1,s,11} = (2^{s+1} - 1, 1, 0, 2^s - 2)$, $b_{1,s,12} = (2^{s+1} - 1, 1, 2^s - 2, 0)$,
$b_{1,s,13} = (0, 1, 2^s - 1, 2^{s+1} - 2)$, $b_{1,s,14} = (0, 1, 2^{s+1} - 2, 2^s - 1)$,
$b_{1,s,15} = (0, 2^s - 1, 1, 2^{s+1} - 2)$, $b_{1,s,16} = (1, 0, 2^s - 1, 2^{s+1} - 2)$,
$b_{1,s,17} = (1, 0, 2^{s+1} - 2, 2^s - 1)$, $b_{1,s,18} = (1, 2^s - 1, 0, 2^{s+1} - 2)$,
$b_{1,s,19} = (1, 2^s - 1, 2^{s+1} - 2, 0)$, $b_{1,s,20} = (1, 2^{s+1} - 2, 0, 2^s - 1)$,
$b_{1,s,21} = (1, 2^{s+1} - 2, 2^s - 1, 0)$, $b_{1,s,22} = (2^s - 1, 0, 1, 2^{s+1} - 2)$,
$b_{1,s,23} = (2^s - 1, 1, 0, 2^{s+1} - 2)$, $b_{1,s,24} = (2^s - 1, 1, 2^{s+1} - 2, 0)$,
$b_{1,s,25} = (0, 0, 2^s - 1, 2^{s+1} - 1)$, $b_{1,s,26} = (0, 0, 2^{s+1} - 1, 2^s - 1)$,
$b_{1,s,27} = (0, 2^s - 1, 0, 2^{s+1} - 1)$, $b_{1,s,28} = (0, 2^s - 1, 2^{s+1} - 1, 0)$,
$b_{1,s,29} = (0, 2^{s+1} - 1, 0, 2^s - 1)$, $b_{1,s,30} = (0, 2^{s+1} - 1, 2^s - 1, 0)$,
$b_{1,s,31} = (2^s - 1, 0, 0, 2^{s+1} - 1)$, $b_{1,s,32} = (2^s - 1, 0, 2^{s+1} - 1, 0)$,
$b_{1,s,33} = (2^s - 1, 2^{s+1} - 1, 0, 0)$, $b_{1,s,34} = (2^{s+1} - 1, 0, 0, 2^s - 1)$,
$b_{1,s,35} = (2^{s+1} - 1, 0, 2^s - 1, 0)$, $b_{1,s,36} = (2^{s+1} - 1, 2^s - 1, 0, 0)$,
$b_{1,s,37} = (0, 3, 2^{s+1} - 3, 2^s - 2)$, $b_{1,s,38} = (3, 0, 2^{s+1} - 3, 2^s - 2)$,
$b_{1,s,39} = (3, 2^{s+1} - 3, 0, 2^s - 2)$, $b_{1,s,40} = (3, 2^{s+1} - 3, 2^s - 2, 0)$,

For $s = 2$,

$b_{1,2,41} = (0, 3, 3, 4)$, $b_{1,2,42} = (3, 0, 3, 4)$,
$b_{1,2,43} = (3, 3, 0, 4)$, $b_{1,2,44} = (3, 3, 4, 0)$.

For $s \geqslant 3$,

$b_{1,s,41} = (0, 3, 2^s - 3, 2^{s+1} - 2)$, $b_{1,s,42} = (3, 0, 2^s - 3, 2^{s+1} - 2)$,
$b_{1,s,43} = (3, 2^s - 3, 0, 2^{s+1} - 2)$, $b_{1,s,44} = (3, 2^s - 3, 2^{s+1} - 2, 0)$.

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}+2^s-2}$. We have

Theorem 7.3.2. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}+2^s-2}$ is an $\mathbb{F}_2$-vector space with a basis consisting of all the classes represented by the following monomials:*

1. $\phi(c_{1,j}), 1 \leqslant j \leqslant 14$ for $s = 2$.

2. $\phi(a_{1,s-2,j}), 1 \leqslant j \leqslant \mu_2(s-2)$ for $s \geqslant 3$. Here $c_{1,j}$, $a_{1,s-2,j}$, $\mu_2(s-2)$ are defined as in Section 5 and Section 6, the homomorphism ϕ is defined as in Definition 2.9.

3. For $s \geqslant 2$,

$b_{1,s,45} = (1, 1, 2^s - 2, 2^{s+1} - 2)$, $b_{1,s,46} = (1, 1, 2^{s+1} - 2, 2^s - 2)$,
$b_{1,s,47} = (1, 2^s - 2, 1, 2^{s+1} - 2)$, $b_{1,s,48} = (1, 2^{s+1} - 2, 1, 2^s - 2)$,
$b_{1,s,49} = (1, 2, 2^{s+1} - 3, 2^s - 2)$, $b_{1,s,50} = (1, 2, 2^s - 1, 2^{s+1} - 4)$,
$b_{1,s,51} = (1, 2, 2^{s+1} - 4, 2^s - 1)$, $b_{1,s,52} = (1, 2^s - 1, 2, 2^{s+1} - 4)$,
$b_{1,s,53} = (2^s - 1, 1, 2, 2^{s+1} - 4)$, $b_{1,s,54} = (1, 3, 2^{s+1} - 4, 2^s - 2)$,
$b_{1,s,55} = (3, 1, 2^{s+1} - 4, 2^s - 2)$.

For $s = 2$, $b_{1,2,56} = (3, 4, 1, 2)$.

For $s \geqslant 3$,

$$\begin{array}{ll} b_{1,s,56} = (1, 3, 2^s - 2, 2^{s+1} - 4), & b_{1,s,57} = (3, 1, 2^s - 2, 2^{s+1} - 4),\\ b_{1,s,58} = (1, 2, 2^s - 4, 2^{s+1} - 1), & b_{1,s,59} = (1, 2, 2^{s+1} - 1, 2^s - 4),\\ b_{1,s,60} = (1, 2^{s+1} - 1, 2, 2^s - 4), & b_{1,s,61} = (2^{s+1} - 1, 1, 2, 2^s - 4),\\ b_{1,s,62} = (1, 2, 2^s - 3, 2^{s+1} - 2), & b_{1,s,63} = (1, 3, 2^s - 4, 2^{s+1} - 2),\\ b_{1,s,64} = (1, 3, 2^{s+1} - 2, 2^s - 4), & b_{1,s,65} = (3, 1, 2^s - 4, 2^{s+1} - 2),\\ b_{1,s,66} = (3, 1, 2^{s+1} - 2, 2^s - 4), & b_{1,s,67} = (3, 2^{s+1} - 3, 2, 2^s - 4),\\ b_{1,s,68} = (3, 2^s - 3, 2, 2^{s+1} - 4), & b_{1,s,69} = (3, 5, 2^{s+1} - 6, 2^s - 4). \end{array}$$

For $s = 3$, $b_{1,3,70} = (3, 5, 6, 8)$.
For $s \geqslant 4$, $b_{1,s,70} = (3, 5, 2^s - 6, 2^{s+1} - 4)$.

To prove this theorem, we prove the following propositions.

Proposition 7.3.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+1}+2^s-2}$ is generated by the elements listed in Theorem 7.3.2.*

The proof of the proposition is based on some lemmas.

Lemma 7.3.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 0 & 1 & 0 & 1\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 1\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 0\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

Proof. See [8]. □

Lemma 7.3.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 0 & 1 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 0\\ 1 & 0 & 1 & 0\\ 1 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Proof. See [8]. □

Lemma 7.3.6. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 1 & 0 & 0\\ 0 & 1 & 1 & 0\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 1\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 0\\ 0 & 1 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 1 & 0\\ 1 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 1\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 1\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 1 & 0\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Proof. The monomials corresponding to these matrices respectively are (1,7,10,4), (7,1,10,4), (1,6,7,8), (1,7,6,8), (7,1,6,8), (3,3,4,12), (3,3,12,4), (7,9,2,4), (7,8,3,4), (3,5,8,6). We prove the lemma for the matrices corresponding to the monomials (7,1,10,4), (1,6,7,8), (3,3,4,12), (7,9,2,4), (7,8,3,4), (3,5,8,6). The others can be obtained by a same computation. We have

$$\begin{aligned}
(7,1,10,4) &= Sq^1\big((7,1,9,4)+(7,4,7,3)+(7,1,8,5)\big)\\
&\quad + Sq^2\big((7,2,7,4)+(7,2,8,3)\big)+Sq^4\big((5,2,7,4)\\
&\quad + (4,4,7,3)\big)+(5,2,11,4)+(5,2,7,8)+(4,4,11,3)\\
&\quad + (4,8,7,3)+(7,1,8,6) \bmod \mathcal{L}_4(2;2;2;1),\\
(1,7,6,8) &= Sq^1\big((1,7,5,8)+(4,7,3,7)+(1,8,5,7)+(1,9,4,7)\\
&\quad + (1,7,4,9)\big)+Sq^2\big((2,7,3,8)+(2,8,3,7)+(2,7,4,7)\big)\\
&\quad + Sq^4\big((1,6,4,7)+(1,4,6,7)\big)+(1,4,10,7)+(1,4,6,11)\\
&\quad + (1,6,8,7)+(1,6,4,11)+(1,7,4,10) \bmod \mathcal{L}_4(2;2;2;1),\\
(3,3,4,12) &= Sq^1\big((3,3,4,11)+(3,8,3,7)+(3,3,8,7)\big)+Sq^2\big((2,3,4,11)\\
&\quad + (2,8,3,7)+(2,3,8,7)\big)+Sq^4(3,4,4,7)+(2,5,4,11)\\
&\quad + (2,3,4,13)+(2,8,5,7)+(2,5,8,7) \quad \bmod \mathcal{L}_4(2;2;2;1),\\
(7,9,2,4) &= Sq^1(7,7,3,4)+Sq^2(7,7,2,4)+Sq^4\big((5,7,2,4)\\
&\quad + (4,7,3,4)\big)+(5,11,2,4)+(5,7,2,8)+(4,11,3,4)\\
&\quad + (4,7,3,8)+(7,8,3,4) \quad \bmod \mathcal{L}_4(2;2;2;1),\\
(7,8,3,4) &= Sq^1(7,8,3,3)+Sq^2(7,8,2,3)\\
&\quad + (7,8,2,5) \quad \bmod \mathcal{L}_4(2;2;2;1),\\
(3,5,8,6) &= Sq^1\big((3,3,9,6)+(3,3,6,9)\big)+Sq^2\big((5,3,6,6)+(2,3,6,9)\\
&\quad + (2,3,9,6)\big)+Sq^4(3,3,6,6)+(3,5,6,8)+(3,4,6,9)\\
&\quad + (3,4,9,6)+(2,5,6,9)+(2,5,9,6) \bmod \mathcal{L}_4(2;2;2;1).
\end{aligned}$$

The lemma is proved. □

Using the results in Section 4, Section 5, Lemma 7.1.5, 7.1.6, 7.1.1, 7.2.4, 7.2.5, 7.2.6, 7.3.4, 7.3.5 and Theorem 2.5, we get Proposition 7.3.3.

Now, we show that the elements listed in Theorem 7.3.2 are linearly independent.

Proposition 7.3.7. *The elements* $[b_{1,2,i}], 45 \leqslant i \leqslant 56$, *and* $[\phi(c_{1,j})], 1 \leqslant j \leqslant 14$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{10}$.

Proof. Suppose that there is a linear relation

$$\sum_{i=45}^{56} \gamma_i [b_{1,2,i}] + \sum_{j=1}^{14} \eta_j [\phi(c_{1,j})] = 0, \tag{7.3.7.1}$$

with $\gamma_i, \eta_j \in \mathbb{F}_2$.

Applying the squaring operation Sq^0_* to (7.3.7.1), we obtain

$$\sum_{j=1}^{14} \eta_j [c_{1,j}] = 0.$$

Since $\{[c_{1,j}]; 1 \leqslant j \leqslant 14\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_3$, we obtain $\eta_j = 0, 1 \leqslant j \leqslant 14$. Hence, (7.3.7.1) becomes

$$\sum_{45 \leqslant i \leqslant 56} \gamma_i [b_{1,2,i}] = 0. \tag{7.3.7.2}$$

Now, we prove $\gamma_i = 0, 45 \leqslant i \leqslant 56$.

Apply the homomorphisms f_1, f_2, f_3 to the relation (7.3.7.2) and we get

$$\begin{aligned}
&\gamma_{\{48,56\}}[7,1,2] + \gamma_{51}[1,3,6] + \gamma_{51}[1,6,3] \\
&\quad + \gamma_{47}[3,1,6] + \gamma_{49}[3,5,2] + \gamma_{\{50,51\}}[3,3,4] = 0, \\
&\gamma_{\{46,55\}}[7,1,2] + \gamma_{51}[1,3,6] + \gamma_{51}[1,6,3] \\
&\quad + \gamma_{\{45,51\}}[3,1,6] + \gamma_{54}[3,5,2] + \gamma_{\{51,52,54\}}[3,3,4] = 0, \\
&\gamma_{\{45,47,53\}}[7,1,2] + \gamma_{50}[1,3,6] + \gamma_{\{46,49,50\}}[3,1,6] \\
&\quad + \gamma_{50}[1,6,3] + \gamma_{\{48,52\}}[3,5,2] + \gamma_{\{50,52,54\}}[3,3,4] = 0.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
\gamma_{47} = \gamma_{49} = \gamma_{50} = \gamma_{51} = \gamma_{54} = 0, \\
\gamma_{\{48,56\}} = \gamma_{\{50,51\}} = \gamma_{\{46,55\}} = \gamma_{\{45,51\}} = \gamma_{\{51,52,54\}} = 0, \\
\gamma_{\{45,47,53\}} = \gamma_{\{46,49,50\}} = \gamma_{\{48,52\}} = \gamma_{\{50,52,54\}} = 0.
\end{cases} \tag{7.3.7.3}$$

The relation (7.3.7.3) implies $\gamma_j = 0$ for $j = 45, \ldots, 56$. So, the proposition is proved. □

Proposition 7.3.8. *The elements* $[b_{1,3,i}], 45 \leqslant i \leqslant 70$, *and* $[\phi(a_{1,1,j})], 1 \leqslant j \leqslant 46$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{22}$.

Proof. Suppose that there is a linear relation

$$\sum_{i=45}^{70} \gamma_i [b_{1,3,i}] + \sum_{j=1}^{46} \eta_j [\phi(a_{1,1,j})] = 0, \tag{7.3.8.1}$$

with $\gamma_i, \eta_j \in \mathbb{F}_2$.

Applying the squaring operation Sq^0_* to (7.3.8.1), we get

$$\sum_{j=1}^{46} \eta_j [a_{1,1,j}] = 0.$$

Since $\{[a_{1,1,j}]; 1 \leqslant j \leqslant 46\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_9$, we obtain $\eta_j = 0, 1 \leqslant j \leqslant 46$. Hence, (7.3.8.1) becomes

$$\sum_{45 \leqslant i \leqslant 70} \gamma_i [b_{1,3,i}] = 0. \tag{7.3.8.2}$$

Now, we prove $\gamma_i = 0, 45 \leqslant i \leqslant 70$.

Applying the homomorphisms $f_t, t = 1, \ldots, 6$, to the relation (7.3.8.2), we obtain

$$\begin{aligned}
&\gamma_{58}[1,6,15] + \gamma_{59}[1,15,6] + \gamma_{48}[15,1,6] + \gamma_{50}[1,7,14]\\
&\quad + \gamma_{51}[1,14,7] + \gamma_{47}[7,1,14] + \gamma_{62}[3,5,14] + \gamma_{49}[3,13,6] = 0,\\
&\gamma_{58}[1,6,15] + \gamma_{60}[1,15,6] + \gamma_{\{46,55,64\}}[15,1,6] + \gamma_{\{45,56,65\}}[7,1,14]\\
&\quad + \gamma_{52}[1,7,14] + \gamma_{51}[1,14,7] + \gamma_{\{51,54,63\}}[3,5,14] + \gamma_{54}[3,13,6] = 0,\\
&\gamma_{59}[1,6,15] + \gamma_{60}[1,15,6] + \gamma_{\{45,47,62,57,63,68\}}[15,1,6]\\
&\quad + \gamma_{52}[1,7,14] + \gamma_{50}[1,14,7] + \gamma_{\{46,48,4954,66,67,69\}}[7,1,14]\\
&\quad + \gamma_{\{48,5056,64,67,70\}}[3,5,14] + \gamma_{\{48,52,56,67,70\}}[3,13,6] = 0,\\
&\gamma_{58}[1,6,15] + \gamma_{\{46,48,49,54,66,67,69\}}[1,15,6] + \gamma_{61}[15,1,6] + \gamma_{51}[1,14,7]\\
&\quad + \gamma_{\{45,47,62,57,63,68\}}[1,7,14] + \gamma_{53}[7,1,14] + \gamma_{65}[3,5,14] + \gamma_{55}[3,13,6] = 0,\\
&\gamma_{59}[1,6,15] + \gamma_{\{45,56,65\}}[1,15,6] + \gamma_{61}[15,1,6] + \gamma_{\{46,55,64\}}[1,7,14]\\
&\quad + \gamma_{50}[1,14,7] + \gamma_{53}[7,1,14] + \gamma_{\{53,66\}}[3,5,14] + \gamma_{\{53,57,70\}}[3,13,6] = 0,\\
&\gamma_{47}[1,6,15] + \gamma_{60}[1,15,6] + \gamma_{61}[15,1,6] + \gamma_{52}[1,7,14]\\
&\quad + \gamma_{48}[1,14,7] + \gamma_{53}[7,1,14] + \gamma_{\{68,69,70\}}[3,5,14] + \gamma_{67}[3,13,6] = 0.
\end{aligned}$$

From the above equalities, we get

$$\begin{cases}
\gamma_j = 0, \ j = 47, \ldots, 63, 65, 66, 54, 55, 67,\\
\gamma_{\{46,64\}} = \gamma_{\{45,56\}} = \gamma_{\{45,57,68\}} = \gamma_{\{46,69\}} = 0,\\
\gamma_{\{56,64,70\}} = \gamma_{\{56,70\}} = \gamma_{\{57,70\}} = \gamma_{\{68,69,70\}} = 0.
\end{cases} \tag{7.3.8.3}$$

Computing directly from (7.3.8.3), we obtain $\gamma_i = 0$ for any i. The proposition is proved. □

Proposition 7.3.9. *For $s \geqslant 4$, the elements $[b_{1,s,i}], 45 \leqslant i \leqslant 70$, and $[\phi(a_{1,s-2,j})], 1 \leqslant j \leqslant \mu_2(s-2)$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{3.2^s-2}$.*

Proof. Suppose that there is a linear relation

$$\sum_{i=45}^{70} \gamma_i [b_{1,s-2,i}] + \sum_{j=1}^{\mu_2(s-2)} \eta_j [\phi(a_{1,s-2,j})] = 0, \tag{7.3.9.1}$$

with $\gamma_i, \eta_j \in \mathbb{F}_2$.

Applying the squaring operation Sq^0_* to (7.3.9.1), we get

$$\sum_{j=1}^{\mu_2(s-2)} \eta_j[a_{1,s-2,j}] = 0.$$

Since $\{[a_{1,s-2,j}]; 1 \leqslant j \leqslant \mu_2(s-2)\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{2^s+2^{s-1}-3}$, we obtain $\eta_j = 0, 1 \leqslant j \leqslant \mu_2(s-2)$. Hence, (7.3.9.1) becomes

$$\sum_{45\leqslant i\leqslant 70} \gamma_i[b_{1,s,i}] = 0. \tag{7.3.9.2}$$

We need to prove $\gamma_i = 0, 45 \leqslant i \leqslant 70$.

Apply the homomorphisms $f_t, t = 1, \ldots, 6$, to the relation (7.3.9.2) and we obtain

$$\begin{aligned}
&\gamma_{58}v_{1,s,1} + \gamma_{59}v_{1,s,2} + \gamma_{48}v_{1,s,3} + \gamma_{50}v_{1,s,4}\\
&\quad + \gamma_{51}v_{1,s,5} + \gamma_{47}v_{1,s,6} + \gamma_{49}v_{1,s,13} + \gamma_{62}v_{1,s,14} = 0,\\
&\gamma_{58}v_{1,s,1} + \gamma_{60}v_{1,s,2} + \gamma_{\{46,64,66\}}v_{1,s,3} + \gamma_{52}v_{1,s,4} + \gamma_{51}v_{1,s,5}\\
&\quad + \gamma_{\{45,56,65\}}v_{1,s,6} + \gamma_{54}v_{1,s,13} + \gamma_{\{51,54,63\}}v_{1,s,14} = 0,\\
&\gamma_{59}v_{1,s,1,60}v_{1,s,2} + \gamma_{\{45,47,57,62,63,68,70\}}v_{1,s,3} + \gamma_{52}v_{1,s,4} + \gamma_{50}v_{1,s,5}\\
&\quad + a_2v_{1,s,6} + \gamma_{\{52,56\}}v_{1,s,13} + \gamma_{\{50,56,64\}}v_{1,s,14} = 0,\\
&\gamma_{58}v_{1,s,1} + \gamma_{\{46,48,49,54,66,67,69\}}v_{1,s,2} + \gamma_{61}v_{1,s,3} + a_1v_{1,s,4}\\
&\quad + \gamma_{51}v_{1,s,5} + \gamma_{55}v_{1,s,13} + \gamma_{53}v_{1,s,6,65}v_{1,s,14} = 0,\\
&\gamma_{59}v_{1,s,1} + \gamma_{\{45,56,65\}}v_{1,s,2} + \gamma_{46,55,64}v_{1,s,4} + \gamma_{61}v_{1,s,3}\\
&\quad + \gamma_{50}v_{1,s,5} + \gamma_{53}v_{1,s,6} + \gamma_{57}v_{1,s,13} + \gamma_{66}v_{1,s,14} = 0,\\
&\gamma_{47}v_{1,s,1} + \gamma_{60}v_{1,s,2} + \gamma_{61}v_{1,s,3} + \gamma_{52}v_{1,s,4}\\
&\quad + \gamma_{48}v_{1,s,5} + \gamma_{53}v_{1,s,6} + \gamma_{67}v_{1,s,13} + \gamma_{68}v_{1,s,14} = 0.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
\gamma_j = 0,\ j = 47, \ldots, 62, 65, 66, 54, \ldots, 68,\\
\gamma_{\{46,64,55\}} = \gamma_{\{45,65,56\}} = \gamma_{\{51,63,54\}} = 0,\\
\gamma_{\{50,64,56\}} = \gamma_{\{52,56\}} = 0.
\end{cases} \tag{7.3.9.3}$$

Computing directly from (7.3.9.3), we obtain $\gamma_i = 0$ for any i. The proposition is proved. □

7.4. **The case $s \geqslant 2$ and $t \geqslant 2$.**

Using the result in Kameko [8], we see that for $t \geqslant 2$ and $s \geqslant 2$, $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{t+s}+2^s-2} = 21$ with a basis given by the following classes:

$v_{t,s,1} = [1, 2^s - 2, 2^{s+t} - 1]$, $v_{t,s,2} = [1, 2^{s+t} - 1, 2^s - 2]$,
$v_{t,s,3} = [2^{s+t} - 1, 1, 2^s - 2]$, $v_{t,s,4} = [1, 2^s - 1, 2^{s+t} - 2]$,
$v_{t,s,5} = [1, 2^{s+t} - 2, 2^s - 1]$, $v_{t,s,6} = [2^s - 1, 1, 2^{s+t} - 2]$,
$v_{t,s,7} = [1, 2^{s+1} - 2, 2^{s+t} - 2^s - 1]$, $v_{t,s,8} = [1, 2^{s+1} - 1, 2^{s+t} - 2^s - 2]$,
$v_{t,s,9} = [2^{s+1} - 1, 1, 2^{s+t} - 2^s - 2]$, $v_{t,s,10} = [3, 2^{s+1} - 3, 2^{s+t} - 2^s - 2]$,
$v_{t,s,11} = [3, 2^{s+t} - 3, 2^s - 2]$, $v_{t,s,12} = [0, 2^s - 1, 2^{s+t} - 1]$,
$v_{t,s,13} = [0, 2^{s+1} - 1, 2^{s+t} - 2^s - 1]$, $v_{t,s,14} = [0, 2^{s+t} - 1, 2^s - 1]$,
$v_{t,s,15} = [2^s - 1, 0, 2^{s+t} - 1]$, $v_{t,s,16} = [2^s - 1, 2^{s+t} - 1, 0]$,
$v_{t,s,17} = [2^{s+1} - 1, 0, 2^{s+t} - 2^s - 1]$, $v_{t,s,18} = [2^{s+1} - 1, 2^{s+t} - 2^s - 1, 0]$,
$v_{t,s,19} = [2^{s+t} - 1, 0, 2^s - 1]$, $v_{t,s,20} = [2^{s+t} - 1, 2^s - 1, 0]$,
$v_{t,2,21} = [3, 3, 2^{t+2} - 4]$, $s = 2$, $v_{t,s,21} = [3, 2^s - 3, 2^{s+t} - 2]$, $s \geqslant 3$.

Hence, we easily obtain

Proposition 7.4.1. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+t}+2^s-2}$ is an $\mathbb{F}_2$-vector space of dimension 66 with a basis consisting of all the classes represented by the monomials $b_{t,s,j}, j \geqslant 1$, which are determined as follows:*

For $s \geqslant 2$,

1. $(0, 1, 2^s - 2, 2^{s+t} - 1)$, 2. $(0, 1, 2^{s+t} - 1, 2^s - 2)$,
3. $(0, 2^{s+t} - 1, 1, 2^s - 2)$, 4. $(1, 0, 2^s - 2, 2^{s+t} - 1)$,
5. $(1, 0, 2^{s+t} - 1, 2^s - 2)$, 6. $(1, 2^s - 2, 0, 2^{s+t} - 1)$,
7. $(1, 2^s - 2, 2^{s+t} - 1, 0)$, 8. $(1, 2^{s+t} - 1, 0, 2^s - 2)$,
9. $(1, 2^{s+t} - 1, 2^s - 2, 0)$, 10. $(2^{s+t} - 1, 0, 1, 2^s - 2)$,
11. $(2^{s+t} - 1, 1, 0, 2^s - 2)$, 12. $(2^{s+t} - 1, 1, 2^s - 2, 0)$,
13. $(0, 1, 2^s - 1, 2^{s+t} - 2)$, 14. $(0, 1, 2^{s+t} - 2, 2^s - 1)$,
15. $(0, 2^s - 1, 1, 2^{s+t} - 2)$, 16. $(1, 0, 2^s - 1, 2^{s+t} - 2)$,
17. $(1, 0, 2^{s+t} - 2, 2^s - 1)$, 18. $(1, 2^s - 1, 0, 2^{s+t} - 2)$,
19. $(1, 2^s - 1, 2^{s+t} - 2, 0)$, 20. $(1, 2^{s+t} - 2, 0, 2^s - 1)$,
21. $(1, 2^{s+t} - 2, 2^s - 1, 0)$, 22. $(2^s - 1, 0, 1, 2^{s+t} - 2)$,
23. $(2^s - 1, 1, 0, 2^{s+t} - 2)$, 24. $(2^s - 1, 1, 2^{s+t} - 2, 0)$,
25. $(0, 0, 2^s - 1, 2^{s+t} - 1)$, 26. $(0, 0, 2^{s+t} - 1, 2^s - 1)$,
27. $(0, 2^s - 1, 0, 2^{s+t} - 1)$, 28. $(0, 2^s - 1, 2^{s+t} - 1, 0)$,
29. $(0, 2^{s+t} - 1, 0, 2^s - 1)$, 30. $(0, 2^{s+t} - 1, 2^s - 1, 0)$,
31. $(2^s - 1, 0, 0, 2^{s+t} - 1)$, 32. $(2^s - 1, 0, 2^{s+t} - 1, 0)$,
33. $(2^s - 1, 2^{s+t} - 1, 0, 0)$, 34. $(2^{s+t} - 1, 0, 0, 2^s - 1)$,
35. $(2^{s+t} - 1, 0, 2^s - 1, 0)$, 36. $(2^{s+t} - 1, 2^s - 1, 0, 0)$,
37. $(0, 1, 2^{s+1} - 2, 2^{s+t} - 2^s - 1)$, 38. $(1, 0, 2^{s+1} - 2, 2^{s+t} - 2^s - 1)$,
39. $(1, 2^{s+1} - 2, 0, 2^{s+t} - 2^s - 1)$, 40. $(1, 2^{s+1} - 2, 2^{s+t} - 2^s - 1, 0)$,
41. $(0, 1, 2^{s+1} - 1, 2^{s+t} - 2^s - 2)$, 42. $(0, 2^{s+1} - 1, 1, 2^{s+t} - 2^s - 2)$,
43. $(1, 0, 2^{s+1} - 1, 2^{s+t} - 2^s - 2)$, 44. $(1, 2^{s+1} - 1, 0, 2^{s+t} - 2^s - 2)$,
45. $(1, 2^{s+1} - 1, 2^{s+t} - 2^s - 2, 0)$, 46. $(2^{s+1} - 1, 0, 1, 2^{s+t} - 2^s - 2)$,
47. $(2^{s+1} - 1, 1, 0, 2^{s+t} - 2^s - 2)$, 48. $(2^{s+1} - 1, 1, 2^{s+t} - 2^s - 2, 0)$,
49. $(0, 0, 2^{s+1} - 1, 2^{s+t} - 2^s - 1)$, 50. $(0, 2^{s+1} - 1, 0, 2^{s+t} - 2^s - 1)$,
51. $(0, 2^{s+1} - 1, 2^{s+t} - 2^s - 1, 0)$, 52. $(2^{s+1} - 1, 0, 0, 2^{s+t} - 2^s - 1)$,
53. $(2^{s+1} - 1, 0, 2^{s+t} - 2^s - 1, 0)$, 54. $(2^{s+1} - 1, 2^{s+t} - 2^s - 1, 0, 0)$,

55. $(0, 3, 2^{s+t} - 3, 2^s - 2)$, 56. $(3, 0, 2^{s+t} - 3, 2^s - 2)$,
57. $(3, 2^{s+t} - 3, 0, 2^s - 2)$, 58. $(3, 2^{s+t} - 3, 2^s - 2, 0)$,
59. $(0, 3, 2^{s+1} - 3, 2^{s+t} - 2^s - 2)$, 60. $(3, 0, 2^{s+1} - 3, 2^{s+t} - 2^s - 2)$,
61. $(3, 2^{s+1} - 3, 0, 2^{s+t} - 2^s - 2)$, 62. $(3, 2^{s+1} - 3, 2^{s+t} - 2^s - 2, 0)$.

For $s = 2$,

63. $(0, 3, 3, 2^{t+2} - 4)$, 64. $(3, 0, 3, 2^{t+2} - 4)$,
65. $(3, 3, 0, 2^{t+2} - 4)$, 66. $(3, 3, 2^{t+2} - 4, 0)$.

For $s \geqslant 3$,

63. $(0, 3, 2^s - 3, 2^{s+t} - 2)$, 64. $(3, 0, 2^s - 3, 2^{s+t} - 2)$,
65. $(3, 2^s - 3, 0, 2^{s+t} - 2)$, 66. $(3, 2^s - 3, 2^{s+t} - 2, 0)$.

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t}+2^s-2}$. We have

Theorem 7.4.2. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t}+2^s-2}$ is an $\mathbb{F}_2$-vector space with a basis consisting of all the classes represented by the following monomials:*

1. $\phi(c_{t,i}), 1 \leqslant i \leqslant \rho_2(t)$ and $\phi(d_{t,j}), 1 \leqslant j \leqslant \rho_3(t)$, for $s = 2$.

2. $\phi(a_{t,s-2,j}), 1 \leqslant j \leqslant \mu_{t+1}(s-2)$ for $s \geqslant 3$.

Here $c_{t,i}, d_{t,j}$, $\rho_2(t), \rho_3(t)$ are defined as in Section 3, $a_{t,s-2,j}$ is defined as in Section 6, the homomorphism ϕ is defined as in Definition 2.9. By convention, we set $\mu_{t+1}(s-2) = \mu_5(s-2)$ for $t \geqslant 4$.

3. The monomials $b_{t,s,j}, j \geqslant 67$, are determined as follows:

For $s \geqslant 2$,

67. $(1, 1, 2^s - 2, 2^{s+t} - 2)$, 68. $(1, 1, 2^{s+t} - 2, 2^s - 2)$,
69. $(1, 2^s - 2, 1, 2^{s+t} - 2)$, 70. $(1, 2^{s+t} - 2, 1, 2^s - 2)$,
71. $(1, 2, 2^{s+t} - 3, 2^s - 2)$, 72. $(1, 1, 2^{s+1} - 2, 2^{s+t} - 2^s - 2)$,
73. $(1, 2^{s+1} - 2, 1, 2^{s+t} - 2^s - 2)$, 74. $(1, 2, 2^s - 1, 2^{s+t} - 4)$,
75. $(1, 2, 2^{s+t} - 4, 2^s - 1)$, 76. $(1, 3, 2^s - 2, 2^{s+t} - 4)$,
77. $(1, 3, 2^{s+t} - 4, 2^s - 2)$, 78. $(3, 1, 2^s - 2, 2^{s+t} - 4)$,
79. $(3, 1, 2^{s+t} - 4, 2^s - 2)$, 80. $(1, 2, 2^{s+1} - 4, 2^{s+t} - 2^s - 1)$,
81. $(1, 2, 2^{s+1} - 3, 2^{s+t} - 2^s - 2)$, 82. $(1, 2, 2^{s+1} - 1, 2^{s+t} - 2^s - 4)$,
83. $(1, 2^{s+1} - 1, 2, 2^{s+t} - 2^s - 4)$, 84. $(2^{s+1} - 1, 1, 2, 2^{s+t} - 2^s - 4)$,
85. $(1, 3, 2^{s+1} - 4, 2^{s+t} - 2^s - 2)$, 86. $(3, 1, 2^{s+1} - 4, 2^{s+t} - 2^s - 2)$,
87. $(1, 3, 2^{s+1} - 2, 2^{s+t} - 2^s - 4)$, 88. $(3, 1, 2^{s+1} - 2, 2^{s+t} - 2^s - 4)$,
89. $(3, 2^{s+1} - 3, 2, 2^{s+t} - 2^s - 4)$.

For $s = 2$ *and* $t \geqslant 2$, $b_{t,2,90} = (3, 3, 4, 2^{t+2} - 8)$.

For $s = 2$ *and* $t = 2$, $b_{2,2,91} = (3, 5, 8, 2)$.

For $s \geqslant 3$,

90. $(1, 2, 2^s - 4, 2^{s+t} - 1)$, 91. $(1, 2, 2^{s+t} - 1, 2^s - 4)$,
92. $(1, 2^{s+t} - 1, 2, 2^s - 4)$, 93. $(2^{s+t} - 1, 1, 2, 2^s - 4)$,
94. $(1, 2, 2^s - 3, 2^{s+t} - 2)$, 95. $(1, 3, 2^s - 4, 2^{s+t} - 2)$,
96. $(1, 3, 2^{s+t} - 2, 2^s - 4)$, 97. $(3, 1, 2^s - 4, 2^{s+t} - 2)$,
98. $(3, 1, 2^{s+t} - 2, 2^s - 4)$, 99. $(1, 2^s - 1, 2, 2^{s+t} - 4)$,

100. $(2^s-1,1,2,2^{s+t}-4)$, 101. $(3,2^{s+t}-3,2,2^s-4)$,
102. $(3,2^s-3,2,2^{s+t}-4)$, 103. $(3,5,2^{s+t}-6,2^s-4)$,
104. $(3,5,2^{s+1}-6,2^{s+t}-2^s-4)$.

For $s=3$, $b_{t,3,105}=(3,5,6,2^{t+3}-8)$.

For $s\geqslant 4$, $b_{t,s,105}=(3,5,2^s-6,2^{s+t}-4)$.

We prove the theorem by proving the following propositions.

Proposition 7.4.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2\underset{\mathcal{A}}{\otimes}R_4)_{2^{s+t}+2^s-2}$ is generated by the elements listed in Theorem 7.4.2.*

The proof of this proposition is based on the following lemmas.

Lemma 7.4.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}0&1&1&0\\0&1&0&1\\0&1&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\1&0&0&1\\1&0&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&0&1\\1&0&0&0\\0&1&0&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&0\\1&0&0&0\\0&1&0&0\end{pmatrix}$$

$$\begin{pmatrix}0&1&1&0\\0&1&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}0&1&1&0\\0&1&1&0\\0&1&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}0&1&0&1\\0&1&0&1\\0&1&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\1&0&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}$$

$$\begin{pmatrix}1&1&0&0\\1&1&0&0\\0&1&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&1&0&0\\0&1&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\1&0&1&0\\1&0&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&0&0&1\\1&0&0&1\\1&0&0&0\\0&0&1&0\end{pmatrix}$$

$$\begin{pmatrix}1&1&0&0\\1&1&0&0\\1&0&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&1&0&0\\1&0&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&0&0&1\\1&0&0&1\\1&0&0&0\\0&1&0&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\1&0&1&0\\1&0&0&0\\0&1&0&0\end{pmatrix}.$$

Proof. See [8] □

Lemma 7.4.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&0&1&0\\1&0&0&1\\0&1&0&0\\0&1&0&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\0&1&0&1\\0&1&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\0&1&0&1\\0&1&0&0\\0&0&1&0\end{pmatrix}$$

$$\begin{pmatrix}1&1&0&0\\1&0&0&1\\1&0&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\1&0&0&1\\1&0&0&0\\0&1&0&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&0\\1&0&0&1\\0&1&0&0\\0&0&0&1\end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(3,12,1,2),(1,6,9,2),(1,7,8,2),(7,1,8,2),(7,8,1,2),$$
$$(3,4,1,10),(1,6,3,8),(1,6,8,3),(3,4,9,2).$$

A direct computation shows

$$\begin{aligned}
(3,12,1,2) &= Sq^1\big((3,11,1,2)+(3,7,3,4)+(1,7,5,4)\big)\\
&\quad + Sq^2\big((2,11,1,2)+(5,7,2,2)+(2,7,3,4)+(1,7,6,2)\big)\\
&\quad + Sq^4(3,7,2,2)+(2,13,1,2)+(2,11,1,4)+(3,9,4,2)+(3,9,2,4)\\
&\quad + (2,9,3,4)+(3,8,3,4)+(1,9,6,2)+(1,7,8,2) \bmod \mathcal{L}_4(2;2;1;1),\\
(1,6,9,2) &= Sq^1\big((1,5,7,4)+(1,3,9,4)+(4,3,7,3)+(1,5,8,3)\\
&\quad + (1,3,8,5)+(1,4,9,3)+(1,4,7,5)\big)+Sq^2\big((1,6,7,2)\\
&\quad + (2,3,7,4)+(2,3,8,3)+(2,4,7,3)+(1,2,7,6)\big)+Sq^4(1,4,7,2)\\
&\quad + (1,4,11,2)+(1,3,10,4)+(1,6,8,3)+(1,3,8,6)\\
&\quad + (1,4,10,3)+(1,2,9,6)+(1,2,7,8) \quad \bmod \mathcal{L}_4(2;2;1;1),\\
(1,7,8,2) &= Sq^1\big((1,7,5,4)+(1,5,3,8)+(4,7,3,3)+(1,8,5,3)\\
&\quad + (1,8,3,5)+(1,9,4,3)+(1,7,4,5)\big)+Sq^2\big((1,7,6,2)+(2,7,3,4)\\
&\quad + (2,8,3,3)+(2,7,4,3)+(1,7,2,6)\big)+Sq^4\big((1,5,6,2)+(2,5,3,4)\\
&\quad + (1,4,6,3)+(1,4,3,6)+(1,6,4,3)+(1,5,2,6)\big)+(1,5,10,2)\\
&\quad + (1,6,3,8)+(1,4,10,3)+(1,4,3,10)+(1,6,8,3)\\
&\quad + (1,7,2,8)+(1,5,2,10) \quad \bmod \mathcal{L}_4(2;2;1;1),\\
(7,1,8,2) &= \overline{\varphi}_1(1,7,8,2),\\
(7,8,1,2) &= Sq^1\big((7,5,1,4)+(7,3,4,3)+(7,1,4,5)\big)+Sq^2\big((7,6,1,2)\\
&\quad + (7,3,2,4)+(7,2,4,3)+(7,1,2,6)\big)+Sq^4\big((5,6,1,2)+(5,3,2,4)\\
&\quad + (4,3,4,3)+(5,2,4,3)+(5,1,2,6)\big)+(5,10,1,2)+(5,3,2,8)\\
&\quad + (4,3,8,3)+(5,2,8,3)+(7,1,2,8)+(5,1,2,10) \quad \bmod \mathcal{L}_4(2;2;1;1),\\
(3,4,1,10) &= Sq^1\big((3,4,1,9)+(3,3,4,7)+(1,5,4,7)+(3,1,2,11)\big)\\
&\quad + Sq^2\big((5,2,2,7)+(2,3,4,7)+(1,6,2,7)+(2,1,2,11)\big)\\
&\quad + Sq^4(3,2,2,7)+(3,2,4,9)+(3,3,4,8)+(2,3,4,9)+(2,1,2,13)\\
&\quad + (1,8,2,7)+(1,6,2,9)+(3,1,2,12)+(2,1,4,11) \bmod \mathcal{L}_4(2;2;1;1),
\end{aligned}$$

$$\begin{aligned}(1,6,3,8) &= Sq^1\big((2,5,3,7)+(2,3,5,7)+(2,3,3,9)\big)\\ &\quad + Sq^2\big((1,5,3,7)+(1,6,2,7)+(1,3,5,7)+(1,2,6,7)+(1,3,3,9)\big)\\ &\quad + Sq^4(1,4,2,7)+(1,6,2,9)+(1,4,2,11)+(1,2,8,7)+(1,2,6,9)\\ &\quad + (1,3,6,8)+(1,4,3,10)+(1,3,4,10) \mod \mathcal{L}_4(2;2;1;1),\\ (1,6,8,3) &= Sq^1\big((1,6,5,5)+(1,8,3,5)+(4,5,3,5)+(1,5,3,8)\\ &\quad + (1,5,5,6)\big)+Sq^2\big((1,6,6,3)+(2,6,3,5)+(2,5,3,6)+(1,3,6,6)\big)\\ &\quad + Sq^4\big((1,4,6,3)+(1,4,3,6)\big)+(1,4,10,3)+(1,4,3,10)\\ &\quad + (1,6,3,8)+(1,3,8,6)+(1,3,6,8) \mod \mathcal{L}_4(2;2;1;1),\\ (3,4,9,2) &= Sq^1\big((3,1,11,2)+(3,4,7,3)+(1,4,7,5)\big)\\ &\quad + Sq^2\big((5,2,7,2)+(2,1,11,2)+(2,4,7,3)+(1,2,7,6)\big)\\ &\quad + Sq^4(3,2,7,2)+(3,2,9,4)+(3,1,12,2)+(2,1,13,2)+(2,1,11,4)\\ &\quad + (3,4,8,3)+(2,4,9,3)+(1,2,7,8)+(1,2,9,6) \mod \mathcal{L}_4(2;2;1;1).\end{aligned}$$

The lemma is proved. □

Lemma 7.4.6. *The following matrix is strictly inadmissible*

$$\begin{pmatrix}1&1&0&0\\1&0&1&0\\0&1&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrix is (3,5,14,16). By a direct calculation, we have

$$\begin{aligned}(3,5,14,16) &= Sq^1\big((3,3,17,14)+(3,3,14,17)\big)\\ &\quad + Sq^2\big((5,3,14,14)+(2,3,17,14)+(2,3,14,17)\big)\\ &\quad + Sq^4(3,3,14,14)+Sq^8(3,5,8,14)+(3,5,8,22)\\ &\quad + (3,4,17,14)+(3,4,14,17)+(2,5,17,14)\\ &\quad + (2,5,14,17) \mod \mathcal{L}_4(2;2;2;1;1).\end{aligned}$$

The lemma follows. □

Lemma 7.4.7. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&1&0&0\\1&0&0&1\\0&1&0&0\\0&0&1&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&0&1\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&0\\1&0&1&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (3,5,24,2), (3,5,8,18), (3,5,10,16). By a direct calculation, we have

$$\begin{aligned}
(3,5,24,2) &= Sq^1(3,3,25,2) + Sq^2\big((5,3,22,2)+(2,3,25,2)\big)\\
&\quad + Sq^4\big((3,9,14,4)+(3,3,22,2)\big) + Sq^8(3,5,14,4)\\
&\quad + (3,5,18,8) + (3,3,24,4) + (3,4,25,2)\\
&\quad + (2,5,25,2) + (2,3,25,4) \mod \mathcal{L}_4(2;2;1;1;1),\\
(3,5,8,18) &= Sq^1(3,3,1,26) + Sq^2\big((5,3,2,22)+(2,3,1,26)\big)\\
&\quad + Sq^4\big((3,9,4,14)+(3,3,2,22)\big) + Sq^8(3,5,4,14)\\
&\quad + (3,5,2,24) + (3,3,4,24) + (3,4,1,26)\\
&\quad + (2,5,1,26) + (2,3,1,28) \mod \mathcal{L}_4(2;2;1;1;1),\\
(3,5,10,16) &= Sq^1(3,3,17,10) + Sq^2\big((5,3,18,6)+(2,3,17,10)\big)\\
&\quad + Sq^4\big((3,3,18,6)+(3,9,12,6)\big) + Sq^8\big((3,5,10,8)+(3,5,12,6)\\
&\quad + (3,5,8,10)\big) + (3,3,20,8) + (3,5,8,18) + (3,4,17,10)\\
&\quad + (2,5,17,10) + (2,3,17,12) \mod \mathcal{L}_4(2;2;1;1;1).
\end{aligned}$$

The lemma is proved. □

Using the results in Section 5, Section 6, Lemmas 7.4.4-7.4.7, 7.1.5, 7.1.1 and Theorem 2.5, we get Proposition 7.4.3.

Now, we show that the elements listed in Theorem 7.4.2 are linearly independent.

Proposition 7.4.8. *The elements* $[b_{2,2,i}], 67 \leqslant i \leqslant 91$, $[\phi(c_{2,j})], 1 \leqslant j \leqslant 26$, *and* $[\phi(d_{2,\ell})], 1 \leqslant \ell \leqslant 9$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{18}$.

Proof. Suppose that there is a linear relation

$$\sum_{i=67}^{91} \gamma_i[b_{2,2,i}] + \sum_{j=1}^{26} \eta_j[\phi(c_{2,j})] + \sum_{\ell=1}^{9} \xi_\ell[\phi(d_{2,\ell})] = 0, \tag{7.4.8.1}$$

with $\gamma_i, \eta_j, \xi_\ell \in \mathbb{F}_2$.

Apply the squaring operation Sq^0_* to (7.4.8.1) and we get

$$\sum_{j=1}^{26} \eta_j[c_{2,j}] + \sum_{\ell=1}^{9} \xi_\ell[d_{2,j}] = 0.$$

Since $\{[c_{3,i}]; 1 \leqslant i \leqslant 26\} \cup \{[d_{3,j}]; 1 \leqslant j \leqslant 9\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_7$, we obtain $\eta_i = 0, 1 \leqslant i \leqslant 26, \xi_j = 0, 1 \leqslant j \leqslant 9$. Hence, (7.4.8.1) becomes

$$\sum_{i=67}^{91} \gamma_i[b_{2,2,i}] = 0. \tag{7.4.8.2}$$

Now, we prove $\gamma_i = 0, 67 \leqslant i \leqslant 91$.

Applying the homomorphisms f_1, f_6, to the relation (7.4.8.2), we obtain

$$\begin{aligned}
&\gamma_{70}[15,1,2] + \gamma_{\{75,80,82\}}[1,3,14] + \gamma_{75}[1,14,3]\\
&\quad + \gamma_{69}[3,1,14] + \gamma_{80}[1,6,11] + \gamma_{82}[1,7,10] + \gamma_{73}[7,1,10]\\
&\quad + \gamma_{71}[3,13,2] + \gamma_{\{74,75,80,82\}}[3,3,12] + \gamma_{81}[3,5,10] = 0,\\
&\gamma_{\{69,71,74,75,80,81,82\}}[1,2,15] + \gamma_{\{76,77,85,87\}}[1,3,14] + \gamma_{70}[1,14,3]\\
&\quad + \gamma_{\{78,79,86,88\}}[3,1,14] + \gamma_{73}[1,6,11] + \gamma_{83}[1,7,10]\\
&\quad + \gamma_{84}[7,1,10] + \gamma_{90}[3,3,12] + \gamma_{\{89,91\}}[3,5,10] = 0.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
\gamma_i = 0, \ i = 69, 70, 71, 73, 74, 75, 81, 82, 83, 84, 90,\\
\gamma_{\{76,77,85,87\}} = \gamma_{\{78,79,86,88\}} = \gamma_{\{89,91\}} = 0.
\end{cases} \tag{7.4.8.3}$$

With the aid of (7.4.8.3), the homomorphisms f_2, f_3 send (7.4.8.2) to

$$\begin{aligned}
&\gamma_{\{68,79\}}[15,1,2] + \gamma_{\{67,87,91\}}[3,1,14] + \gamma_{\{72,86,87,91\}}[7,1,10]\\
&\quad + \gamma_{\{77,91\}}[3,13,2] + \gamma_{\{76,77,85,87\}}[3,3,12] + \gamma_{85}[3,5,10] = 0,\\
&\gamma_{\{67,78\}}[15,1,2] + \gamma_{\{68,85,89\}}[3,1,14] + \gamma_{\{72,85,88,89\}}[7,1,10]\\
&\quad + \gamma_{\{76,89\}}[3,13,2] + \gamma_{\{76,77,85,87\}}[3,3,12] + \gamma_{87}[3,5,10] = 0.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
\gamma_{85} = \gamma_{87} = 0,\\
\gamma_{\{68,79\}} = \gamma_{\{67,91\}} = \gamma_{\{72,86,91\}} = \gamma_{\{77,91\}} = \gamma_{\{76,77\}} = 0,\\
\gamma_{\{67,78\}} = \gamma_{\{68,89\}} = \gamma_{\{72,88,89\}} = \gamma_{\{76,89\}} = 0.
\end{cases} \tag{7.4.8.4}$$

With the aid of (7.4.8.3) and (7.4.8.4), the homomorphism f_4 send (7.4.8.2) to

$$\gamma_{88}[1,3,14] + \gamma_{\{67,72,88\}}[1,7,10] + \gamma_{88}[3,3,12] + \gamma_{86}[3,5,10] = 0.$$

From this equality, we get

$$\gamma_{86} = \gamma_{88} = \gamma_{\{67,72\}} = 0. \tag{7.4.8.5}$$

Combining (7.4.8.3), (7.4.8.4) and (7.4.8.5), we obtain

$$\begin{cases}
\gamma_i = 0, i \neq 67, 68, 72, 76, 77, 78, 79, 89, 91,\\
\gamma_{67} = \gamma_{68} = \gamma_{72} = \gamma_{76} = \gamma_{77} = \gamma_{78} = \gamma_{79} = \gamma_{89} = \gamma_{91}
\end{cases} \tag{7.4.8.6}$$

Substituting (7.4.8.6) into the relation (7.4.8.2), we obtain

$$\gamma_{67}[\theta] = 0, \tag{7.4.8.7}$$

where $\theta = b_{2,2,67} + b_{2,2,68} + b_{2,2,72} + b_{2,2,76} + b_{2,2,77} + b_{2,2,78} + b_{2,2,79} + b_{2,2,89} + b_{2,2,91}$.

Now, we prove that the polynomial θ is non-hit. Suppose the contrary, that θ is hit. Then, we have

$$\theta = Sq^1(A) + Sq^2(B) + Sq^4(C) + Sq^8(D),$$

for some polynomials A, B, C, D in R_4. Let $(Sq^2)^3$ act on the both sides of the above equality. We get

$$(Sq^2)^3(\theta) = (Sq^2)^3Sq^4(C) + (Sq^2)^3Sq^8(D). \tag{7.4.8.8}$$

On the other hand, by a direct computation, it is not difficult to check that the relation (7.4.8.8) is not true for all $C \in (R_4)_{14}$ and $D \in (R_4)_{10}$. This is a contradiction. Hence, $[\theta] \neq 0$ and the relation (7.4.8.7) implies $\gamma_{67} = 0$. The proposition is proved. □

Proposition 7.4.9. *For $t \geqslant 3$, the elements $[b_{t,2,i}]$, for $67 \leqslant i \leqslant 90$, $[\phi(c_{t,j})]$, for $1 \leqslant j \leqslant \rho_2(t)$, and $[\phi(d_{t,\ell})]$, for $1 \leqslant \ell \leqslant \rho_3(t)$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+2}+2}$.*

Proof. By a simple computation, we have

$$Sq^0_*([b_{t,2,i}]) = 0, Sq^0_*([\phi(c_{t,j}]) = [c_{t,j}], Sq^0_*([\phi(d_{t,\ell}]) = [d_{t,\ell}].$$

Since $\{[c_{t,j}]; 1 \leqslant j \leqslant \rho_2(t)\} \cup \{[d_{t,\ell}]; 1 \leqslant \ell \leqslant \rho_3(t)\}$ is a basis of the space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{t+1}-1}$, it suffices to show that the elements $[b_{t,2,i}], 67 \leqslant i \leqslant 90$, are linearly independent.

Suppose that there is a linear relation

$$\sum_{i=67}^{90} \gamma_i[b_{t,2,i}] = 0, \tag{7.4.9.1}$$

with $\gamma_i \in \mathbb{F}_2$. We prove $\gamma_i = 0, 67 \leqslant i \leqslant 90$.

Apply the homomorphisms f_1, f_6 to the relation (7.4.9.1) and we obtain

$$\begin{aligned}
&\gamma_{70}v_{t,2,3} + \gamma_{\{75,80,82\}}v_{t,2,4} + \gamma_{75}v_{t,2,5} + \gamma_{69}v_{t,2,6} + \gamma_{80}v_{t,2,7}\\
&\quad + \gamma_{82}v_{t,2,8} + \gamma_{73}v_{t,2,9} + \gamma_{81}v_{t,2,10} + \gamma_{71}v_{t,2,11} + \gamma_{\{74,75,80,82\}}v_{t,2,21} = 0,\\
&\gamma_{\{69,71,74,75,80,81,82\}}v_{t,2,1} + \gamma_{\{76,77,85,87\}}v_{t,2,4} + \gamma_{70}v_{t,2,5} + \gamma_{73}v_{t,2,7}\\
&\quad + \gamma_{\{78,79,86,88\}}v_{t,2,6} + \gamma_{83}v_{t,2,8} + \gamma_{84}v_{t,2,9} + \gamma_{89}v_{t,2,10} + \gamma_{90}v_{t,2,21} = 0.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
\gamma_i = 0, \ i = 69, 70, 71, 73, 74, 75, 80, 81, 82, 83, 84, 89, 90,\\
\gamma_{\{76,77,85,87\}} = \gamma_{\{78,79,86,88\}} = 0.
\end{cases} \tag{7.4.9.2}$$

With the aid of (7.4.9.2), the homomorphisms f_2, f_3, f_4 send (7.4.9.1) to

$$\begin{aligned}
&\gamma_{\{68,79\}}v_{t,2,3} + \gamma_{\{67,87\}}v_{t,2,6} + \gamma_{\{72,86,87\}}v_{t,2,9}\\
&\quad + \gamma_{85}v_{t,2,10} + \gamma_{77}v_{t,2,11} + \gamma_{\{76,77,85,87\}}v_{t,2,21} = 0,\\
&\gamma_{\{67,78\}}v_{t,2,3} + \gamma_{\{68,85\}}v_{t,2,6} + \gamma_{\{72,85,88\}}v_{t,2,9}\\
&\quad + \gamma_{87}v_{t,2,10} + \gamma_{76}v_{t,2,11} + \gamma_{\{76,77,85,87\}}v_{t,2,21} = 0,\\
&\gamma_{\{68,77\}}v_{t,2,2} + \gamma_{\{67,88\}}v_{t,2,4} + \gamma_{\{72,85,88\}}v_{t,2,8}\\
&\quad + \gamma_{86}v_{t,2,10} + \gamma_{79}v_{t,2,11} + \gamma_{\{78,88\}}v_{t,2,21} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\begin{cases} \gamma_i = 0,\ i = 76, 77, 79, 85, 86, 87, \\ \gamma_{\{68,79\}} = \gamma_{\{67,87\}} = \gamma_{\{72,86,87\}} = \gamma_{\{76,77,85,87\}} = 0, \\ \gamma_{\{67,78\}} = \gamma_{\{68,85\}} = \gamma_{\{72,85,88\}} = \gamma_{\{76,77,85,87\}} = 0, \\ \gamma_{\{68,77\}} = \gamma_{\{67,88\}} = \gamma_{\{72,85,88\}} = \gamma_{\{78,88\}} = 0. \end{cases} \tag{7.4.9.3}$$

Combining (7.4.9.2) and (7.4.9.3), we get $\gamma_i = 0$ for any i. The proposition is proved. □

Proposition 7.4.10. *For $t \geqslant 2$, the elements $[b_{t,3,i}], 67 \leqslant i \leqslant 105$, and $[\phi(a_{t,1,j})], 1 \leqslant j \leqslant \nu(t,3)$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+3}+6}$. Here $\nu(2,3) = 87, \nu(3,3) = 136, \nu(t,3) = 150$ for $t \geqslant 4$.*

Proof. Suppose that there is a linear relation

$$\sum_{i=67}^{105} \gamma_i [b_{t,3,i}] + \sum_{j=1}^{\nu(t,3)} \eta_j [\phi(a_{t,1,j})] = 0, \tag{7.4.10.1}$$

with $\gamma_i, \eta_j \in \mathbb{F}_2$.

Applying the squaring operation Sq^0_* to (7.4.10.1), we get

$$\sum_{j=1}^{\nu(t,3)} \eta_j [a_{t,1,j}] = 0.$$

Since $\{[a_{t,1,j}]; 1 \leqslant j \leqslant \nu(t,3)\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{2^{t+2}+1}$, we obtain $\eta_j = 0, 1 \leqslant j \leqslant \nu(t,3)$. Hence, (7.4.10.1) becomes

$$\sum_{i=67}^{105} \gamma_i [b_{t,2,i}] = 0. \tag{7.4.10.2}$$

Now, we prove $\gamma_i = 0, 67 \leqslant i \leqslant 105$.

Applying the homomorphisms f_1, f_6 to the relation (7.4.10.2), we obtain

$$\begin{aligned} &\gamma_{90} v_{t,3,1} + \gamma_{91} v_{t,3,2} + \gamma_{70} v_{t,3,3} + \gamma_{74} v_{t,3,4} \\ &\quad + \gamma_{75} v_{t,3,5} + \gamma_{69} v_{t,3,6} + \gamma_{80} v_{t,3,7} + \gamma_{82} v_{t,3,8} \\ &\quad + \gamma_{73} v_{t,3,9} + \gamma_{81} v_{t,3,10} + \gamma_{71} v_{t,3,11} + \gamma_{94} v_{t,3,21} = 0, \\ &\gamma_{69} v_{t,3,1} + \gamma_{92} v_{t,3,2} + \gamma_{93} v_{t,3,3} + \gamma_{99} v_{t,3,4} + \gamma_{70} v_{t,3,5} \\ &\quad + \gamma_{100} v_{t,3,6} + \gamma_{73} v_{t,3,7} + \gamma_{83} v_{t,3,8} + \gamma_{84} v_{t,3,9} \\ &\quad + \gamma_{89} v_{t,3,10} + \gamma_{101} v_{t,3,11} + \gamma_{\{102,103,104,105\}} v_{t,3,21} = 0. \end{aligned}$$

From these equalities, we get

$$\begin{cases} \gamma_i = 0,\ i = 69, 70, 71, 73, 74, 75, 80, 81, \\ 82, 83, 84, 89, 90, 91, 92, 93, 94, 99, 100, 101, \\ \gamma_{\{102,103,104,105\}} = 0. \end{cases} \tag{7.4.10.3}$$

With the aid of (7.4.10.3), the homomorphisms f_4, f_5 send (7.4.10.2) to

$$\begin{aligned}
&\gamma_{\{68,77,98,103\}}v_{t,3,2} + \gamma_{\{67,78,95,102\}}v_{t,3,4} + \gamma_{\{72,85,88,104\}}v_{t,3,8}\\
&\quad + \gamma_{86}v_{t,3,10} + \gamma_{79}v_{t,3,11} + \gamma_{97}v_{t,3,21} = 0,\\
&\gamma_{\{67,76,97\}}v_{t,3,2} + \gamma_{\{68,79,96\}}v_{t,3,4} + \gamma_{\{72,86,87\}}v_{t,3,8}\\
&\quad + \gamma_{88}v_{t,3,10} + \gamma_{\{78,105\}}v_{t,3,11} + \gamma_{98}v_{t,3,21} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\begin{cases}
\gamma_{79} = \gamma_{86} = \gamma_{88} = \gamma_{97} = \gamma_{98} = 0,\\
\gamma_{\{68,77,98,103\}} = \gamma_{\{67,78,95,102\}} = \gamma_{\{72,85,88,104\}} = 0,\\
\gamma_{\{67,76,97\}} = \gamma_{\{68,79,96\}} = \gamma_{\{72,86,87\}} = \gamma_{\{78,105\}} = 0.
\end{cases} \tag{7.4.10.4}$$

With the aid of (7.4.10.3) and (7.4.10.4), the homomorphisms f_2, f_3 send the relation (7.4.10.2) respectively to

$$\begin{aligned}
&\gamma_{\{68,96\}}v_{t,3,3} + \gamma_{\{67,76\}}v_{t,3,6} + \gamma_{\{72,87\}}v_{t,3,9}\\
&\quad + \gamma_{85}v_{t,3,10} + \gamma_{77}v_{t,3,11} + \gamma_{\{77,85,95\}}v_{t,3,21} = 0,\\
&\gamma_{\{67,78,95,102\}}v_{t,3,3} + \gamma_{\{68,77,103\}}v_{t,3,6} + \gamma_{\{72,85,104\}}v_{t,3,9}\\
&\quad + \gamma_{87}v_{t,3,10} + \gamma_{\{76,105\}}v_{t,3,11} + \gamma_{\{76,87,96,105\}}v_{t,3,21} = 0.
\end{aligned}$$

The above equalities imply

$$\begin{cases}
\gamma_{77} = \gamma_{85} = \gamma_{87} = 0,\\
\gamma_{\{68,96\}} = \gamma_{\{67,76\}} = \gamma_{\{72,87\}} = \gamma_{\{77,85,95\}} = 0,\\
\gamma_{\{67,78,95,102\}} = \gamma_{\{68,77,103\}} = \gamma_{\{72,85,104\}} = 0,\\
\gamma_{\{76,87,96,105\}} = \gamma_{\{76,105\}} = 0.
\end{cases} \tag{7.4.10.5}$$

Combining (7.4.10.3), (7.4.10.4) and (7.4.10.5), we get $\gamma_i = 0$ for any i. The proposition is proved. □

Proposition 7.4.11. *For $t \geqslant 2, s \geqslant 4$, the elements $[b_{t,s,i}]$, for $67 \leqslant i \leqslant 105$, and $[\phi(a_{t,s-2,j})]$, for $1 \leqslant j \leqslant \nu(t,s)$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t}+2^s-2}$. Here $\nu(2,4) = 135, \nu(2,s) = 150$ for $s \geqslant 5$, $\nu(3,4) = 180, \nu(3,s) = 195$ for $s \geqslant 5$, $\nu(t,4) = 195, \nu(t,s) = 210$ for $t \geqslant 4, s \geqslant 5$.*

Proof. Suppose that there is a linear relation

$$\sum_{i=67}^{105} \gamma_i[b_{t,s,i}] + \sum_{j=1}^{\nu(t,s)} \eta_j[\phi(a_{t,s-2,j})] = 0, \tag{7.4.11.1}$$

with $\gamma_i, \eta_j \in \mathbb{F}_2$.

Applying the squaring operation Sq^0_* to (7.4.11.1), we get

$$\sum_{j=1}^{\nu(t,s)} \eta_j[a_{t,s-2,j}] = 0.$$

Since $\{[a_{t,s-2,j}]; 1 \leqslant j \leqslant \nu(t,s)\}$ is a basis of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{2^{s+t-1}+2^{s-1}-3}$, we obtain $\eta_j = 0, 1 \leqslant j \leqslant \nu(t,s)$. Hence, (7.4.11.1) becomes

$$\sum_{i=67}^{105} \gamma_i [b_{t,s,i}] = 0. \tag{7.4.11.2}$$

Now, we prove $\gamma_i = 0, 67 \leqslant i \leqslant 105$.

Apply the homomorphisms f_1, f_6 to the relation (7.4.11.2) and we obtain

$$\begin{aligned}
&\gamma_{90}v_{t,s,1} + \gamma_{91}v_{t,s,2} + \gamma_{70}v_{t,s,3} + \gamma_{74}v_{t,s,4} + \gamma_{75}v_{t,s,5} + \gamma_{69}v_{t,s,6}\\
&\quad + \gamma_{80}v_{t,s,7} + \gamma_{82}v_{t,s,8} + \gamma_{73}v_{t,s,9} + \gamma_{81}v_{t,s,10} + \gamma_{71}v_{t,s,11} + \gamma_{94}v_{t,s,21} = 0,\\
&\gamma_{69}v_{t,s,1} + \gamma_{92}v_{t,s,2} + \gamma_{93}v_{t,s,3} + \gamma_{99}v_{t,s,4} + \gamma_{70}v_{t,s,5} + \gamma_{100}v_{t,s,6}\\
&\quad + \gamma_{73}v_{t,s,7} + \gamma_{83}v_{t,s,8} + \gamma_{84}v_{t,s,9} + \gamma_{89}v_{t,s,10} + \gamma_{101}v_{t,s,11} + \gamma_{102}v_{t,s,21} = 0.
\end{aligned}$$

From these equalities, we get

$$\begin{cases} \gamma_i = 0, \ i = 69, 70, 71, 73, 74, 75, 80, 81, 82, 83, 84,\\ \qquad\qquad 89, 90, 91, 92, 93, 94, 99, 100, 101, 102. \end{cases} \tag{7.4.11.3}$$

With the aid of (7.4.11.3), the homomorphisms f_4, f_5 send (7.4.11.2) to

$$\begin{aligned}
&\gamma_{\{68,77,98,103\}}v_{t,s,2} + \gamma_{\{67,78,95,105\}}v_{t,s,4}\\
&\quad + \gamma_{\{72,85,88,104\}}v_{t,s,8} + \gamma_{86}v_{t,s,10} + \gamma_{79}v_{t,s,11} + \gamma_{97}v_{t,s,21} = 0,\\
&\gamma_{91}v_{t,s,1} + \gamma_{\{67,76,97\}}v_{t,s,2} + \gamma_{93}v_{t,s,3} + \gamma_{\{68,79,96\}}v_{t,s,4}\\
&\quad + \gamma_{74}v_{t,s,5} + \gamma_{100}v_{t,s,6} + \gamma_{82}v_{t,s,7} + \gamma_{\{72,86,87\}}v_{t,s,8} + \gamma_{84}v_{t,s,9}\\
&\quad + \gamma_{88}v_{t,s,10} + \gamma_{\{78,84,100\}}v_{t,s,11} + \gamma_{\{84,98,100\}}v_{t,s,21} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\begin{cases} \gamma_i = 0, \ i = 74, 79, 82, 84, 86, 88, 91, 93, 97, 100,\\ \gamma_{\{68,77,98,103\}} = \gamma_{\{67,78,95,105\}} = \gamma_{\{72,85,88,104\}} = 0,\\ \gamma_{\{67,76,97\}} = \gamma_{\{68,79,96\}} = \gamma_{\{72,86,87\}} = 0,\\ \gamma_{\{84,98,100\}} = \gamma_{\{78,84,100\}} = 0. \end{cases} \tag{7.4.11.4}$$

With the aid of (7.4.11.3) and (7.4.11.4), the homomorphisms f_2, f_3 send the relation (7.4.11.2) respectively to

$$\begin{aligned}
&\gamma_{\{68,96\}}v_{t,s,3} + \gamma_{\{67,76\}}v_{t,s,6} + \gamma_{\{72,87\}}v_{t,s,9}\\
&\quad + \gamma_{85}v_{t,s,10} + \gamma_{77}v_{t,s,11} + \gamma_{\{77,85,95\}}v_{t,s,21} = 0,\\
&\gamma_{\{67,78,95,105\}}v_{t,s,3} + \gamma_{\{68,77,98,103\}}v_{t,s,6} + \gamma_{87}v_{t,s,10}\\
&\quad + \gamma_{\{72,85,104\}}v_{t,s,9} + \gamma_{76}v_{t,s,11} + \gamma_{\{76,87,96\}}v_{t,s,21} = 0.
\end{aligned}$$

The above equalities imply

$$\begin{cases} \gamma_{76} = \gamma_{77} = \gamma_{85} = \gamma_{87} = 0,\\ \gamma_{\{68,96\}} = \gamma_{\{67,76\}} = \gamma_{\{77,85,95\}} = \gamma_{\{67,78,95,105\}} = 0,\\ \gamma_{\{68,77,98,103\}} = \gamma_{\{72,85,104\}} = \gamma_{\{76,87,96\}} = \gamma_{\{72,87\}} = 0. \end{cases} \tag{7.4.11.5}$$

Combining (7.4.11.3), (7.4.11.4) and (7.4.11.5), we get $\gamma_i = 0$ for any i. The proposition is proved. □

Remark 7.4.12. The element $[\theta]$ determined as in the proof of Proposition 7.4.8, is an $GL_4(\mathbb{F}_2)$-invariant in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{18}$.

8. The indecomposables of P_4 in degree $2^{s+t+u} + 2^{s+t} + 2^s - 3$

8.1. The τ-sequences of the admissible monomials.

In this subsection, we prove the following.

Lemma 8.1.1. *If x is an admissible monomial of degree $2^{s+t+u} + 2^{s+t} + 2^s - 3$ in P_4 then*

$$\tau(x) = (\underbrace{3; 3; \ldots; 3}_{s \text{ times}}; \underbrace{2; 2; \ldots; 2}_{t \text{ times}}; \underbrace{1; 1; \ldots; 1}_{u \text{ times}}).$$

We need the following for the proof of this lemma. From Lemmas 4.4, 2.6, Theorem 2.5 and the results in Section 7, we get

Lemma 8.1.2. *Let x be a monomial of degree $2^{s+1} + 2^s - 2$ in P_4. If x is inadmissible then there is a strictly inadmissible matrix Δ such that $\Delta \triangleright x$.*

Lemma 8.1.3. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (3,4,5,7), (3,4,7,5), (3,5,4,7), (3,5,7,4), (3,7,4,5), (3,7,5,4), (7,3,4,5), (7,3,5,4), (3,5,5,6), (3,5,6,5). We proved the lemma for the matrices associated to the monomials (3,4,5,7) and (3,5,5,6). By a direct computation, we have

$$\begin{aligned} (3,4,5,7) &= Sq^1(3,1,3,11) + Sq^2(5,2,3,7) \\ &\quad + Sq^4(3,2,3,7) \bmod L_4(3;2;3), \\ (3,5,5,6) &= Sq^1(3,3,3,9) + Sq^2(5,3,3,6) \\ &\quad + Sq^4(3,3,3,6) \bmod L_4(3;2;3). \end{aligned}$$

The lemma is proved. □

Proof of Lemma 8.1.1. Observe that $z = (2^{s+t+u}-1, 2^{s+t}-1, 2^s-1, 0)$ is the minimal spike of degree $2^{s+t+u}+2^{s+t}+2^s-3$ and

$$\tau(z) = (\underbrace{3;3;\ldots;3}_{s \text{ times}};\underbrace{2;2;\ldots;2}_{t \text{ times}};\underbrace{1;1;\ldots;1}_{u \text{ times}}).$$

Since $2^{s+t+u}+2^{s+t}+2^s-3$ is odd and x is admissible, using Proposition 2.7 and Theorem 2.13, we get $\tau_i(x) = 3$ for $1 \leqslant i \leqslant s$. Let $M = (\varepsilon_{ij}(x)), i \geqslant 1, 1 \leqslant j \leqslant 4$, be the matrix associated with x. We set $M' = (\varepsilon_{ij}(x)), i > s, 1 \leqslant j \leqslant 4$ and denote by x' the monomial corresponding to M'. We have $\tau_j(x') = \tau_{j+s}(x)$ for all $j \geqslant 1$ and

$$\begin{aligned} 2^{s+t+u}+2^{s+t}+2^s-3 = \deg x &= \sum_{i\geqslant 1} 2^{i-1}\tau_i(x) \\ &= 3(2^s-1) + 2^s\sum_{j\geqslant 1} 2^{j-1}\tau_{j+s}(x) \\ &= 3.2^s - 3 + 2^s \deg x'. \end{aligned}$$

This equality implies $\deg x' = 2^{t+u}+2^u-2$. Since x is admissible, using Lemma 8.1.2, Theorem 2.5, we see that x' is also admissible. Now, using Lemmas 7.1.1, 8.1.3, Proposition 2.7 and Theorem 2.5, we obtain

$$\tau(x') = (\underbrace{2;2;\ldots;2}_{t \text{ times}};\underbrace{1;1;\ldots;1}_{u \text{ times}}).$$

The lemma is proved. □

8.2. **The case $s = 1$ and $t = 1$.**

According to Kameko [8], $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{u+2}+3}$ is an $\mathbb{F}_2$-vector space with a basis given by the following classes:

For $u \geqslant 1$,

$$\begin{aligned} w_{u,1,1,1} &= [1,3,2^{u+2}-1], & w_{u,1,1,2} &= [1,2^{u+2}-1,3], \\ w_{u,1,1,3} &= [3,1,2^{u+2}-1], & w_{u,1,1,4} &= [3,2^{u+2}-1,1], \\ w_{u,1,1,5} &= [2^{u+2}-1,1,3], & w_{u,1,1,6} &= [2^{u+2}-1,3,1], \\ w_{u,1,1,7} &= [1,7,2^{u+2}-5], & w_{u,1,1,8} &= [7,1,2^{u+2}-5], \end{aligned}$$

For $u \geqslant 2$,

$$\begin{aligned} w_{u,1,1,9} &= [7,2^{u+2}-5,1], & w_{u,1,1,10} &= [3,3,2^{u+2}-3], \\ w_{u,1,1,11} &= [3,2^{u+2}-3,3], & w_{u,1,1,12} &= [3,5,2^{u+2}-5], \\ w_{u,1,1,13} &= [3,7,2^{u+2}-7], & w_{u,1,1,14} &= [7,3,2^{u+2}-7]. \end{aligned}$$

For $u = 2$, $w_{2,1,1,15} = [7,9,3]$.

So, we easily obtain

Proposition 8.2.1. *For any positive integer u, the dimension of the space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{u+2}+3}$ is given by*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{u+2}+3} = \begin{cases} 32, & u = 1, \\ 60, & u = 2, \\ 56, & u \geqslant 3. \end{cases}$$

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{u+2}+3}$. We have

Theorem 8.2.2. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{u+2}+3}$ is an $\mathbb{F}_2$-vector space with a basis consisting of all the classes represented by the monomials $a_{u,1,1,j}, j \geqslant 1$, which are determined as follows:*

For $u \geqslant 1$,

1. $(1,1,2,2^{u+2}-1)$, 2. $(1,1,2^{u+2}-1,2)$, 3. $(1,2,1,2^{u+2}-1)$,
4. $(1,2,2^{u+2}-1,1)$, 5. $(1,2^{u+2}-1,1,2)$, 6. $(1,2^{u+2}-1,2,1)$,
7. $(2^{u+2}-1,1,1,2)$, 8. $(2^{u+2}-1,1,2,1)$, 9. $(1,1,3,2^{u+2}-2)$,
10. $(1,1,2^{u+2}-2,3)$, 11. $(1,3,1,2^{u+2}-2)$, 12. $(1,3,2^{u+2}-2,1)$,
13. $(1,2^{u+2}-2,1,3)$, 14. $(1,2^{u+2}-2,3,1)$, 15. $(3,1,1,2^{u+2}-2)$,
16. $(3,1,2^{u+2}-2,1)$, 17. $(1,2,3,2^{u+2}-3)$, 18. $(1,2,2^{u+2}-3,3)$,
19. $(1,3,2,2^{u+2}-3)$, 20. $(1,3,2^{u+2}-3,2)$, 21. $(3,1,2,2^{u+2}-3)$,
22. $(3,1,2^{u+2}-3,2)$, 23. $(3,2^{u+2}-3,1,2)$, 24. $(3,2^{u+2}-3,2,1)$,
25. $(1,3,3,2^{u+2}-4)$, 26. $(1,3,2^{u+2}-4,3)$, 27. $(3,1,3,2^{u+2}-4)$,
28. $(3,1,2^{u+2}-4,3)$, 29. $(3,3,1,2^{u+2}-4)$, 30. $(3,3,2^{u+2}-4,1)$.

For $u = 1$, $a_{1,1,1,31} = (3,4,1,3)$, $a_{1,1,1,32} = (3,4,3,1)$.

For $u \geqslant 2$,

31. $(1,1,6,2^{u+2}-5)$, 32. $(1,6,1,2^{u+2}-5)$, 33. $(1,6,2^{u+2}-5,1)$,
34. $(1,1,7,2^{u+2}-6)$, 35. $(1,7,1,2^{u+2}-6)$, 36. $(1,7,2^{u+2}-6,1)$,
37. $(7,1,1,2^{u+2}-6)$, 38. $(7,1,2^{u+2}-6,1)$, 39. $(1,2,5,2^{u+2}-5)$,
40. $(1,3,4,2^{u+2}-5)$, 41. $(3,1,4,2^{u+2}-5)$, 42. $(3,4,1,2^{u+2}-5)$,
43. $(3,4,2^{u+2}-5,1)$, 44. $(1,2,7,2^{u+2}-7)$, 45. $(1,7,2,2^{u+2}-7)$,
46. $(7,1,2,2^{u+2}-7)$, 47. $(1,3,5,2^{u+2}-6)$, 48. $(3,1,5,2^{u+2}-6)$,
49. $(3,5,1,2^{u+2}-6)$, 50. $(3,5,2^{u+2}-6,1)$, 51. $(1,3,6,2^{u+2}-7)$,
52. $(1,6,3,2^{u+2}-7)$, 53. $(3,1,6,2^{u+2}-7)$, 54. $(1,3,7,2^{u+2}-8)$,
55. $(1,7,3,2^{u+2}-8)$, 56. $(3,1,7,2^{u+2}-8)$, 57. $(3,7,1,2^{u+2}-8)$,
58. $(7,1,3,2^{u+2}-8)$, 59. $(7,3,1,2^{u+2}-8)$, 60. $(3,5,2,2^{u+2}-7)$,
61. $(3,3,4,2^{u+2}-7)$, 62. $(3,4,3,2^{u+2}-7)$, 63. $(3,3,5,2^{u+2}-8)$,
64. $(3,5,3,2^{u+2}-8)$.

For $u = 2$,

65. $(3,12,1,3)$, 66. $(3,12,3,1)$, 67. $(1,7,9,2)$, 68. $(7,1,9,2)$,
69. $(7,9,1,2)$, 70. $(7,9,2,1)$, 71. $(1,6,9,3)$, 72. $(1,7,8,3)$,
73. $(3,7,8,1)$, 74. $(7,1,8,3)$, 75. $(7,3,8,1)$, 76. $(7,8,1,3)$,
77. $(7,8,3,1)$, 78. $(3,5,9,2)$, 79. $(3,4,9,3)$, 80. $(3,5,8,3)$.

We prove the theorem by proving some propositions.

Proposition 8.2.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{u+2}+3}$ is generated by the elements listed in Theorem 8.2.2.*

The proposition is prove by combining Theorem 2.5 and the following.

Lemma 8.2.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(3,28,1,3),(3,28,3,1),(3,7,24,1),(7,3,24,1),(3,5,25,2),(3,4,25,3),$$
$$(3,5,24,3),(3,4,9,19),(3,4,11,17),(3,5,8,19),(3,5,9,18),(3,5,10,17),$$
$$(3,5,11,16),(3,7,8,17),(7,3,8,17),(3,7,9,16),(7,3,9,16).$$

We prove the lemma for the matrices associated to the monomials

$$(3,28,1,3),(3,7,24,1),(3,5,25,2),(3,4,25,3),(3,5,24,3),$$
$$(3,4,9,19),(3,4,11,17),(3,5,8,19),(3,5,10,17),(3,7,9,16).$$

The others can be proved by a similar computation. We have

$$\begin{aligned}
(3,28,1,3) &= Sq^1(3,27,1,3) + Sq^2\big((2,27,1,3)+(5,23,2,3)\big)\\
&\quad + Sq^4\big((3,23,2,3)+(3,15,8,5)\big) + Sq^8(3,15,4,5)\\
&\quad + (2,29,1,3)+(2,27,1,5)+(3,27,1,4)+(3,25,4,3)\\
&\quad + (3,25,2,5)+(3,19,4,9) \mod \mathcal{L}_4(3;2;1;1;1),\\
(3,7,24,1) &= Sq^1(3,7,23,1) + Sq^2\big((2,7,23,1)+(5,7,19,2)\big)\\
&\quad + Sq^4\big((3,4,23,1)+(3,11,15,2)+(3,5,21,2)+(3,5,19,4)\\
&\quad + (3,11,13,4)\big) + Sq^8\big((3,7,15,2)+(3,7,13,4)\big)\\
&\quad + (2,9,23,1)+(2,7,25,1)+(3,4,27,1)+(3,5,25,2)\\
&\quad + (3,5,19,8)+(3,7,17,8) \mod \mathcal{L}_4(3;2;1;1;1),\\
(3,5,25,2) &= Sq^1(3,3,27,1) + Sq^2\big((5,3,23,2)+(2,3,27,1)\big)\\
&\quad + Sq^4\big((3,3,23,2)+(3,9,15,4)\big) + Sq^8(3,5,15,4)\\
&\quad + (3,3,25,4)+(3,3,28,1)+(3,4,27,1)+(2,5,27,1)\\
&\quad + (2,3,29,1)+(3,5,19,8) \mod \mathcal{L}_4(3;2;1;1;1),\\
(3,4,25,3) &= Sq^1(3,1,27,3) + Sq^2\big((5,2,23,3)+(2,1,27,3)\big)\\
&\quad + Sq^4\big((3,2,23,3)+(3,8,15,5)\big) + Sq^8(3,4,15,5)\\
&\quad + (3,2,25,5)+(3,1,28,3)+(3,1,27,4)+(2,1,27,5)\\
&\quad + (2,1,29,3)+(3,4,19,9) \mod \mathcal{L}_4(3;2;1;1;1),\\
(3,5,24,3) &= Sq^1(3,3,25,3) + Sq^2\big((5,3,22,3)+(2,3,25,3)\big)\\
&\quad + Sq^4\big((3,3,22,3)+(3,9,14,5)\big) + Sq^8(3,5,14,5)\\
&\quad + (3,3,24,5)+(3,3,25,4)+(3,4,25,3)+(2,5,25,3)\\
&\quad + (2,3,25,5)+(3,5,18,9) \mod \mathcal{L}_4(3;2;1;1;1),\\
(3,4,9,19) &= Sq^1(3,1,3,27) + Sq^2\big((5,2,3,23)+(2,1,3,27)\big)\\
&\quad + Sq^4\big((3,8,5,15)+(3,2,3,23)\big) + Sq^8(3,4,5,15)\\
&\quad + (3,4,3,25)+(3,2,5,25)+(3,1,3,28)+(3,1,4,27)\\
&\quad + (2,1,5,27)+(2,1,3,29) \mod \mathcal{L}_4(3;2;1;1;1),\\
(3,4,11,17) &= Sq^1(3,1,11,19) + Sq^2\big((5,2,7,19)+(2,1,11,19)\big)\\
&\quad + Sq^4\big((3,8,7,13)+(3,2,7,19)\big) + Sq^8(3,4,7,13)\\
&\quad + (3,4,9,19)+(3,2,9,21)+(3,1,12,19)+(3,1,11,20)\\
&\quad + (2,1,13,19)+(2,1,11,21) \mod \mathcal{L}_4(3;2;1;1;1),\\
(3,5,8,19) &= Sq^1(3,3,1,27) + Sq^2\big((5,3,2,23)+(2,3,1,27)\big)\\
&\quad + Sq^4\big((3,9,4,15)+(3,3,2,23)\big) + Sq^8(3,5,4,15)\\
&\quad + (3,5,2,25)+(3,3,4,25)+(3,3,1,28)+(3,4,1,27)\\
&\quad + (2,5,1,27)+(2,3,1,29) \mod \mathcal{L}_4(3;2;1;1;1),
\end{aligned}$$

$$\begin{aligned}(3,5,9,18) &= Sq^1(3,3,3,25) + Sq^2\big((5,3,3,22) + (2,3,3,25)\big)\\ &\quad + Sq^4\big((3,9,5,14) + (3,3,3,22)\big) + Sq^8(3,5,5,14)\\ &\quad + (3,5,3,24) + (3,3,5,24) + (3,4,3,25) + (3,3,4,25)\\ &\quad + (2,5,3,25) + (2,3,5,25) \mod \mathcal{L}_4(3;2;1;1;1),\\ (3,5,10,17) &= Sq^1(3,3,9,19) + Sq^2\big((5,3,6,19) + (2,3,9,19)\big)\\ &\quad + Sq^4\big((3,9,6,13) + (3,3,6,19)\big) + Sq^8(3,5,6,13)\\ &\quad + (3,3,8,21) + (3,5,8,19) + (3,3,9,20) + (3,4,9,19)\\ &\quad + (2,5,9,19) + (2,3,9,21) \mod \mathcal{L}_4(3;2;1;1;1),\\ (3,7,8,17) &= Sq^1(3,7,1,23) + Sq^2\big((5,7,2,19) + (2,7,1,23)\big)\\ &\quad + Sq^4\big((3,11,4,13) + (3,11,2,15) + (3,5,4,19) + (3,5,2,21)\\ &\quad + (3,4,1,23)\big) + Sq^8\big((3,7,4,13) + (3,7,2,15)\big)\\ &\quad + (3,5,8,19) + (3,5,2,25) + (3,7,1,24) + (2,9,1,23)\\ &\quad + (2,7,1,25) + (3,4,1,27) \mod \mathcal{L}_4(3;2;1;1;1),\\ (3,7,9,16) &= Sq^1\big((3,7,3,21) + (1,7,5,21) + (3,7,1,23)\big)\\ &\quad + Sq^2\big((5,7,3,18) + (2,7,3,21) + (1,7,6,19) + (1,7,3,22)\\ &\quad + (5,7,2,19) + (2,7,1,23)\big) + Sq^4\big((3,11,5,12) + (3,11,3,14)\\ &\quad + (3,5,5,18) + (3,5,3,20) + (3,4,3,21) + (3,11,4,13)\big)\\ &\quad + Sq^8\big((3,7,5,12) + (3,7,3,14) + (3,7,4,13)\big) + (3,5,9,18)\\ &\quad + (3,5,3,24) + (3,4,3,25) + (2,9,3,21) + (1,9,3,22) + (1,7,3,24)\\ &\quad + (1,7,8,19) + (1,9,6,19) + (3,7,8,17) \mod \mathcal{L}_4(3;2;1;1;1).\end{aligned}$$

The lemma is proved. □

Combining Lemmas 5.10, 8.1.3, 8.1.1, 8.2.4, Theorem 2.5 and the results in Section 7, we get Proposition 8.2.3.

Now, we show that the elements listed in Theorem 8.2.2 are linearly independent.

Proposition 8.2.5. *The elements* $[a_{1,1,1,j}], 1 \leqslant j \leqslant 32$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{11}$.

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{32} \gamma_j [a_{1,1,1,j}] = 0, \tag{8.2.5.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms f_1, f_2, f_3 to the relation (8.2.5.1), we obtain

$$\begin{aligned}
&\gamma_3[3,1,7] + \gamma_4[3,7,1] + \gamma_{\{13,31\}}[7,1,3]\\
&\quad + \gamma_{\{14,32\}}[7,3,1] + \gamma_{17}[3,3,5] + \gamma_{18}[3,5,3] = 0,\\
&\gamma_1[3,1,7] + \gamma_6[3,7,1] + \gamma_{\{10,28\}}[7,1,3]\\
&\quad + \gamma_{\{12,30\}}[7,3,1] + \gamma_{\{19,26\}}[3,3,5] + \gamma_{26}[3,5,3] = 0,\\
&\gamma_2[3,1,7] + \gamma_5[3,7,1] + \gamma_{\{9,27\}}[7,1,3]\\
&\quad + \gamma_{\{11,29\}}[7,3,1] + \gamma_{\{20,25\}}[3,3,5] + \gamma_{25}[3,5,3] = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\begin{cases}
\gamma_i = 0,\ i = 1,2,3,4,5,6,17,18,19,20,25,26,\\
\gamma_{\{13,31\}} = \gamma_{\{14,32\}} = \gamma_{\{10,28\}} = 0,\\
\gamma_{\{12,30\}} = \gamma_{\{9,27\}} = \gamma_{\{11,29\}} = 0.
\end{cases} \tag{8.2.5.2}$$

With the aid of (8.2.5.2), the homomorphisms f_4, f_5, f_6 send (8.2.5.1) to

$$\begin{aligned}
&\gamma_{\{10,13\}}[1,7,3] + \gamma_{\{16,24,30,32\}}[3,7,1]\\
&\quad + \gamma_8[7,3,1] + \gamma_{21}[3,3,5] + \gamma_{\{28,31\}}[3,5,3] = 0,\\
&\gamma_{\{9,14\}}[1,7,3] + \gamma_{\{15,23,29,31\}}[3,7,1]\\
&\quad + \gamma_7[7,3,1] + \gamma_{22}[3,3,5] + \gamma_{\{27,32\}}[3,5,3] = 0,\\
&\gamma_{\{11,12\}}[1,3,7] + \gamma_{\{15,16,21,22,27,28\}}[3,1,7]\\
&\quad + \gamma_{\{7,8\}}[7,1,3] + \gamma_{\{29,30\}}[3,3,5] + \gamma_{\{23,24\}}[3,5,3] = 0.
\end{aligned}$$

From these equalities, we get

$$\begin{cases}
\gamma_j = 0,\ i = 7,8,21,22,\\
\gamma_{\{10,13\}} = \gamma_{\{16,24,30,32\}} = \gamma_{\{28,31\}} = \gamma_{\{9,14\}} = 0,\\
\gamma_{\{15,23,29,31\}} = \gamma_{\{27,32\}} = \gamma_{\{11,12\}} = 0,\\
\gamma_{\{15,16,27,28\}} = \gamma_{\{29,30\}} = \gamma_{\{23,24\}} = 0.
\end{cases} \tag{8.2.5.3}$$

With the aid of (8.2.5.2) and (8.2.5.3), the homomorphisms g_1, g_2 send (8.2.5.1) to

$$\begin{aligned}
&\gamma_{16}[3,7,1] + \gamma_{23}[7,3,1] + \gamma_{10}[3,3,5] = 0,\\
&\gamma_{15}[3,7,1] + \gamma_{23}[7,3,1] + \gamma_9[3,3,5] = 0.
\end{aligned}$$

These equalities imply

$$\gamma_j = 0, j = 9,10,15,16,23. \tag{8.2.5.4}$$

Combining (8.2.5.2), (8.2.5.3) and (8.2.5.4), we get $\gamma_j = 0$ for $1 \leqslant j \leqslant 32$. The proposition is proved. □

Proposition 8.2.6. *The elements* $[a_{2,1,1,j}], 1 \leqslant j \leqslant 80$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{19}$.

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{80} \gamma_j [a_{2,1,1,j}] = 0, \tag{8.2.6.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms f_1, f_2, f_3 to the relation (8.2.6.1), we obtain

$$\begin{aligned}
&\gamma_3[3,1,15] + \gamma_4[3,15,1] + \gamma_{\{13,65,76\}}[15,1,3] + \gamma_{\{14,66,77\}}[15,3,1]\\
&\quad + \gamma_{17}[3,3,13] + \gamma_{18}[3,13,3] + \gamma_{\{32,42\}}[7,1,11] + \gamma_{\{33,43\}}[7,11,1]\\
&\quad + \gamma_{39}[3,5,11] + \gamma_{44}[3,7,9] + \gamma_{\{52,62\}}[7,3,9] + \gamma_{\{71,79\}}[7,9,3] = 0,\\
&\gamma_1[3,1,15] + \gamma_6[3,15,1] + \gamma_{\{10,28,74\}}[15,1,3] + \gamma_{\{12,30,75\}}[15,3,1]\\
&\quad + \gamma_{\{19,40\}}[3,3,13] + \gamma_{72}[3,13,3] + \gamma_{\{31,41\}}[7,1,11] + \gamma_{\{36,73\}}[7,11,1]\\
&\quad + \gamma_{40}[3,5,11] + \gamma_{\{45,72\}}[3,7,9] + \gamma_{\{26,51,61\}}[7,3,9] + \gamma_{\{26,80\}}[7,9,3] = 0,\\
&\gamma_2[3,1,15] + \gamma_5[3,15,1] + \gamma_{\{9,27,58\}}[15,1,3] + \gamma_{\{11,29,59\}}[15,3,1]\\
&\quad + \gamma_{\{20,54\}}[3,3,13] + \gamma_{55}[3,13,3] + \gamma_{\{34,56\}}[7,1,11] + \gamma_{\{35,57\}}[7,11,1]\\
&\quad + \gamma_{54}[3,5,11] + \gamma_{\{55,67\}}[3,7,9] + \gamma_{\{25,47,63\}}[7,3,9] + \gamma_{\{25,64\}}[7,9,3] = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\begin{cases}
\gamma_i = 0,\ i = 1,2,3,4,5,6,17,18,\\
19,20,39,40,44,45,54,55,67,72,\\
\gamma_{\{13,65,76\}} = \gamma_{\{14,66,77\}} = \gamma_{\{32,42\}} = \gamma_{\{33,43\}} = \gamma_{\{52,62\}} = 0,\\
\gamma_{\{71,79\}} = \gamma_{\{10,28,74\}} = \gamma_{\{12,30,75\}} = \gamma_{\{31,41\}} = \gamma_{\{36,73\}} = 0,\\
\gamma_{\{26,51,61\}} = \gamma_{\{26,80\}} = \gamma_{\{9,27,58\}} = \gamma_{\{11,29,59\}} = 0,\\
\gamma_{\{34,56\}} = \gamma_{\{35,57\}} = \gamma_{\{25,47,63\}} = \gamma_{\{25,64\}} = 0.
\end{cases} \tag{8.2.6.2}$$

With the aid of (8.2.6.2), the homomorphisms f_4, f_5, f_6 send (8.2.6.1) to

$$\begin{aligned}
&\gamma_{\{10,13,26,71\}}[1,15,3] + a_1[3,15,1] + \gamma_8[15,3,1] + \gamma_{21}[3,3,13]\\
&\quad + a_2[3,13,3] + \gamma_{\{31,32\}}[1,7,11] + \gamma_{\{38,70,75,77\}}[7,11,1] + \gamma_{\{41,42\}}[3,5,11]\\
&\quad + \gamma_{\{53,60,61,62\}}[3,7,9] + \gamma_{46}[7,3,9] + \gamma_{\{74,76\}}[7,9,3] = 0,\\
&\gamma_{\{9,14,25,52\}}[1,15,3] + a_3[3,15,1] + \gamma_7[15,3,1] + \gamma_{22}[3,3,13] + \gamma_{68}[7,3,9]\\
&\quad + \gamma_{\{27,62,64,66\}}[3,13,3] + \gamma_{\{33,34\}}[1,7,11] + \gamma_{\{37,59,69,76\}}[7,11,1]\\
&\quad + \gamma_{\{43,56\}}[3,5,11] + \gamma_{\{48,63,78,79\}}[3,7,9] + \gamma_{\{58,77\}}[7,9,3] = 0,\\
&a_4[1,3,15] + a_5[3,1,15] + \gamma_{\{7,8\}}[15,1,3] + \gamma_{\{29,30,61,63\}}[3,3,13]\\
&\quad + \gamma_{\{23,24\}}[3,13,3] + \gamma_{\{35,36\}}[1,7,11] + a_6[7,1,11] + a_7[3,5,11]\\
&\quad + \gamma_{\{57,73\}}[3,7,9] + \gamma_{\{59,75\}}[7,3,9] + \gamma_{\{69,70\}}[7,9,3] = 0.
\end{aligned}$$

where

$$a_1 = \gamma_{\{16,24,30,43,50,66,73\}},\quad a_2 = \gamma_{\{28,65,79,80\}},\quad a_3 = \gamma_{\{15,23,29,42,49,57,65\}},$$
$$a_4 = \gamma_{\{11,12,25,26,47,51\}},\quad a_6 = \gamma_{\{37,38,46,58,68,74\}},\quad a_7 = \gamma_{\{49,50,60,64,78,80\}},$$
$$a_5 = \gamma_{\{15,16,21,22,27,28,41,48,53,56\}}.$$

From these equalities, we get

$$\text{(8.2.6.3)}\qquad \begin{cases} a_i = 0,\ i = 1,2,3,4,5,6,7, \\ \gamma_j = 0,\ j = 7,8,21,22,46,68, \\ \gamma_{\{10,13,26,71\}} = \gamma_{\{31,32\}} = \gamma_{\{38,58,59,69\}} = \gamma_{\{52,53,60,61\}} = 0, \\ \gamma_{\{74,76\}} = \gamma_{\{9,14,25,52\}} = \gamma_{\{25,27,52,66\}} = \gamma_{\{33,34\}} = 0, \\ \gamma_{\{37,59,69,74\}} = \gamma_{\{48,63,71,78\}} = \gamma_{\{58,77\}} = \gamma_{\{23,24\}} = 0, \\ \gamma_{\{29,30,61,63\}} = \gamma_{\{35,36\}} = \gamma_{\{59,75\}} = \gamma_{\{69,70\}} = 0. \end{cases}$$

With the aid of (8.2.6.2) and (8.2.6.3), the homomorphisms g_1, g_2 send (8.2.6.1) respectively to

$$\begin{aligned} &\gamma_{\{10,28,74\}}[1,15,3] + a_8[3,15,1] + \gamma_{\{13,65,74\}}[15,1,3] + a_9[15,3,1] \\ &\quad + \gamma_{31}[3,3,13] + \gamma_{\{28,71,74\}}[3,13,3] + a_{10}[7,11,1] \\ &\quad + \gamma_{\{26,51,53,61,71\}}[3,7,9] + \gamma_{\{60,65,74\}}[7,3,9] + \gamma_{\{26,65,74\}}[7,9,3] = 0, \\ &\gamma_{\{9,27,58\}}[1,15,3] + a_{11}[3,15,1] + \gamma_{\{14,58,66\}}[15,1,3] \\ &\quad + a_{12}[15,3,1] + \gamma_{33}[3,3,13] + \gamma_{\{27,52,58\}}[3,13,3] + a_{13}[7,11,1] \\ &\quad + a_{14}[3,7,9] + \gamma_{\{58,66,78\}}[7,3,9] + \gamma_{\{25,58,66\}}[7,9,3] = 0. \end{aligned}$$

where

$$a_8 = \gamma_{\{12,16,30,38,59\}},\quad a_9 = \gamma_{\{14,23,58,66,69\}},\quad a_{10} = \gamma_{\{38,50,58,59,69\}},$$
$$a_{11} = \gamma_{\{11,15,29,37,59\}},\quad a_{12} = \gamma_{\{13,23,65,69,74\}},\quad a_{13} = \gamma_{\{37,49,59,69,74\}},$$
$$a_{14} = \gamma_{\{25,47,48,52,63\}}$$

From these equalities, we obtain

$$\text{(8.2.6.4)}\qquad \begin{cases} \gamma_{31} = \gamma_{33} = a_j = 0,\ j = 8,9,\ldots,14, \\ \gamma_{\{10,28,74\}} = \gamma_{\{13,65,74\}} = \gamma_{\{28,71,74\}} = \gamma_{\{60,65,74\}} = 0, \\ \gamma_{\{26,51,53,61,71\}} = \gamma_{\{26,65,74\}} = \gamma_{\{9,27,58\}} = 0, \\ \gamma_{\{14,58,66\}} = \gamma_{\{27,52,58\}} = \gamma_{\{58,66,78\}} = \gamma_{\{25,58,66\}} = 0. \end{cases}$$

With the aid of (8.2.6.2), (8.2.6.3) and (8.2.6.4), the homomorphisms g_3, g_4 send (8.2.6.1) to

$$\begin{aligned} &\gamma_{\{11,29,59\}}[1,15,3] + a_{15}[3,15,1] + \gamma_{\{12,30,59\}}[15,1,3] \\ &\quad + a_{16}[15,3,1] + \gamma_{35}[3,3,13] + a_{17}[3,13,3] + a_{18}[7,11,1] \\ &\quad + a_{19}[3,7,9] + \gamma_{\{30,50,59,78\}}[7,3,9] + a_{20}[7,9,3] = 0, \end{aligned}$$

$$\begin{aligned}a_{21}[1,15,3]+a_{22}[3,15,1]+a_{23}[15,1,3]+a_{24}[15,3,1]+\gamma_{\{59,69\}}[3,3,13]\\ +a_{25}[3,13,3]+\gamma_{\{37,59,69,74\}}[1,7,11]+\gamma_{\{38,58,59,69\}}[7,1,11]\\ +\gamma_{\{47,51,52,71\}}[7,11,1]+a_{26}[3,7,9]+a_{27}[7,3,9]+a_{28}[7,9,3]=0.\end{aligned}$$

where

$$\begin{aligned}&a_{15}=\gamma_{\{9,15,27,37,58\}},\quad a_{16}=\gamma_{\{10,16,28,38,74\}},\quad a_{17}=\gamma_{\{25,29,47,59,61,63\}},\\ &a_{18}=\gamma_{\{37,38,48,53,58,74\}},\quad a_{19}=\gamma_{\{25,47,49,60,61,63\}},\quad a_{20}=\gamma_{\{26,30,51,59,61,63\}},\\ &a_{21}=\gamma_{\{15,23,29,35,49,65\}},\quad a_{22}=\gamma_{\{9,11,13,14,25,35,52\}},\quad a_{23}=\gamma_{\{16,23,30,35,50,66\}},\\ &a_{24}=\gamma_{\{10,12,13,14,26,35,71\}},\quad a_{25}=\gamma_{\{23,25,26,27,29,35,48,49,52,60,61,63,66,71,78\}},\\ &a_{26}=\gamma_{\{26,48,49,61,63,71,78\}},\quad a_{27}=\gamma_{\{23,26,28,30,35,65,71,78\}},\\ &a_{28}=\gamma_{\{23,25,26,28,30,35,50,52,53,60,61,63,65,71,78\}}.\end{aligned}$$

So, we obtain

$$\begin{cases}\gamma_{35}=a_j=0,\ j=15,16,\ldots,28,\\ \gamma_{\{11,29,59\}}=\gamma_{\{12,30,59\}}=\gamma_{\{30,50,59,78\}}=\gamma_{\{59,69\}}=0,\\ \gamma_{\{37,59,69,74\}}=\gamma_{\{38,58,59,69\}}=\gamma_{\{47,51,52,71\}}=0.\end{cases}\tag{8.2.6.5}$$

Substituting (8.2.6.2), (8.2.6.3), (8.2.6.4), and (8.2.6.5) into the relation (8.2.6.1), we have

$$\gamma_{15}[\theta_1]+\gamma_{16}[\theta_2]+\gamma_{12}[\theta_3]+\gamma_{11}[\theta_4]+\gamma_9[\theta_5]+\gamma_{23}[\theta_6]=0,\tag{8.2.6.6}$$

where

$$\begin{aligned}\theta_1&=a_{2,1,1,15}+a_{2,1,1,28}+a_{2,1,1,37}+a_{2,1,1,65}+a_{2,1,1,74}+a_{2,1,1,76},\\ \theta_2&=a_{2,1,1,16}+a_{2,1,1,27}+a_{2,1,1,38}+a_{2,1,1,58}+a_{2,1,1,66}+a_{2,1,1,77},\\ \theta_3&=a_{2,1,1,12}+a_{2,1,1,14}+a_{2,1,1,25}+a_{2,1,1,30}\\ &\quad+a_{2,1,1,63}+a_{2,1,1,64}+a_{2,1,1,66}+a_{2,1,1,78},\\ \theta_4&=a_{2,1,1,11}+a_{2,1,1,13}+a_{2,1,1,26}+a_{2,1,1,29}\\ &\quad+a_{2,1,1,60}+a_{2,1,1,61}+a_{2,1,1,65}+a_{2,1,1,80},\\ \theta_5&=a_{2,1,1,9}+a_{2,1,1,10}+a_{2,1,1,27}+a_{2,1,1,28}+a_{2,1,1,48}\\ &\quad+a_{2,1,1,52}+a_{2,1,1,53}+a_{2,1,1,62}+a_{2,1,1,71}+a_{2,1,1,79},\\ \theta_6&=a_{2,1,1,23}+a_{2,1,1,24}+a_{2,1,1,29}+a_{2,1,1,30}\\ &\quad+a_{2,1,1,59}+a_{2,1,1,69}+a_{2,1,1,70}+a_{2,1,1,75}.\end{aligned}$$

We need to prove $\gamma_9=\gamma_{11}=\gamma_{12}=\gamma_{15}=\gamma_{16}=\gamma_{23}=0$. The proof is divided into 4 steps.

Step 1. First we prove $\gamma_{23}=0$ by showing the polynomial $\theta=\beta_1\theta_1+\beta_2\theta_2+\beta_3\theta_3+\beta_4\theta_4+\beta_5\theta_5+\theta_6$ is non-hit for all $\beta_1,\beta_2,\beta_3,\beta_4,\beta_5\in\mathbb{F}_2$. Suppose the contrary that this polynomial is hit. Then we have

$$\theta=Sq^1(A)+Sq^2(B)+Sq^4(C)+Sq^8(D),$$

for some polynomials A, B, C, D in R_4. Let $(Sq^2)^3$ act on the both sides of this equality. Using the relations $(Sq^2)^3Sq^1 = 0, (Sq^2)^3Sq^2 = 0$, we get

$$(Sq^2)^3(\theta) = (Sq^2)^3Sq^4(C) + (Sq^2)^3Sq^8(D).$$

The polynomial $(7, 12, 4, 2) + (7, 12, 2, 4)$ is a summand of $(Sq^2)^3(\theta)$. This polynomial is not a summand of $(Sq^2)^3Sq^8(D)$ for all D. It is also not a summand of $(Sq^2)^3Sq^4(C)$ if either $(7, 5, 1, 2) + (7, 5, 2, 1)$ or $(7, 6, 1, 1)$ is not a summand of C.

Suppose $(7, 5, 1, 2) + (7, 5, 2, 1)$ is a summand of C. Then

$$(Sq^2)^3(\theta + Sq^4((7, 5, 1, 2) + (7, 5, 2, 1))) = (Sq^2)^3(Sq^4(C') + Sq^8(D)),$$

where $C' = C + (7, 5, 1, 2) + (7, 5, 2, 1)$. We see that the polynomial $(16, 6, 1, 2) + (16, 6, 2, 1)$ is a summand of $(Sq^2)^3(\theta + Sq^4((7, 5, 1, 2) + (7, 5, 2, 1)))$. This polynomial is not a summand of $(Sq^2)^3Sq^8(D)$ for all D. It is also not a summand of $(Sq^2)^3Sq^4(C')$ for $C' \neq (7, 5, 1, 2) + (7, 5, 2, 1), (7, 6, 1, 1)$. So, either $(7, 5, 1, 2) + (7, 5, 2, 1)$ or $(7, 6, 1, 1)$ is a summand of C'. Since $(7, 5, 1, 2) + (7, 5, 2, 1)$ is not a summand of C', $(7, 6, 1, 1)$ is a summand of C'. Hence, we obtain

$$\begin{aligned}(Sq^2)^3(\theta + Sq^4((7, 5, 1, 2) + (7, 5, 2, 1) + (7, 6, 1, 1)))&\\ = (Sq^2)^3(Sq^4(C'') + Sq^8(D)),&\end{aligned}$$

where $C'' = C + (7, 5, 1, 2) + (7, 5, 2, 1) + (7, 6, 1, 1)$. Now $(7, 12, 2, 4) + (7, 12, 4, 2)$ is a summand of

$$(Sq^2)^3(\theta + Sq^4((7, 5, 1, 2) + (7, 5, 2, 1) + (7, 6, 1, 1))).$$

So, either $(7, 5, 1, 2) + (7, 5, 2, 1)$ or $(7, 6, 1, 1)$ is a summand of C'' and we have a contradiction. Hence, $[\theta] \neq 0$ and $\gamma_{23} = 0$.

Step 2. By a direct computation, we see that the homomorphism φ_2 sends (8.2.6.6) to $\gamma_{15}[\theta_1] + \gamma_{16}[\theta_6] + \gamma_{12}[\theta_3] + \gamma_{11}[\theta_5] + \gamma_9[\theta_4] = 0$. By Step 1, we obtain $\gamma_{16} = 0$.

Step 3. The homomorphism $\varphi_2\varphi_1$ sends (8.2.6.6) to $\gamma_{15}[\theta_5] + \gamma_{12}[\theta_6] + \gamma_{11}[\theta_1] + \gamma_9[\theta_4] = 0$. By Step 1, we obtain $\gamma_{12} = 0$.

Step 4. Now the homomorphism φ_3 sends (8.2.6.6) to $\gamma_{15}[\theta_2] + \gamma_{11}[\theta_3] + \gamma_9[\theta_5] = 0$. Combining Step 2 and Step 3, we obtain $\gamma_{11} = \gamma_{15} = 0$.

Since $\varphi_2([\theta_5]) = [\theta_2]$, we get $\gamma_9 = 0$. So, we obtain $\gamma_j = 0$ for all j. The proposition follows. □

Proposition 8.2.7. *For $u \geqslant 3$, the elements $[a_{u,1,1,j}], 1 \leqslant j \leqslant 64$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{u+2}+3}$.*

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{64} \gamma_j[a_{u,1,1,j}] = 0, \tag{8.2.7.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms f_1, f_2, f_3 to the relation (8.2.7.1), we obtain

$$\begin{aligned}
&\gamma_3 w_{u,1,1,3} + \gamma_4 w_{u,1,1,4} + \gamma_{13} w_{u,1,1,5} + \gamma_{14} w_{u,1,1,6}\\
&\quad + \gamma_{\{32,42\}} w_{u,1,1,8} + \gamma_{\{33,43\}} w_{u,1,1,9} + \gamma_{17} w_{u,1,1,10} + \gamma_{18} w_{u,1,1,11}\\
&\quad + \gamma_{39} w_{u,1,1,12} + \gamma_{44} w_{u,1,1,13} + \gamma_{\{52,62\}} w_{u,1,1,14} = 0,\\
&\gamma_1 w_{u,1,1,3} + \gamma_6 w_{u,1,1,4} + \gamma_{\{10,28\}} w_{u,1,1,5} + \gamma_{\{12,30\}} w_{u,1,1,6}\\
&\quad + \gamma_{\{31,41\}} w_{u,1,1,8} + \gamma_{36} w_{u,1,1,9} + \gamma_{\{19,26,40\}} w_{u,1,1,10}\\
&\quad + \gamma_{26} w_{u,1,1,11} + \gamma_{40} w_{u,1,1,12} + \gamma_{45} w_{u,1,1,13} + \gamma_{\{51,61\}} w_{u,1,1,14} = 0,\\
&\gamma_2 w_{u,1,1,3} + \gamma_5 w_{u,1,1,4} + \gamma_{\{9,27,58\}} w_{u,1,1,5} + \gamma_{\{11,29,59\}} w_{u,1,1,6}\\
&\quad + \gamma_{\{34,56\}} w_{u,1,1,8} + \gamma_{\{35,57\}} w_{u,1,1,9} + \gamma_{\{20,25,54,64\}} w_{u,1,1,10}\\
&\quad + \gamma_{\{25,55,64\}} w_{u,1,1,11} + \gamma_{54} w_{u,1,1,12} + \gamma_{55} w_{u,1,1,13} + \gamma_{\{47,63,64\}} w_{u,1,1,14} = 0.
\end{aligned}$$

Computing from these equalities, we get

$$\text{(8.2.7.2)}\quad \begin{cases}
\gamma_j = 0,\ j = 1,2,3,4,5,6,13,14,17,18,\\
\qquad\qquad 19,26,36,39,40,44,45,54,55,\\
\gamma_{\{32,42\}} = \gamma_{\{33,43\}} = \gamma_{\{52,62\}} = \gamma_{\{10,28\}} = \gamma_{\{12,30\}} = 0,\\
\gamma_{\{31,41\}} = \gamma_{\{51,61\}} = \gamma_{\{9,27,58\}} = \gamma_{\{11,29,59\}} = 0,\\
\gamma_{\{34,56\}} = \gamma_{\{35,57\}} = \gamma_{\{20,25,64\}} = \gamma_{\{25,64\}} = \gamma_{\{47,63,64\}} = 0.
\end{cases}$$

With the aid of (8.2.7.2), the homomorphisms f_4, f_5, f_6 send (8.2.7.1) to

$$\begin{aligned}
&\gamma_{10} w_{u,1,1,2} + \gamma_{\{16,24,30,43,50\}} w_{u,1,1,4} + \gamma_8 w_{u,1,1,6}\\
&\quad + \gamma_{\{31,32\}} w_{u,1,1,7} + \gamma_{38} w_{u,1,1,9} + \gamma_{21} w_{u,1,1,10} + \gamma_{28} w_{u,1,1,11}\\
&\quad + \gamma_{\{41,42\}} w_{u,1,1,12} + \gamma_{\{53,60,61,62\}} w_{u,1,1,13} + \gamma_{46} w_{u,1,1,14} = 0,\\
&\gamma_{\{9,25,52\}} w_{u,1,1,2} + \gamma_{\{15,23,29,42,49,57\}} w_{u,1,1,4} + \gamma_7 w_{u,1,1,6} + \gamma_{\{33,34\}} w_{u,1,1,7}\\
&\quad + \gamma_{\{37,59\}} w_{u,1,1,9} + \gamma_{\{22,58\}} w_{u,1,1,10} + \gamma_{\{27,58,62,64\}} w_{u,1,1,11}\\
&\quad + \gamma_{\{43,56\}} w_{u,1,1,12} + \gamma_{\{48,63\}} w_{u,1,1,13} + \gamma_{58} w_{u,1,1,14} = 0,\\
&\gamma_{\{11,12,20,25,47,51\}} w_{u,1,1,1} + \gamma_{\{15,16,21,22,27,28,41,48,53,56\}} w_{u,1,1,3} + \gamma_{\{7,8\}} w_{u,1,1,5}\\
&\quad + \gamma_{35} w_{u,1,1,7} + \gamma_{\{37,38,46,58\}} w_{u,1,1,8} + \gamma_{\{29,30,61,63\}} w_{u,1,1,10}\\
&\quad + \gamma_{\{23,24\}} w_{u,1,1,11} + \gamma_{\{49,50,60,64\}} w_{u,1,1,12} + \gamma_{57} w_{u,1,1,13} + \gamma_{59} w_{u,1,1,14} = 0.
\end{aligned}$$

Computing from these equalities and (8.2.7.2), we get

$$\text{(8.2.7.3)}\quad \begin{cases}
\gamma_j = 0,\ j = 7,8,10,20,21,22,28,35,37,38,46,57,58,59,\\
\gamma_{\{16,24,30,43,50\}} = \gamma_{\{31,32\}} = \gamma_{\{41,42\}} = \gamma_{\{53,60,61,62\}} = 0,\\
\gamma_{\{9,25,52\}} = \gamma_{\{15,23,29,42,49\}} = \gamma_{\{33,34\}} = \gamma_{\{27,62,64\}} = 0,\\
\gamma_{\{43,56\}} = \gamma_{\{48,63\}} = \gamma_{\{29,30,61,63\}} = \gamma_{\{49,50,60,64\}} = 0,\\
\gamma_{\{23,24\}} = \gamma_{\{11,12,20,25,47,51\}} = \gamma_{\{15,16,21,22,27,28,41,48,53,56\}} = 0.
\end{cases}$$

With the aid of (8.2.7.2) and (8.2.7.3), the homomorphisms g_1, g_2, g_3 send (8.2.7.1) to

$$\begin{aligned}
&\gamma_{16}w_{u,1,1,4} + \gamma_{23}w_{u,1,1,6} + \gamma_{50}w_{u,1,1,9}\\
&\quad + \gamma_{31}w_{u,1,1,10} + \gamma_{53}w_{u,1,1,13} + \gamma_{60}w_{u,1,1,14} = 0,\\
&\gamma_{15}w_{u,1,1,4} + \gamma_{23}w_{u,1,1,6} + \gamma_{49}w_{u,1,1,9} + \gamma_{\{25,33\}}w_{u,1,1,10}\\
&\quad + \gamma_{\{9,25,52\}}w_{u,1,1,11} + \gamma_{\{25,47,52\}}w_{u,1,1,13} + \gamma_{25}w_{u,1,1,14} = 0,\\
&\gamma_{15}w_{u,1,1,4} + \gamma_{16}w_{u,1,1,6} + \gamma_{\{48,53\}}w_{u,1,1,9} + \gamma_{\{11,12,25,47,51\}}w_{u,1,1,11}\\
&\quad + \gamma_{\{12,48\}}w_{u,1,1,10} + \gamma_{\{25,47,48,49,51,60\}}w_{u,1,1,13} + \gamma_{\{48,50\}}w_{u,1,1,14} = 0.
\end{aligned}$$

Computing directly from these equalities, we get

$$\text{(8.2.7.4)}\qquad \begin{cases} \gamma_j = 0,\ j = 12, 15, 16, 23, 25, 31, 33, 48, 49, 50, 53, 60,\\ \gamma_{\{9,52\}} = \gamma_{\{47,52\}} = \gamma_{\{11,47,51\}} = \gamma_{\{47,51\}} = 0. \end{cases}$$

The relations (8.2.7.2), (8.2.7.3) and (8.2.7.4) imply $\gamma_j = 0$ for $1 \leqslant j \leqslant 64$. The proposition is proved. □

8.3. **The case $s = 1, t \geqslant 2$ and $u = 1$.**

It is well known that for $t \geqslant 2$, $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{t+2}+2^{t+1}-1} = 14$ with a basis given by the classes $w_{1,t,1,j}, 1 \leqslant j \leqslant 14$, which are determined as follows:

For $t \geqslant 2$,

1. $[1, 2^{t+1}-1, 2^{t+2}-1]$, 2. $[1, 2^{t+2}-1, 2^{t+1}-1]$,
3. $[2^{t+1}-1, 1, 2^{t+2}-1]$, 4. $[2^{t+1}-1, 2^{t+2}-1, 1]$,
5. $[2^{t+2}-1, 1, 2^{t+1}-1]$, 6. $[2^{t+2}-1, 2^{t+1}-1, 1]$,
7. $[3, 2^{t+1}-3, 2^{t+2}-1]$, 8. $[3, 2^{t+2}-1, 2^{t+1}-3]$,
9. $[2^{t+2}-1, 3, 2^{t+1}-3]$, 10. $[3, 2^{t+1}-1, 2^{t+2}-3]$,
11. $[3, 2^{t+2}-3, 2^{t+1}-1]$, 12. $[2^{t+1}-1, 3, 2^{t+2}-3]$,
13. $[7, 2^{t+2}-5, 2^{t+1}-3]$.

For $t = 2$, $w_{1,2,1,14} = [7, 7, 9]$.
For $t \geqslant 3$, $w_{1,t,1,14} = [7, 2^{t+1}-5, 2^{t+2}-3]$.

So, we easily obtain

Proposition 8.3.1. *For any integer $t \geqslant 2$,*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{t+2}+2^{t+1}-1} = 56.$$

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+2}+2^{t+1}-1}$. We have

Theorem 8.3.2. $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+2}+2^{t+1}-1}$ *is an $\mathbb{F}_2$-vector space with a basis consisting of all the classes represented by the monomials $a_{1,t,1,j}, j \geqslant 1$, which are determined as follows:*

For $t \geqslant 2$,

1. $(1, 1, 2^{t+1} - 2, 2^{t+2} - 1)$, 2. $(1, 1, 2^{t+2} - 1, 2^{t+1} - 2)$,
3. $(1, 2^{t+1} - 2, 1, 2^{t+2} - 1)$, 4. $(1, 2^{t+1} - 2, 2^{t+2} - 1, 1)$,
5. $(1, 2^{t+2} - 1, 1, 2^{t+1} - 2)$, 6. $(1, 2^{t+2} - 1, 2^{t+1} - 2, 1)$,
7. $(2^{t+2} - 1, 1, 1, 2^{t+1} - 2)$, 8. $(2^{t+2} - 1, 1, 2^{t+1} - 2, 1)$,
9. $(1, 1, 2^{t+1} - 1, 2^{t+2} - 2)$, 10. $(1, 1, 2^{t+2} - 2, 2^{t+1} - 1)$,
11. $(1, 2^{t+1} - 1, 1, 2^{t+2} - 2)$, 12. $(1, 2^{t+1} - 1, 2^{t+2} - 2, 1)$,
13. $(1, 2^{t+2} - 2, 1, 2^{t+1} - 1)$, 14. $(1, 2^{t+2} - 2, 2^{t+1} - 1, 1)$,
15. $(2^{t+1} - 1, 1, 1, 2^{t+2} - 2)$, 16. $(2^{t+1} - 1, 1, 2^{t+2} - 2, 1)$,
17. $(1, 2, 2^{t+1} - 3, 2^{t+2} - 1)$, 18. $(1, 2, 2^{t+2} - 1, 2^{t+1} - 3)$,
19. $(1, 2^{t+2} - 1, 2, 2^{t+1} - 3)$, 20. $(2^{t+2} - 1, 1, 2, 2^{t+1} - 3)$,
21. $(1, 2, 2^{t+1} - 1, 2^{t+2} - 3)$, 22. $(1, 2, 2^{t+2} - 3, 2^{t+1} - 1)$,
23. $(1, 2^{t+1} - 1, 2, 2^{t+2} - 3)$, 24. $(2^{t+1} - 1, 1, 2, 2^{t+2} - 3)$,
25. $(1, 3, 2^{t+1} - 4, 2^{t+2} - 1)$, 26. $(1, 3, 2^{t+2} - 1, 2^{t+1} - 4)$,
27. $(1, 2^{t+2} - 1, 3, 2^{t+1} - 4)$, 28. $(3, 1, 2^{t+1} - 4, 2^{t+2} - 1)$,
29. $(3, 1, 2^{t+2} - 1, 2^{t+1} - 4)$, 30. $(3, 2^{t+2} - 1, 1, 2^{t+1} - 4)$,
31. $(2^{t+2} - 1, 1, 3, 2^{t+1} - 4)$, 32. $(2^{t+2} - 1, 3, 1, 2^{t+1} - 4)$,
33. $(1, 3, 2^{t+1} - 3, 2^{t+2} - 2)$, 34. $(1, 3, 2^{t+2} - 2, 2^{t+1} - 3)$,
35. $(1, 2^{t+2} - 2, 3, 2^{t+1} - 3)$, 36. $(3, 1, 2^{t+1} - 3, 2^{t+2} - 2)$,
37. $(3, 1, 2^{t+2} - 2, 2^{t+1} - 3)$, 38. $(3, 2^{t+1} - 3, 1, 2^{t+2} - 2)$,
39. $(3, 2^{t+1} - 3, 2^{t+2} - 2, 1)$, 40. $(1, 3, 2^{t+1} - 2, 2^{t+2} - 3)$,
41. $(1, 3, 2^{t+2} - 3, 2^{t+1} - 2)$, 42. $(1, 2^{t+1} - 2, 3, 2^{t+2} - 3)$,
43. $(3, 1, 2^{t+1} - 2, 2^{t+2} - 3)$, 44. $(3, 1, 2^{t+2} - 3, 2^{t+1} - 2)$,
45. $(3, 2^{t+2} - 3, 1, 2^{t+1} - 2)$, 46. $(3, 2^{t+2} - 3, 2^{t+1} - 2, 1)$,
47. $(1, 3, 2^{t+1} - 1, 2^{t+2} - 4)$, 48. $(1, 3, 2^{t+2} - 4, 2^{t+1} - 1)$,
49. $(1, 2^{t+1} - 1, 3, 2^{t+2} - 4)$, 50. $(3, 1, 2^{t+1} - 1, 2^{t+2} - 4)$,
51. $(3, 1, 2^{t+2} - 4, 2^{t+1} - 1)$, 52. $(3, 2^{t+1} - 1, 1, 2^{t+2} - 4)$,
53. $(2^{t+1} - 1, 1, 3, 2^{t+2} - 4)$, 54. $(2^{t+1} - 1, 3, 1, 2^{t+2} - 4)$,
55. $(3, 2^{t+1} - 3, 2, 2^{t+2} - 3)$, 56. $(3, 2^{t+2} - 3, 2, 2^{t+1} - 3)$,
57. $(3, 3, 2^{t+1} - 4, 2^{t+2} - 3)$, 58. $(3, 3, 2^{t+2} - 3, 2^{t+1} - 4)$,
59. $(3, 2^{t+2} - 3, 3, 2^{t+1} - 4)$, 60. $(3, 3, 2^{t+1} - 3, 2^{t+2} - 4)$,
61. $(3, 3, 2^{t+2} - 4, 2^{t+1} - 3)$, 62. $(3, 2^{t+1} - 3, 3, 2^{t+2} - 4)$.
63. $(3, 4, 2^{t+2} - 5, 2^{t+1} - 3)$, 64. $(3, 4, 2^{t+1} - 5, 2^{t+2} - 3)$,
65. $(3, 5, 2^{t+2} - 5, 2^{t+1} - 4)$, 66. $(3, 5, 2^{t+2} - 6, 2^{t+1} - 3)$,
67. $(3, 7, 2^{t+2} - 7, 2^{t+1} - 4)$, 68. $(7, 3, 2^{t+2} - 7, 2^{t+1} - 4)$,
69. $(1, 6, 2^{t+2} - 5, 2^{t+1} - 3)$, 70. $(1, 7, 2^{t+2} - 6, 2^{t+1} - 3)$,
71. $(7, 1, 2^{t+2} - 6, 2^{t+1} - 3)$. 72. $(1, 7, 2^{t+2} - 5, 2^{t+1} - 4)$,
73. $(7, 1, 2^{t+2} - 5, 2^{t+1} - 4)$, 74. $(7, 2^{t+2} - 5, 1, 2^{t+1} - 4)$,

For $t = 2$,

75. $(3, 5, 6, 9)$, 76. $(3, 5, 7, 8)$, 77. $(3, 7, 5, 8)$,
78. $(7, 3, 5, 8)$, 79. $(1, 6, 7, 9)$, 80. $(1, 7, 7, 8)$,
81. $(7, 1, 7, 8)$, 82. $(7, 7, 1, 8)$, 83. $(1, 7, 6, 9)$,
84. $(7, 1, 6, 9)$, 85. $(3, 7, 8, 5)$, 86. $(7, 3, 8, 5)$,
87. $(3, 4, 1, 15)$, 88. $(3, 4, 15, 1)$, 89. $(3, 15, 4, 1)$,
90. $(15, 3, 4, 1)$, 91. $(3, 7, 12, 1)$, 92. $(7, 3, 12, 1)$,

93. $(7,11,4,1)$, 94. $(7,7,8,1)$, 95. $(7,9,2,5)$,
96. $(3,4,7,9)$, *97.* $(3,7,4,9)$, 98. $(7,3,4,9)$,
99. $(7,9,3,4)$.

For $t \geqslant 3$,

75. $(3,5,2^{t+1}-6,2^{t+2}-3)$, 76. $(3,5,2^{t+1}-5,2^{t+2}-4)$,
77. $(3,7,2^{t+1}-7,2^{t+2}-4)$, 78. $(7,3,2^{t+1}-7,2^{t+2}-4)$,
79. $(1,6,2^{t+1}-5,2^{t+2}-3)$, 80. $(1,7,2^{t+1}-5,2^{t+2}-4)$,
81. $(7,1,2^{t+1}-5,2^{t+2}-4)$, 82. $(7,2^{t+1}-5,1,2^{t+2}-4)$,
83. $(1,7,2^{t+1}-6,2^{t+2}-3)$, 84. $(7,1,2^{t+1}-6,2^{t+2}-3)$,

We prove the theorem by proving the following propositions.

Proposition 8.3.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+2}+2^{t+1}-1}$ is generated by the elements listed in Theorem 8.3.2.*

The proof of this proposition is based on the following lemmas.

Lemma 8.3.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&0&1&1\\1&0&0&1\\0&1&0&1\\0&1&0&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&1&0\\0&1&1&0\\0&1&0&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&1&0\\0&1&0&1\\0&1&0&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&0&1\\0&1&0&1\\0&0&1&0\end{pmatrix}$$

$$\begin{pmatrix}1&1&1&0\\1&0&0&1\\0&1&0&1\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&0&0&1\\0&1&0&1\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&1&0\\1&0&0&1\\0&1&0&0\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are (3,12,1,7), (3,12,7,1), (3,12,3,5), (3,4,9,7), (3,5,9,6), (3,5,8,7), (7,8,3,5). By a direct computation, we have

$$\begin{aligned}(3,12,1,7) &= Sq^1\big((3,11,1,7)+(3,7,1,11)\big)+Sq^2\big((2,11,1,7)\\&\quad+(5,7,2,7)+(2,7,1,11)\big)+Sq^4(3,7,2,7)+(2,13,1,7)+(3,9,4,7)\\&\quad+(3,7,4,9)+(2,7,1,13)+(3,7,1,12)\mod \mathcal{L}_4(3;2;2;1),\\(3,12,7,1) &= \overline{\varphi}_3(3,12,1,7),\\(3,12,3,5) &= Sq^1\big((3,11,3,5)+(3,7,3,9)\big)+Sq^2\big((2,11,3,5)\\&\quad+(5,7,3,6)+(2,7,3,9)\big)+Sq^4(3,7,3,6)+(2,13,3,5)\\&\quad+(2,11,5,5)+(3,11,4,5)+(3,9,5,6)+(3,7,5,8)\\&\quad+(2,7,5,9)+(3,7,4,9)\mod \mathcal{L}_4(3;2;2;1),\\(3,4,9,7) &= Sq^1\big((3,1,11,7)+(3,1,7,11)\big)+Sq^2\big((2,1,11,7)\\&\quad+(5,2,7,7)+(2,1,7,11)\big)+Sq^4(3,2,7,7)+(2,1,13,7)+(3,4,7,9)\\&\quad+(2,1,7,13)+(3,1,12,7)+(3,1,7,12)\mod \mathcal{L}_4(3;2;2;1),\end{aligned}$$

$$\begin{aligned}(3,5,9,6) &= Sq^1\big((3,3,11,5)+(3,3,7,9)\big)+Sq^2\big((5,3,7,6)\\ &\quad+(2,3,11,5)+(2,3,7,9)\big)+Sq^4(3,3,7,6)+(3,5,7,8)\\ &\quad+(3,4,11,5)+(3,3,12,5)+(3,4,7,9)+(2,5,11,5)\\ &\quad+(2,3,13,5)+(2,5,7,9) \mod \mathcal{L}_4(3;2;2;1),\\ (3,5,8,7) &= Sq^1\big((3,3,5,11)+(3,3,9,7)\big)+Sq^2\big((5,3,6,7)\\ &\quad+(2,3,5,11)+(2,3,9,7)\big)+Sq^4(3,3,6,7)+(3,5,6,9)\\ &\quad+(3,4,5,11)+(3,3,5,12)+(3,4,9,7)+(2,5,5,11)\\ &\quad+(2,3,5,13)+(2,5,9,7) \mod \mathcal{L}_4(3;2;2;1),\\ (7,8,3,5) &= Sq^1(7,5,5,5)+Sq^2\big((7,6,3,5)+(7,3,6,5)+(7,3,5,6)\big)\\ &\quad+Sq^4\big((5,6,3,5)+(5,3,5,6)+(5,3,6,5)\big)+(5,10,3,5)\\ &\quad+(5,6,3,9)+(7,3,5,8)+(7,3,8,5)+(5,3,9,6)\\ &\quad+(5,3,5,10)+(5,3,10,5)+(5,3,6,9) \mod \mathcal{L}_4(3;2;2;1).\end{aligned}$$

The lemma is proved. □

Lemma 8.3.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&1&0&1\\1&1&0&0\\0&1&0&1\\0&0&1&1\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&0&1\\0&0&1&1\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\0&1&0&1\\0&0&1&1\\0&0&1&0\end{pmatrix}$$

$$\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&0&1\\0&0&1&1\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&0\\1&0&1&0\\0&1&1&0\\0&0&0&1\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are $(3,7,8,29),(7,3,8,29),(3,7,24,13),(7,3,24,13),(7,11,13,16)$.

For simplicity, we prove the lemma for the matrices associated to the monomials $(7,3,24,13),(3,7,8,29),(7,11,13,16)$. By a direct computation, we have

$$\begin{aligned}(7,3,24,13) &= Sq^1\big((7,3,23,13)+(7,3,19,17)+(7,3,15,21)\big)\\ &\quad+Sq^2\big((7,2,23,13)+(7,5,19,14)+(7,5,15,18)+(7,2,15,21)\big)\\ &\quad+Sq^4\big((4,3,23,13)+(5,2,23,13)+(11,3,15,14)+(5,3,21,14)\\ &\quad+(5,3,15,20)+(4,3,15,21)+(5,2,15,21)\big)+Sq^8(7,3,15,14)\\ &\quad+(4,3,27,13)+(5,2,27,13)+(7,2,25,13)+(5,3,25,14)\\ &\quad+(5,3,15,24)+(4,3,15,25)+(5,2,15,25) \mod \mathcal{L}_4(3;2;2;2;1),\end{aligned}$$

$$\begin{aligned}(3,7,8,29) &= Sq^1\big((3,7,9,27)+(3,7,11,25)+(3,7,13,23)\\
&\quad+(3,7,21,15)+(3,7,17,19)\big)+Sq^2\big((5,7,10,23)+(5,7,6,27)\\
&\quad+(2,7,9,27)+(5,7,11,22)+(5,7,7,26)+(2,7,11,25)\\
&\quad+(2,7,13,23)+(5,7,18,15)+(5,7,14,19)+(2,7,21,15)\big)\\
&\quad+Sq^4\big((3,5,12,23)+(3,5,6,29)+(3,11,6,23)+(3,11,7,22)\\
&\quad+(3,5,7,28)+(3,5,13,22)+(3,4,13,23)+(3,11,14,15)\\
&\quad+(3,5,20,15)+(3,5,14,21)+(3,4,21,15)\big)\\
&\quad+Sq^8\big((3,7,6,23)+(3,7,7,22)+(3,7,14,15)\big)+(3,5,12,27)\\
&\quad+(3,5,10,29)+(2,7,9,29)+(3,5,13,26)+(3,5,11,28)\\
&\quad+(2,9,13,23)+(3,4,13,27)+(3,5,24,15)+(3,5,14,25)\\
&\quad+(2,9,21,15)+(3,4,25,15) \bmod \mathcal{L}_4(3;2;2;2;1),\\
(7,11,13,16) &= Sq^1(7,13,13,13)+Sq^2\big((7,11,13,14)+(7,14,11,13)\\
&\quad+(7,11,14,13)+(7,7,18,13)+(7,7,10,21)\big)+Sq^4\big((5,11,13,14)\\
&\quad+(5,14,11,13)+(5,11,14,13)+(11,7,12,13)+(5,7,18,13)\\
&\quad+(5,7,10,21)\big)+Sq^8\big((7,8,11,13)+(7,7,12,13)\big)+(5,11,17,14)\\
&\quad+(5,11,13,18)+(5,18,11,13)+(5,14,11,17)+(7,8,19,13)\\
&\quad+(7,8,11,21)+(5,11,14,17)+(7,11,12,17)+(7,9,18,13)\\
&\quad+(7,9,10,21)+(5,11,10,21)+(5,7,10,25) \quad \bmod \mathcal{L}_4(3;2;2;2;1).\end{aligned}$$

The lemma is proved. □

Using the results in Section 7, Lemmas 5.10, 5.12, 5.13, 8.3.4, 8.3.5 and Theorem 2.5, we get Proposition 8.3.3.

Now, we show that the elements listed in Theorem 8.3.2 are linearly independent.

Proposition 8.3.6. *The elements* $[a_{1,2,1,j}], 1 \leqslant j \leqslant 99$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{23}$.

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{99} \gamma_j [a_{1,2,1,j}] = 0, \tag{8.3.6.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms f_1, f_2, f_3 to the relation (8.3.6.1) and we get

$$\begin{aligned}&\gamma_{\{3,87\}}[7,1,15]+\gamma_{\{4,88\}}[7,15,1]+\gamma_{13}[15,1,7]+\gamma_{14}[15,7,1]\\
&\quad+\gamma_{17}[3,5,15]+\gamma_{18}[3,15,5]+\gamma_{35}[15,3,5]+\gamma_{21}[3,7,13]+\gamma_{22}[3,13,7]\\
&\quad+\gamma_{\{42,64\}}[7,3,13]+\gamma_{\{63,69\}}[7,11,5]+\gamma_{\{79,96\}}[7,7,9]=0,\end{aligned}$$

$$\begin{aligned}
&\gamma_{\{1,28\}}[7,1,15] + \gamma_{\{6,89\}}[7,15,1] + \gamma_{\{10,51\}}[15,1,7]\\
&\quad + \gamma_{\{12,91,94\}}[15,7,1] + \gamma_{25}[3,5,15] + \gamma_{19}[3,15,5]\\
&\quad + \gamma_{\{34,61,86\}}[15,3,5] + \gamma_{\{48,70,83,85,97\}}[7,7,9] + \gamma_{48}[3,13,7]\\
&\quad + \gamma_{\{40,48,57\}}[7,3,13] + \gamma_{\{70,85\}}[7,11,5] + \gamma_{\{23,48\}}[3,7,13] = 0,\\
&\gamma_{\{2,29\}}[7,1,15] + \gamma_{\{5,30\}}[7,15,1] + \gamma_{\{9,50,81\}}[15,1,7] + \gamma_{\{11,52,82\}}[15,7,1]\\
&\quad + \gamma_{26}[3,5,15] + \gamma_{27}[3,15,5] + \gamma_{\{33,60,62,78\}}[15,3,5]\\
&\quad + \gamma_{\{47,72,76,80\}}[3,7,13] + \gamma_{\{47,76,80\}}[3,13,7] + \gamma_{\{41,47,58,65,76\}}[7,3,13]\\
&\quad + \gamma_{\{49,59,77\}}[7,11,5] + \gamma_{\{47,49,67,76,77\}}[7,7,9] = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\text{(8.3.6.2)}\quad \begin{cases}
\gamma_j = 0,\ j = 13,14,17,18,19,21,22,23,25,26,27,35,48,\\
\gamma_{\{3,87\}} = \gamma_{\{4,88\}} = \gamma_{\{42,64\}} = \gamma_{\{63,69\}} = 0,\\
\gamma_{\{79,96\}} = \gamma_{\{1,28\}} = \gamma_{\{6,89\}} = \gamma_{\{10,51\}} = 0,\\
\gamma_{\{12,91,94\}} = \gamma_{\{34,61,86\}} = \gamma_{\{40,57\}} = \gamma_{\{70,85\}} = 0,\\
\gamma_{\{70,83,85,97\}} = \gamma_{\{2,29\}} = \gamma_{\{5,30\}} = \gamma_{\{9,50,81\}} = 0,\\
\gamma_{\{11,52,82\}} = \gamma_{\{33,60,62,78\}} = \gamma_{\{47,72,76,80\}} = \gamma_{\{47,76,80\}} = 0,\\
\gamma_{\{41,47,58,65,76\}} = \gamma_{\{49,59,77\}} = \gamma_{\{47,49,67,76,77\}} = 0.
\end{cases}$$

With the aid of (8.3.6.2), the homomorphisms f_4, f_5, f_6 send (8.3.6.1) to

$$\begin{aligned}
&\gamma_{\{1,3\}}[1,7,15] + \gamma_{10}[1,15,7] + \gamma_{\{16,92,93,94\}}[7,15,1] + \gamma_{\{8,90\}}[15,7,1]\\
&\quad + \gamma_{\{28,87\}}[3,5,15] + a_1[3,15,5] + \gamma_{20}[15,3,5] + \gamma_{\{43,55,57,64\}}[3,7,13]\\
&\quad + \gamma_{51}[3,13,7] + \gamma_{24}[7,3,13] + \gamma_{\{71,86,95\}}[7,11,5] + \gamma_{\{84,98\}}[7,7,9] = 0,\\
&\gamma_{\{2,4\}}[1,7,15] + \gamma_{\{9,47,79,80\}}[1,15,7] + \gamma_{\{15,54,74,82\}}[7,15,1]\\
&\quad + \gamma_{\{7,32\}}[15,7,1] + \gamma_{\{29,88\}}[3,5,15] + \gamma_{\{36,60,77\}}[3,15,5]\\
&\quad + \gamma_{31}[15,3,5] + \gamma_{\{44,58,81\}}[3,7,13] + \gamma_{\{50,76,81,96\}}[3,13,7]\\
&\quad + \gamma_{73}[7,3,13] + \gamma_{\{53,78,99\}}[7,11,5] + \gamma_{\{68,81\}}[7,7,9] = 0,\\
&a_2[1,7,15] + \gamma_{\{5,6\}}[1,15,7] + a_3[7,1,15] + \gamma_{\{7,8,20,31\}}[15,1,7]\\
&\quad + a_4[3,5,15] + \gamma_{\{30,89\}}[3,15,5] + \gamma_{\{32,90\}}[15,3,5] + a_5[3,7,13]\\
&\quad + a_6[3,13,7] + a_7[7,3,13] + \gamma_{\{74,93\}}[7,11,5] + \gamma_{\{82,94,95,99\}}[7,7,9] = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{37,56,61,63,66,85\}},\quad a_2 = \gamma_{\{11,12,49,70,72,80,83\}},\\
&a_3 = \gamma_{\{15,16,24,53,71,73,81,84\}},\quad a_4 = \gamma_{\{38,39,55,62,65,66,75,76\}},\\
&a_5 = \gamma_{\{52,67,77,85,91,95,97,99\}},\quad a_6 = \gamma_{\{45,46,56,59,95,99\}},\\
&a_7 = \gamma_{\{54,68,78,86,92,98\}}.
\end{aligned}$$

From these equalities, we get

$$(8.3.6.3)\quad \begin{cases} a_i = 0,\ i = 1, 2, \ldots, 7,\ \gamma_j = 0,\ j = 10, 20, 51, 24, 31, 73, \\ \gamma_{\{15,54,74,82\}} = \gamma_{\{16,92,93,94\}} = \gamma_{\{1,3\}} = \gamma_{\{2,4\}} = 0, \\ \gamma_{\{28,87\}} = \gamma_{\{29,88\}} = \gamma_{\{30,89\}} = \gamma_{\{32,90\}} = \gamma_{\{36,60,77\}} = 0, \\ \gamma_{\{43,55,57,64\}} = \gamma_{\{44,58,81\}} = \gamma_{\{5,6\}} = \gamma_{\{50,76,81,96\}} = 0, \\ \gamma_{\{53,78,99\}} = \gamma_{\{7,32\}} = \gamma_{\{7,8\}} = \gamma_{\{68,81\}} = \gamma_{\{71,86,95\}} = 0, \\ \gamma_{\{74,93\}} = \gamma_{\{8,90\}} = \gamma_{\{82,94,95,99\}} = \gamma_{\{84,98\}} = \gamma_{\{9,47,79,80\}} = 0. \end{cases}$$

With the aid of (8.3.6.2) and (8.3.6.3), the homomorphisms g_1, g_2 send (8.3.6.1) to

$$\begin{aligned} &a_8[7,15,1] + \gamma_{\{46,74\}}[15,7,1] + a_9[3,15,5] \\ &\quad + \gamma_{\{56,95\}}[15,3,5] + \gamma_{43}[3,7,13] + \gamma_{55}[7,3,13] \\ &\quad + \gamma_{\{66,71,86,95\}}[7,11,5] + \gamma_{\{75,95\}}[7,7,9] = 0, \\ &\gamma_{\{9,50,68\}}[1,15,7] + a_{10}[7,15,1] + \gamma_{\{45,74\}}[15,7,1] \\ &\quad + a_{11}[3,15,5] + \gamma_{\{59,99\}}[15,3,5] + a_{12}[3,7,13] + a_{13}[3,13,7] \\ &\quad + \gamma_{\{65,72\}}[7,3,13] + a_{14}[7,11,5] + a_{15}[7,7,9] = 0, \end{aligned}$$

where

$$\begin{aligned} &a_8 = \gamma_{\{12,16,39,91,92,94\}},\quad a_9 = \gamma_{\{34,37,61,71,86\}},\quad a_{10} = \gamma_{\{11,15,38,52,54,82\}}, \\ &a_{11} = \gamma_{\{33,36,42,53,60,62,78\}},\quad a_{12} = \gamma_{\{41,44,58,63,65,68,76,80\}}, \\ &a_{13} = \gamma_{\{47,50,68,79,80\}},\quad a_{14} = \gamma_{\{49,53,59,62,77,78,99\}},\quad a_{15} = \gamma_{\{59,67,76,80,99\}}. \end{aligned}$$

These equalities imply

$$(8.3.6.4)\quad \begin{cases} \gamma_j = 0, j = 43, 55,\ a_i = 0, i = 8, 9, \ldots, 15, \\ \gamma_{\{45,74\}} = \gamma_{\{46,74\}} = \gamma_{\{56,95\}} = \gamma_{\{59,99\}} = 0, \\ \gamma_{\{66,71,86,95\}} = \gamma_{\{65,72\}} = \gamma_{\{75,95\}} = \gamma_{\{9,50,68\}} = 0. \end{cases}$$

With the aid of (8.3.6.2), (8.3.6.3) and (8.3.6.4), the homomorphisms g_3, g_4 send (8.3.6.1) to

$$\begin{aligned} &\gamma_{\{11,52,82\}}[1,15,7] + a_{16}[7,15,1] + \gamma_{\{12,91,94\}}[15,1,7] + a_{17}[3,15,5] \\ &\quad + \gamma_{\{16,37,44,71\}}[15,7,1] + a_{18}[15,3,5] + a_{19}[3,7,13] + a_{20}[3,13,7] \\ &\quad + \gamma_{\{46,67,72,91\}}[7,3,13] + a_{21}[7,11,5] + a_{22}[7,7,9] = 0, \\ &\gamma_{\{15,54,74,82\}}[1,15,7] + a_{23}[7,15,1] + \gamma_{\{16,74,92,94\}}[15,1,7] \\ &\quad + a_{24}[15,7,1] + a_{25}[3,15,5] + a_{26}[15,3,5] + a_{27}[3,7,13] \\ &\quad + a_{28}[3,13,7] + a_{29}[7,3,13] + a_{30}[7,11,5] + a_{31}[7,7,9] = 0, \end{aligned}$$

where

$$\begin{aligned} &a_{16} = \gamma_{\{9,15,36,50,53,68,84\}},\quad a_{17} = \gamma_{\{33,38,40,54,60,62,78,84\}}, \\ &a_{18} = \gamma_{\{34,39,58,61,68,86,92\}},\quad a_{19} = \gamma_{\{45,56,59,70,72,77,80,82,91,94\}}, \\ &a_{20} = \gamma_{\{49,52,70,72,80,82,83,91,94\}},\quad a_{21} = \gamma_{\{41,47,54,58,60,61,65,66,68,75,76,78,84,86,92\}}, \end{aligned}$$

$$a_{22} = \gamma_{\{41,58,61,65,66,67,68,72,77,80,82,86,91,92,95,99\}},$$
$$a_{23} = \gamma_{\{9,11,33,40,42,47,49,79,80,83\}}, \quad a_{24} = \gamma_{\{12,34,41,63,70,72\}},$$
$$a_{25} = \gamma_{\{36,38,40,45,52,60,77,83\}}, \quad a_{26} = \gamma_{\{37,39,46,56,58,59,61,63,65,66,67,70,91\}},$$
$$a_{27} = \gamma_{\{68,71,78,82,92,94,95,99\}}, \quad a_{28} = \gamma_{\{53,54,68,71,82,84,92,94\}},$$
$$a_{29} = \gamma_{\{68,71,86,92,95,99\}}, \quad a_{31} = \gamma_{\{44,45,58,61,67,70,71,74,78,82,86,91,92,95,99\}}$$
$$a_{30} = \gamma_{\{40,42,44,45,46,50,52,56,58,59,60,61,62,67,70,75,76,77,79,83,91\}}.$$

From these equalities, we get

$$\begin{cases} a_i = 0, i = 16, 17, \ldots, 31, \\ \gamma_{\{11,52,82\}} = \gamma_{\{15,54,74,82\}} = \gamma_{\{16,37,44,71\}} = 0, \\ \gamma_{\{12,91,94\}} = \gamma_{\{16,74,92,94\}} = \gamma_{\{46,67,72,91\}} = 0. \end{cases} \tag{8.3.6.5}$$

Substituting (8.3.6.2), (8.3.6.3), (8.3.6.4) and (8.3.6.5) into the relation (8.3.6.1), we obtain

$$\sum_{i=1}^{15} c_i[\theta_i] = 0, \tag{8.3.6.6}$$

where $c_1 = \gamma_1, c_2 = \gamma_2, c_3 = \gamma_5, c_4 = \gamma_7, c_5 = \gamma_9, c_6 = \gamma_{11}, c_7 = \gamma_{12}, c_8 = \gamma_{15}, c_9 = \gamma_{16}, c_{10} = \gamma_{33}, c_{11} = \gamma_{37}, c_{12} = \gamma_{38}, c_{13} = \gamma_{40}, c_{14} = \gamma_{45}, c_{15} = \gamma_{59}$ and

$$\begin{aligned}
\theta_1 &= a_{1,2,1,1} + a_{1,2,1,3} + a_{1,2,1,28} + a_{1,2,1,87}, \\
\theta_2 &= a_{1,2,1,2} + a_{1,2,1,4} + a_{1,2,1,29} + a_{1,2,1,88}, \\
\theta_3 &= a_{1,2,1,5} + a_{1,2,1,6} + a_{1,2,1,30} + a_{1,2,1,89}, \\
\theta_4 &= a_{1,2,1,7} + a_{1,2,1,8} + a_{1,2,1,32} + a_{1,2,1,90}, \\
\theta_5 &= a_{1,2,1,9} + a_{1,2,1,50} + a_{1,2,1,79} + a_{1,2,1,96}, \\
\theta_6 &= a_{1,2,1,11} + a_{1,2,1,52} + a_{1,2,1,83} + a_{1,2,1,97}, \\
\theta_7 &= a_{1,2,1,12} + a_{1,2,1,41} + a_{1,2,1,47} + a_{1,2,1,67} + a_{1,2,1,80} + a_{1,2,1,91}, \\
\theta_8 &= a_{1,2,1,15} + a_{1,2,1,54} + a_{1,2,1,84} + a_{1,2,1,98}, \\
\theta_9 &= a_{1,2,1,16} + a_{1,2,1,44} + a_{1,2,1,50} + a_{1,2,1,68} + a_{1,2,1,81} + a_{1,2,1,92}, \\
\theta_{10} &= a_{1,2,1,33} + a_{1,2,1,34} + a_{1,2,1,49} + a_{1,2,1,60} \\
&\quad + a_{1,2,1,61} + a_{1,2,1,70} + a_{1,2,1,77} + a_{1,2,1,85}, \\
\theta_{11} &= a_{1,2,1,36} + a_{1,2,1,37} + a_{1,2,1,53} + a_{1,2,1,60} \\
&\quad + a_{1,2,1,61} + a_{1,2,1,71} + a_{1,2,1,78} + a_{1,2,1,86}, \\
\theta_{12} &= a_{1,2,1,38} + a_{1,2,1,52} + a_{1,2,1,54} + a_{1,2,1,56} + a_{1,2,1,61} \\
&\quad + a_{1,2,1,75} + a_{1,2,1,82} + a_{1,2,1,86} + a_{1,2,1,95},
\end{aligned}$$

$$\begin{aligned}
\theta_{13} &= a_{1,2,1,36} + a_{1,2,1,39} + a_{1,2,1,40} + a_{1,2,1,42} + a_{1,2,1,47} + a_{1,2,1,49}\\
&\quad + a_{1,2,1,50} + a_{1,2,1,56} + a_{1,2,1,57} + a_{1,2,1,58} + a_{1,2,1,61}\\
&\quad + a_{1,2,1,63} + a_{1,2,1,64} + a_{1,2,1,67} + a_{1,2,1,68} + a_{1,2,1,69}\\
&\quad + a_{1,2,1,70} + a_{1,2,1,75} + a_{1,2,1,77} + a_{1,2,1,80} + a_{1,2,1,81}\\
&\quad + a_{1,2,1,83} + a_{1,2,1,84} + a_{1,2,1,85} + a_{1,2,1,86} + a_{1,2,1,91}\\
&\quad + a_{1,2,1,92} + a_{1,2,1,94} + a_{1,2,1,95} + a_{1,2,1,97} + a_{1,2,1,98},\\
\theta_{14} &= a_{1,2,1,45} + a_{1,2,1,46} + a_{1,2,1,52} + a_{1,2,1,74}\\
&\quad + a_{1,2,1,82} + a_{1,2,1,91} + a_{1,2,1,93} + a_{1,2,1,94},\\
\theta_{15} &= a_{1,2,1,34} + a_{1,2,1,39} + a_{1,2,1,47} + a_{1,2,1,59} + a_{1,2,1,60} + a_{1,2,1,61}\\
&\quad + a_{1,2,1,63} + a_{1,2,1,67} + a_{1,2,1,76} + a_{1,2,1,77} + a_{1,2,1,78} + a_{1,2,1,69}\\
&\quad + a_{1,2,1,79} + a_{1,2,1,91} + a_{1,2,1,92} + a_{1,2,1,94} + a_{1,2,1,96} + a_{1,2,1,99}.
\end{aligned}$$

Now, we show that $c_i = 0$ for $i = 1, 2, \ldots, 15$. The proof is divided into 6 steps.

Step 1. Set $\theta = \theta_1 + \sum_{i=2}^{15} \beta_i\theta_i$ for $\beta_i \in \mathbb{F}_2, i = 2, 3, \ldots, 15$. We prove that θ is non-hit. Suppose the contrary that θ is hit. Then we have

$$\theta = Sq^1(A) + Sq^2(B) + Sq^4(C) + Sq^8(D)$$

for some polynomials $A, B, C, D \in R_4$. Let $(Sq^2)^3$ act to the both sides of the above equality, we obtain

$$(Sq^2)^3(\theta) = (Sq^2)^3Sq^4(C) + (Sq^2)^3Sq^8(D).$$

By a direct computation we see that the monomial $(8, 4, 2, 15)$ is a term of $(Sq^2)^3(\theta)$. This monomial is not a term of $(Sq^2)^3(Sq^4(C) + Sq^8(D))$ for all polynomials C, D and we have a contradiction. So, θ is non-hit and we get $c_1 = \gamma_1 = 0$.

By an argument analogous to the previous one, we get $c_2 = c_3 = c_4 = 0$. Now, the relation (8.3.6.6) becomes

$$\sum_{i=5}^{15} c_i[\theta_i] = 0. \tag{8.3.6.7}$$

Step 2. The homomorphism φ_4 sends (8.3.6.7) to

$$c_{14}[\theta_3] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.3.6.8}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.3.6.8) and by a same argument as given in Step 1, we get $c_{14} = \gamma_{45} = 0$.

Under the homomorphism induced by the linear transformation $x_1 \mapsto x_1 + x_3, x_2 \mapsto x_2, x_3 \mapsto x_3, x_4 \mapsto x_4$, the linear relation (8.3.6.7) is sent to

$$c_9[\theta_2] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.3.6.9}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.3.6.9) and by a same argument as given in Step 1, we get $c_9 = \gamma_{16} = 0$.

Consider the homomorphism $\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4 \to \mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4$ induced by the linear transformation $x_1 \mapsto x_1 + x_4, x_2 \mapsto x_2, x_3 \mapsto x_3, x_4 \mapsto x_4$. Under this homomorphism, the linear relation (8.3.6.7) is sent to

$$c_8[\theta_1] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.3.6.10}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.3.6.10) and by a same argument as given in Step 1, we get $c_8 = \gamma_{15} = 0$.

The homomorphism induced by the linear transformation $x_1 \mapsto x_1, x_2 \mapsto x_2 + x_3, x_3 \mapsto x_3, x_4 \mapsto x_4$ sends the linear relation (8.3.6.7) to

$$c_7[\theta_3] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.3.6.11}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.3.6.11) and by a same argument as given in Step 1, we get $c_7 = \gamma_{12} = 0$.

Applying the homomorphism induced by the linear map $x_1 \mapsto x_1, x_2 \mapsto x_2 + x_4, x_3 \mapsto x_3, x_4 \mapsto x_4$ to the linear relation (8.3.6.7), we get

$$c_6[\theta_1] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.3.6.12}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.3.6.12) and by a same argument as given in Step 1, we get $c_6 = \gamma_{11} = 0$.

Consider the homomorphism induced by the linear map $x_1 \mapsto x_1, x_2 \mapsto x_2, x_3 \mapsto x_3 + x_4, x_4 \mapsto x_4$. This homomorphism sends the linear relation (8.3.6.7) to

$$c_5[\theta_1] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.3.6.13}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.3.6.13) and by a same argument as given in Step 1, we get $c_5 = \gamma_9 = 0$.

So, the relation (8.3.6.7) becomes

$$c_{10}[\theta_{10}] + c_{11}[\theta_{11}] + c_{12}[\theta_{12}] + c_{13}[\theta_{13}] + c_{15}[\theta_{15}] = 0. \tag{8.3.6.14}$$

Step 3. The homomorphism φ_4 sends (8.3.6.14) to

$$\begin{aligned} &c_{12}[\theta_6] + (c_{13} + c_{15})[\theta_7] + (c_{10} + c_{11})[\theta_{10}] \\ &\quad + c_{11}[\theta_{11}] + c_{12}[\theta_{12}] + c_{13}[\theta_{13}] + c_{15}[\theta_{15}] = 0. \end{aligned}$$

Using this equality and by a same argument as given in Step 2, we get $c_{12} = 0$ and $c_{13} = c_{15}$. So, the relation (8.3.6.14) becomes

$$c_{10}[\theta_{10}] + c_{11}[\theta_{11}] + c_{13}[\theta_{13}] + c_{13}[\theta_{15}] = 0. \tag{8.3.6.15}$$

Step 4. The homomorphism φ_2 sends (8.3.6.15) to

$$c_{10}[\theta_{10}] + c_{13}[\theta_{11}] + (c_{11} + c_{13})[\theta_{12}] + c_{13}[\theta_{13}] + c_{13}[\theta_{15}] = 0.$$

Using this equality and by a same argument as given in Step 2, we get $c_{11} = c_{13}$. Then the relation (8.3.6.15) becomes

$$c_{10}[\theta_{10}] + c_{11}[\theta_{11}] + c_{11}[\theta_{13}] + c_{11}[\theta_{15}] = 0. \tag{8.3.6.16}$$

Step 5. The homomorphism φ_1 send (8.3.6.16) to

$$(c_{10} + c_{11})[\theta_{11}] + c_{11}[\theta_{13}] + c_{11}[\theta_{15}] = 0.$$

Using this equality and by a same argument as given in Step 4, we get $c_{10} = 0$. Then the relation (8.3.6.16) becomes

$$c_{11}([\theta_{11}] + [\theta_{13}] + [\theta_{15}]) = 0. \tag{8.3.6.17}$$

Step 6. The homomorphism φ_4 sends (8.3.6.17) to

$$c_{11}[\theta_{10}] + c_{11}([\theta_{11}] + [\theta_{13}] + [\theta_{15}]) = 0.$$

By an argument analogous to Step 5, we have $c_{11} = 0$.

The proposition is completely proved. □

Proposition 8.3.7. *For $t \geqslant 3$, the elements $[a_{1,t,1,j}], 1 \leqslant j \leqslant 84$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+2}+2^{t+1}-1}$.*

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{84} \gamma_j [a_{1,t,1,j}] = 0, \tag{8.3.7.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms f_1, f_2, f_3 to the relation (8.3.7.1), we obtain

$$\begin{aligned}
&\gamma_3 w_{1,t,1,3} + \gamma_4 w_{1,t,1,4} + \gamma_{13} w_{1,t,1,5} + \gamma_{14} w_{1,t,1,6} + \gamma_{17} w_{1,t,1,7}\\
&\quad + \gamma_{18} w_{1,t,1,8} + \gamma_{35} w_{1,t,1,9} + \gamma_{21} w_{1,t,1,10} + \gamma_{22} w_{1,t,1,11}\\
&\quad + \gamma_{42} w_{1,t,1,12} + \gamma_{\{63,69\}} w_{1,t,1,13} + \gamma_{\{64,79\}} w_{1,t,1,14} = 0,\\
&\gamma_{\{1,28\}} w_{1,t,1,3} + \gamma_6 w_{1,t,1,4} + \gamma_{\{10,51\}} w_{1,t,1,5} + \gamma_{12} w_{1,t,1,6}\\
&\quad + \gamma_{25} w_{1,t,1,7} + \gamma_{19} w_{1,t,1,8} + \gamma_{\{34,61\}} w_{1,t,1,9} + \gamma_{23} w_{1,t,1,10} + \gamma_{48} w_{1,t,1,11}\\
&\quad + \gamma_{\{40,57\}} w_{1,t,1,12} + \gamma_{70} w_{1,t,1,13} + \gamma_{\{48,70,83\}} w_{1,t,1,14} = 0,\\
&\gamma_{\{2,29\}} w_{1,t,1,3} + \gamma_{\{5,30\}} w_{1,t,1,4} + \gamma_{\{9,50\}} w_{1,t,1,5} + \gamma_{\{11,52\}} w_{1,t,1,6}\\
&\quad + \gamma_{26} w_{1,t,1,7} + \gamma_{27} w_{1,t,1,8} + \gamma_{\{33,60,62,76,77\}} w_{1,t,1,9}\\
&\quad + \gamma_{49} w_{1,t,1,10} + \gamma_{47} w_{1,t,1,11} + \gamma_{\{41,58,59,65,67\}} w_{1,t,1,12}\\
&\quad + \gamma_{\{49,59,80\}} w_{1,t,1,13} + \gamma_{\{47,59,72,80\}} w_{1,t,1,14} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\begin{cases} \gamma_j = 0,\ j = 3,4,6,12,13,14,17,18,19,21,22,23,\\ \qquad\qquad 25,26,27,35,42,47,48,49,70,83,\\ \gamma_{\{64,79\}} = \gamma_{\{63,69\}} = \gamma_{\{34,61\}} = \gamma_{\{33,60,62,76,77\}} = 0,\\ \gamma_{\{40,57\}} = \gamma_{\{59,72,80\}} = \gamma_{\{41,58,59,65,67\}} = \gamma_{\{59,80\}} = 0. \end{cases} \tag{8.3.7.2}$$

With the aid of (8.3.7.2), the homomorphisms f_4, f_5, f_6 send (8.3.7.1) to

$$\begin{aligned} &\gamma_1 w_{1,t,1,1} + \gamma_{10} w_{1,t,1,2} + \gamma_{16} w_{1,t,1,4} + \gamma_8 w_{1,t,1,6} + \gamma_{28} w_{1,t,1,7}\\ &\quad + \gamma_{\{37,56,61,63,66\}} w_{1,t,1,8} + \gamma_{20} w_{1,t,1,9} + \gamma_{\{43,55,57,64,75\}} w_{1,t,1,10}\\ &\quad + \gamma_{51} w_{1,t,1,11} + \gamma_{24} w_{1,t,1,12} + \gamma_{71} w_{1,t,1,13} + \gamma_{84} w_{1,t,1,14} = 0,\\ &\gamma_2 w_{1,t,1,1} + \gamma_9 w_{1,t,1,2} + \gamma_{\{15,54\}} w_{1,t,1,4} + \gamma_{\{7,32\}} w_{1,t,1,6} + \gamma_{29} w_{1,t,1,7}\\ &\quad + \gamma_{\{36,60,78\}} w_{1,t,1,8} + \gamma_{31} w_{1,t,1,9} + \gamma_{\{44,58,68\}} w_{1,t,1,10}\\ &\quad + \gamma_{50} w_{1,t,1,11} + \gamma_{53} w_{1,t,1,12} + \gamma_{\{53,81\}} w_{1,t,1,13} + \gamma_{\{53,73\}} w_{1,t,1,14} = 0,\\ &\gamma_{11} w_{1,t,1,1} + \gamma_5 w_{1,t,1,2} + \gamma_{\{15,16,24,53\}} w_{1,t,1,3} + \gamma_{\{7,8,20,31\}} w_{1,t,1,5}\\ &\quad + \gamma_{\{38,39,55,62\}} w_{1,t,1,7} + \gamma_{30} w_{1,t,1,8} + \gamma_{32} w_{1,t,1,9} + \gamma_{52} w_{1,t,1,10}\\ &\quad + \gamma_{\{45,46,56,59\}} w_{1,t,1,11} + \gamma_{54} w_{1,t,1,12} + \gamma_{74} w_{1,t,1,13} + \gamma_{82} w_{1,t,1,14} = 0. \end{aligned}$$

Computing from these equalities and using (8.3.7.2), we get

$$\begin{cases} \gamma_j = 0,\ j = 1,2,5,7,8,9,10,11,15,16,20,24,28,29,\\ \qquad\qquad 30,31,32,50,51,52,53,54,71,73,74,81,82,84,\\ \gamma_{\{37,56,61,63,66\}} = \gamma_{\{36,60,78\}} = \gamma_{\{43,55,57,64,75\}} = 0,\\ \gamma_{\{44,58,68\}} = \gamma_{\{38,39,55,62\}} = \gamma_{\{45,46,56,59\}} = 0. \end{cases} \tag{8.3.7.3}$$

With the aid of (8.3.7.2) and (8.3.7.3), the homomorphisms g_1, g_2 send (8.3.7.1) to

$$\begin{aligned} &\gamma_{39} w_{1,t,1,4} + \gamma_{46} w_{1,t,1,6} + \gamma_{37} w_{1,t,1,8} + \gamma_{\{40,43,57\}} w_{1,t,1,10}\\ &\quad + \gamma_{56} w_{1,t,1,9} + \gamma_{55} w_{1,t,1,12} + \gamma_{66} w_{1,t,1,13} + \gamma_{75} w_{1,t,1,14} = 0,\\ &\gamma_{38} w_{1,t,1,4} + \gamma_{45} w_{1,t,1,6} + \gamma_{\{33,36,60,62,64,76,77,78\}} w_{1,t,1,8}\\ &\quad + \gamma_{59} w_{1,t,1,9} + \gamma_{\{41,44,58,59,63,65,67,68\}} w_{1,t,1,10}\\ &\quad + \gamma_{62} w_{1,t,1,12} + \gamma_{\{62,76\}} w_{1,t,1,13} + \gamma_{\{62,65\}} w_{1,t,1,14} = 0. \end{aligned}$$

These equalities imply

$$\begin{cases} \gamma_j = 0,\ j = 37,38,39,45,46,55,56,59,62,65,66,75,76,\\ \gamma_{\{40,43,57\}} = \gamma_{\{33,36,60,64,77,78\}} = \gamma_{\{41,44,58,63,67,68\}} = 0\\ \gamma_{\{33,36,60,62,64,76,77,78\}} = \gamma_{\{41,44,58,59,63,65,67,68\}} = 0. \end{cases} \tag{8.3.7.4}$$

With the aid of (8.3.7.2), (8.3.7.3) and (8.3.7.4), the homomorphisms g_3, g_4 send (8.3.7.1) to

$$\begin{aligned}
&\gamma_{\{36,43\}}w_{1,t,1,4} + \gamma_{44}w_{1,t,1,6} + \gamma_{\{33,57,60\}} + \gamma_{\{77,78\}}w_{1,t,1,8}\\
&\quad + \gamma_{\{58,68\}}w_{1,t,1,9} + \gamma_{\{34,41,58\}} + \gamma_{\{67,68\}}w_{1,t,1,10}\\
&\quad + \gamma_{\{40,57,60,78\}}w_{1,t,1,12} + a_2w_{1,t,1,13} + a_1w_{1,t,1,14} = 0,\\
&\gamma_{\{33,40,64\}}w_{1,t,1,4} + \gamma_{\{34,41,63\}}w_{1,t,1,6} + \gamma_{\{36,57,60,77,78\}}w_{1,t,1,8} + a_3w_{1,t,1,10}\\
&\quad + \gamma_{\{34,58,63,67\}}w_{1,t,1,9} + a_4w_{1,t,1,12} + a_6w_{1,t,1,13} + a_5w_{1,t,1,14} = 0,
\end{aligned}$$

where

$$a_1 = \gamma_{\{40,57,60,67,78\}},\quad a_2 = \gamma_{\{34,40,41,57,58,60,68,77,78\}},\quad a_3 = \gamma_{\{34,44,58,67,68\}},$$
$$a_4 = \gamma_{\{43,57,60,64,77\}},\quad a_5 = \gamma_{\{43,57,60,64,68,77\}},\quad a_6 = \gamma_{\{34,43,44,57,58,60,64,67,77,78\}}.$$

From these equalities, we get

$$\text{(8.3.7.5)}\qquad \begin{cases}
\gamma_{44} = 0,\ a_i = 0, i = 1, 2, \ldots, 6,\\
\gamma_{\{36,43\}} = \gamma_{\{33,57,60,77,78\}} = \gamma_{\{58,68\}} = \gamma_{\{34,41,58,67,68\}} = 0,\\
\gamma_{\{40,57,60,78\}} = \gamma_{\{33,40,64\}} = \gamma_{\{34,58,63,67\}} = 0,\\
\gamma_{\{34,41,63\}} = \gamma_{\{36,57,60,77,78\}} = 0.
\end{cases}$$

Combining (8.3.7.2), (8.3.7.3), (8.3.7.4) and (8.3.7.5), we get $\gamma_j = 0$ for all j. The proposition is proved. □

Remark 8.3.8. From the proof of Proposition 8.3.6, we see that the $\mathbb{F}_2$-subspace of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{23}$ generated by $[\theta_i], i = 1, 2, \ldots, 15$, is an $GL_4(\mathbb{F}_2)$-submodule of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{23}$.

8.4. **The case $s = 1, t \geqslant 2$ and $u \geqslant 2$.**

Using the results in [8], for $t \geqslant 2$ and $u \geqslant 2$, we see that the dimension of the space $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{t+u+1}+2^{t+1}-1}$ is 21 with a basis given by the classes $w_{u,t,1,j}, 1 \leqslant j \leqslant 21$, which are determined as follows:

For $t \geqslant 2$,

1. $[1, 2^{t+1} - 1, 2^{t+u+1} - 1]$,
2. $[1, 2^{t+u+1} - 1, 2^{t+1} - 1]$,
3. $[2^{t+1} - 1, 1, 2^{t+u+1} - 1]$,
4. $[2^{t+1} - 1, 2^{t+u+1} - 1, 1]$,
5. $[2^{t+u+1} - 1, 1, 2^{t+1} - 1]$,
6. $[2^{t+u+1} - 1, 2^{t+1} - 1, 1]$,
7. $[1, 2^{t+2} - 1, 2^{t+u+1} - 2^{t+1} - 1]$,
8. $[2^{t+2} - 1, 1, 2^{t+u+1} - 2^{t+1} - 1]$,
9. $[2^{t+2} - 1, 2^{t+u+1} - 2^{t+1} - 1, 1]$,
10. $[3, 2^{t+1} - 3, 2^{t+u+1} - 1]$,
11. $[3, 2^{t+u+1} - 1, 2^{t+1} - 3]$,
12. $[2^{t+u+1} - 1, 3, 2^{t+1} - 3]$,
13. $[3, 2^{t+1} - 1, 2^{t+u+1} - 3]$,
14. $[3, 2^{t+u+1} - 3, 2^{t+1} - 1]$,
15. $[2^{t+1} - 1, 3, 2^{t+u+1} - 3]$,
16. $[3, 2^{t+2} - 3, 2^{t+u+1} - 2^{t+1} - 1]$,
17. $[3, 2^{t+2} - 1, 2^{t+u+1} - 2^{t+1} - 3]$,
18. $[2^{t+2} - 1, 3, 2^{t+u+1} - 2^{t+1} - 3]$,
19. $[7, 2^{t+u+1} - 5, 2^{t+1} - 3]$,
20. $[7, 2^{t+2} - 5, 2^{t+u+1} - 2^{t+1} - 3]$.

For $t = 2$, $w_{u,2,1,21} = [7, 7, 2^{u+3} - 7]$.

For $t \geqslant 3$, $w_{u,t,1,21} = [7, 2^{t+1} - 5, 2^{t+u+1} - 3]$.

So, we easily obtain

Proposition 8.4.1. *For any integer $t \geqslant 2$ and $u \geqslant 2$,*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{t+u+1}+2^{t+1}-1} = 84.$$

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+u+1}+2^{t+1}-1}$. We have

Theorem 8.4.2. *For any $u \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+u+1}+2^{t+1}-1}$ is an $\mathbb{F}_2$-vector space with a basis consisting of all the classes represented by the monomials $a_{u,t,1,j}, j \geqslant 1$, which are determined as follows:*

For $t \geqslant 2$,

1. $(1, 1, 2^{t+1} - 2, 2^{t+u+1} - 1)$,
2. $(1, 1, 2^{t+u+1} - 1, 2^{t+1} - 2)$,
3. $(1, 2^{t+1} - 2, 1, 2^{t+u+1} - 1)$,
4. $(1, 2^{t+1} - 2, 2^{t+u+1} - 1, 1)$,
5. $(1, 2^{t+u+1} - 1, 1, 2^{t+1} - 2)$,
6. $(1, 2^{t+u+1} - 1, 2^{t+1} - 2, 1)$,
7. $(2^{t+u+1} - 1, 1, 1, 2^{t+1} - 2)$,
8. $(2^{t+u+1} - 1, 1, 2^{t+1} - 2, 1)$,
9. $(1, 1, 2^{t+1} - 1, 2^{t+u+1} - 2)$,
10. $(1, 1, 2^{t+u+1} - 2, 2^{t+1} - 1)$,
11. $(1, 2^{t+1} - 1, 1, 2^{t+u+1} - 2)$,
12. $(1, 2^{t+1} - 1, 2^{t+u+1} - 2, 1)$,
13. $(1, 2^{t+u+1} - 2, 1, 2^{t+1} - 1)$,
14. $(1, 2^{t+u+1} - 2, 2^{t+1} - 1, 1)$,
15. $(2^{t+1} - 1, 1, 1, 2^{t+u+1} - 2)$,
16. $(2^{t+1} - 1, 1, 2^{t+u+1} - 2, 1)$,
17. $(1, 1, 2^{t+2} - 2, 2^{t+u+1} - 2^{t+1} - 1)$,
18. $(1, 2^{t+2} - 2, 1, 2^{t+u+1} - 2^{t+1} - 1)$,
19. $(1, 2^{t+2} - 2, 2^{t+u+1} - 2^{t+1} - 1, 1)$,
20. $(1, 1, 2^{t+2} - 1, 2^{t+u+1} - 2^{t+1} - 2)$,
21. $(1, 2^{t+2} - 1, 1, 2^{t+u+1} - 2^{t+1} - 2)$,
22. $(1, 2^{t+2} - 1, 2^{t+u+1} - 2^{t+1} - 2, 1)$,
23. $(2^{t+2} - 1, 1, 1, 2^{t+u+1} - 2^{t+1} - 2)$,
24. $(2^{t+2} - 1, 1, 2^{t+u+1} - 2^{t+1} - 2, 1)$,
25. $(1, 2, 2^{t+1} - 3, 2^{t+u+1} - 1)$,
26. $(1, 2, 2^{t+u+1} - 1, 2^{t+1} - 3)$,
27. $(1, 2^{t+u+1} - 1, 2, 2^{t+1} - 3)$,
28. $(2^{t+u+1} - 1, 1, 2, 2^{t+1} - 3)$,
29. $(1, 2, 2^{t+1} - 1, 2^{t+u+1} - 3)$,
30. $(1, 2, 2^{t+u+1} - 3, 2^{t+1} - 1)$,
31. $(1, 2^{t+1} - 1, 2, 2^{t+u+1} - 3)$,
32. $(2^{t+1} - 1, 1, 2, 2^{t+u+1} - 3)$,
33. $(1, 3, 2^{t+1} - 4, 2^{t+u+1} - 1)$,
34. $(1, 3, 2^{t+u+1} - 1, 2^{t+1} - 4)$,
35. $(1, 2^{t+u+1} - 1, 3, 2^{t+1} - 4)$,
36. $(3, 1, 2^{t+1} - 4, 2^{t+u+1} - 1)$,
37. $(3, 1, 2^{t+u+1} - 1, 2^{t+1} - 4)$,
38. $(3, 2^{t+u+1} - 1, 1, 2^{t+1} - 4)$,
39. $(2^{t+u+1} - 1, 1, 3, 2^{t+1} - 4)$,
40. $(2^{t+u+1} - 1, 3, 1, 2^{t+1} - 4)$,
41. $(1, 3, 2^{t+1} - 3, 2^{t+u+1} - 2)$,
42. $(1, 3, 2^{t+u+1} - 2, 2^{t+1} - 3)$,
43. $(1, 2^{t+u+1} - 2, 3, 2^{t+1} - 3)$,
44. $(3, 1, 2^{t+1} - 3, 2^{t+u+1} - 2)$,
45. $(3, 1, 2^{t+u+1} - 2, 2^{t+1} - 3)$,
46. $(3, 2^{t+1} - 3, 1, 2^{t+u+1} - 2)$,
47. $(3, 2^{t+1} - 3, 2^{t+u+1} - 2, 1)$,
48. $(1, 3, 2^{t+1} - 2, 2^{t+u+1} - 3)$,
49. $(1, 3, 2^{t+u+1} - 3, 2^{t+1} - 2)$,
50. $(1, 2^{t+1} - 2, 3, 2^{t+u+1} - 3)$,
51. $(3, 1, 2^{t+1} - 2, 2^{t+u+1} - 3)$,
52. $(3, 1, 2^{t+u+1} - 3, 2^{t+1} - 2)$,
53. $(3, 2^{t+u+1} - 3, 1, 2^{t+1} - 2)$,
54. $(3, 2^{t+u+1} - 3, 2^{t+1} - 2, 1)$,
55. $(1, 3, 2^{t+1} - 1, 2^{t+u+1} - 4)$,
56. $(1, 3, 2^{t+u+1} - 4, 2^{t+1} - 1)$,
57. $(1, 2^{t+1} - 1, 3, 2^{t+u+1} - 4)$,
58. $(3, 1, 2^{t+1} - 1, 2^{t+u+1} - 4)$,
59. $(3, 1, 2^{t+u+1} - 4, 2^{t+1} - 1)$,
60. $(3, 2^{t+1} - 1, 1, 2^{t+u+1} - 4)$,
61. $(2^{t+1} - 1, 1, 3, 2^{t+u+1} - 4)$,
62. $(2^{t+1} - 1, 3, 1, 2^{t+u+1} - 4)$,
63. $(1, 2, 2^{t+2} - 3, 2^{t+u+1} - 2^{t+1} - 1)$,
64. $(1, 2, 2^{t+2} - 1, 2^{t+u+1} - 2^{t+1} - 3)$,
65. $(1, 2^{t+2} - 1, 2, 2^{t+u+1} - 2^{t+1} - 3)$,
66. $(2^{t+2} - 1, 1, 2, 2^{t+u+1} - 2^{t+1} - 3)$,
67. $(1, 3, 2^{t+2} - 4, 2^{t+u+1} - 2^{t+1} - 1)$,
68. $(3, 1, 2^{t+2} - 4, 2^{t+u+1} - 2^{t+1} - 1)$,
69. $(1, 3, 2^{t+2} - 3, 2^{t+u+1} - 2^{t+1} - 2)$,
70. $(3, 1, 2^{t+2} - 3, 2^{t+u+1} - 2^{t+1} - 2)$,

71. $(3, 2^{t+2}-3, 1, 2^{t+u+1}-2^{t+1}-2)$,
72. $(3, 2^{t+2}-3, 2^{t+u+1}-2^{t+1}-2, 1)$,
73. $(1, 3, 2^{t+2}-2, 2^{t+u+1}-2^{t+1}-3)$,
74. $(1, 2^{t+2}-2, 3, 2^{t+u+1}-2^{t+1}-3)$,
75. $(3, 1, 2^{t+2}-2, 2^{t+u+1}-2^{t+1}-3)$,
76. $(1, 3, 2^{t+2}-1, 2^{t+u+1}-2^{t+1}-4)$,
77. $(1, 2^{t+2}-1, 3, 2^{t+u+1}-2^{t+1}-4)$,
78. $(3, 1, 2^{t+2}-1, 2^{t+u+1}-2^{t+1}-4)$,
79. $(3, 2^{t+2}-1, 1, 2^{t+u+1}-2^{t+1}-4)$,
80. $(2^{t+2}-1, 1, 3, 2^{t+u+1}-2^{t+1}-4)$,
81. $(2^{t+2}-1, 3, 1, 2^{t+u+1}-2^{t+1}-4)$,
82. $(1, 6, 2^{t+u+1}-5, 2^{t+1}-3)$,
83. $(3, 2^{t+1}-3, 2, 2^{t+u+1}-3)$,
84. $(3, 2^{t+u+1}-3, 2, 2^{t+1}-3)$,
85. $(1, 7, 2^{t+u+1}-5, 2^{t+1}-4)$,
86. $(7, 1, 2^{t+u+1}-5, 2^{t+1}-4)$,
87. $(7, 2^{t+u+1}-5, 1, 2^{t+1}-4)$,
88. $(1, 7, 2^{t+u+1}-6, 2^{t+1}-3)$,
89. $(7, 1, 2^{t+u+1}-6, 2^{t+1}-3)$,
90. $(3, 3, 2^{t+1}-4, 2^{t+u+1}-3)$,
91. $(3, 3, 2^{t+u+1}-3, 2^{t+1}-4)$,
92. $(3, 2^{t+u+1}-3, 3, 2^{t+1}-4)$,
93. $(3, 3, 2^{t+1}-3, 2^{t+u+1}-4)$,
94. $(3, 3, 2^{t+u+1}-4, 2^{t+1}-3)$,
95. $(3, 2^{t+1}-3, 3, 2^{t+u+1}-4)$,
96. $(1, 6, 2^{t+2}-5, 2^{t+u+1}-2^{t+1}-3)$,
97. $(3, 2^{t+2}-3, 2, 2^{t+u+1}-2^{t+1}-3)$,
98. $(3, 4, 2^{t+u+1}-5, 2^{t+1}-3)$,
99. $(1, 7, 2^{t+2}-6, 2^{t+u+1}-2^{t+1}-3)$,
100. $(7, 1, 2^{t+2}-6, 2^{t+u+1}-2^{t+1}-3)$,
101. $(1, 7, 2^{t+2}-5, 2^{t+u+1}-2^{t+1}-4)$,
102. $(7, 1, 2^{t+2}-5, 2^{t+u+1}-2^{t+1}-4)$,
103. $(7, 2^{t+2}-5, 1, 2^{t+u+1}-2^{t+1}-4)$,
104. $(3, 3, 2^{t+2}-4, 2^{t+u+1}-2^{t+1}-3)$,
105. $(3, 3, 2^{t+2}-3, 2^{t+u+1}-2^{t+1}-4)$,
106. $(3, 2^{t+2}-3, 3, 2^{t+u+1}-2^{t+1}-4)$,
107. $(3, 7, 2^{t+u+1}-7, 2^{t+1}-4)$,
108. $(7, 3, 2^{t+u+1}-7, 2^{t+1}-4)$,
109. $(3, 4, 2^{t+2}-5, 2^{t+u+1}-2^{t+1}-3)$,
110. $(3, 5, 2^{t+2}-6, 2^{t+u+1}-2^{t+1}-3)$,
111. $(3, 5, 2^{t+2}-5, 2^{t+u+1}-2^{t+1}-4)$,
112. $(3, 7, 2^{t+2}-7, 2^{t+u+1}-2^{t+1}-4)$,
113. $(7, 3, 2^{t+2}-7, 2^{t+u+1}-2^{t+1}-4)$.

For $t = 2$,

114. $(1, 6, 7, 2^{u+3}-7)$,
115. $(1, 7, 6, 2^{u+3}-7)$,
116. $(7, 1, 6, 2^{u+3}-7)$,
117. $(1, 7, 7, 2^{u+3}-8)$,
118. $(7, 1, 7, 2^{u+3}-8)$,
119. $(7, 7, 1, 2^{u+3}-8)$,
120. $(3, 4, 7, 2^{u+3}-7)$,
121. $(3, 5, 6, 2^{u+3}-7)$,
122. $(3, 5, 7, 2^{u+3}-8)$,
123. $(3, 5, 2^{u+3}-5, 4)$,
124. $(3, 5, 2^{u+3}-6, 5)$,
125. $(3, 7, 5, 2^{u+3}-8)$,
126. $(7, 3, 5, 2^{u+3}-8)$,
127. $(3, 7, 8, 2^{u+3}-11)$,
128. $(7, 3, 8, 2^{u+3}-11)$,
129. $(3, 4, 1, 2^{u+3}-1)$,
130. $(3, 4, 2^{u+3}-1, 1)$,
131. $(3, 2^{u+3}-1, 4, 1)$,
132. $(2^{u+3}-1, 3, 4, 1)$,
133. $(3, 7, 2^{u+3}-4, 1)$,
134. $(7, 3, 2^{u+3}-4, 1)$,
135. $(7, 2^{u+3}-5, 4, 1)$,
136. $(3, 4, 3, 2^{u+3}-3)$,
137. $(7, 7, 2^{u+3}-8, 1)$,
138. $(3, 7, 4, 2^{u+3}-7)$,
139. $(7, 3, 4, 2^{u+3}-7)$,
140. $(7, 7, 8, 2^{u+3}-15)$,
141. $(7, 7, 9, 2^{u+3}-16)$.

For $t \geqslant 3$,

114. $(1, 6, 2^{t+1}-5, 2^{t+u+1}-3)$,
115. $(1, 7, 2^{t+1}-6, 2^{t+u+1}-3)$,
116. $(7, 1, 2^{t+1}-6, 2^{t+u+1}-3)$,
117. $(1, 7, 2^{t+1}-5, 2^{t+u+1}-4)$,
118. $(7, 1, 2^{t+1}-5, 2^{t+u+1}-4)$,
119. $(7, 2^{t+1}-5, 1, 2^{t+u+1}-4)$,
120. $(3, 4, 2^{t+1}-5, 2^{t+u+1}-3)$,
121. $(3, 5, 2^{t+1}-6, 2^{t+u+1}-3)$,

122. $(3,5,2^{t+1}-5,2^{t+u+1}-4)$, 123. $(3,5,2^{t+u+1}-5,2^{t+1}-4)$,
124. $(3,5,2^{t+u+1}-6,2^{t+1}-3)$, 125. $(3,7,2^{t+1}-7,2^{t+u+1}-4)$,
126. $(7,3,2^{t+1}-7,2^{t+u+1}-4)$.

We prove the theorem by proving the following propositions.

Proposition 8.4.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+u+1}+2^{t+1}-1}$ is generated by the elements listed in Theorem 8.4.2.*

The proof of this proposition is based on Theorem 2.5 and the following lemma.

Lemma 8.4.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&1&0&1\\1&1&0&0\\0&1&1&0\\0&1&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&1&0\\1&0&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&1&0\\0&1&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\0&1&0&1\\0&0&1&0\\0&0&1&0\end{pmatrix}$$
$$\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&0&1\\0&0&1&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&0&1&1\\1&0&1&0\\0&1&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\0&1&1&0\\0&1&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&1&0\\1&0&0&0\\0&0&0&1\end{pmatrix}$$
$$\begin{pmatrix}1&1&0&1\\1&0&1&0\\0&1&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&0&1&0\\0&1&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&0\\0&1&1&0\\0&1&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&0\\1&0&1&0\\1&0&0&0\\0&0&0&1\end{pmatrix}$$
$$\begin{pmatrix}1&1&0&1\\1&1&0&0\\0&1&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&0\\0&1&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&0\\1&0&1&0\\0&0&1&0\\0&0&0&1\end{pmatrix}$$
$$\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&1&0\\0&1&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&0\\1&0&0&1\\0&1&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&0\\1&0&1&0\\0&1&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&0\\1&0&0&1\\0&1&0&0\\0&0&1&0\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(3,15,20,1),(15,3,20,1),(7,11,20,1),(3,7,24,5),(7,3,24,5),$$
$$(3,4,15,17),(3,15,4,17),(15,3,4,17),(3,5,14,17),(3,5,15,16),$$
$$(3,15,5,16),(15,3,5,16),(3,7,12,17),(7,3,12,17),(3,7,13,16),$$

$$(7,3,13,16), (7,11,4,17), (7,11,17,4), (7,11,5,16), (7,11,16,5).$$

For simplicity, we prove the lemma for the matrices associated to the monomials

$$(3,15,20,1), (7,11,20,1), (7,3,24,5), (3,15,4,17), (3,5,14,17),$$
$$(15,3,5,16), (7,3,12,17), (7,3,13,16), (7,11,17,4), (7,11,5,16).$$

By a direct computation, we have

$$\begin{aligned}
(3,15,20,1) &= Sq^1\big((3,15,19,1) + (3,19,15,1)\big) + Sq^2\big((2,15,19,1)\\
&\quad + (2,19,15,1) + (5,15,15,2)\big) + Sq^4(3,15,15,2)\\
&\quad + Sq^8\big((3,9,15,4) + (3,12,15,1)\big) + (2,15,21,1) + (2,21,15,1)\\
&\quad + (3,15,17,4) + (3,9,23,4) + (3,12,23,1) \mod \mathcal{L}_4(3;2;2;1;1),\\
(7,11,20,1) &= Sq^1(7,13,17,1) + Sq^2\big((7,11,18,1) + (7,14,15,1)\\
&\quad + (7,11,17,2)\big) + Sq^4(5,14,15,1) + Sq^8(7,8,15,1) + (5,18,15,1)\\
&\quad + (5,14,19,1) + (7,8,23,1) + (7,11,17,4) \mod \mathcal{L}_4(3;2;2;1;1),\\
(7,3,24,5) &= Sq^1\big((7,3,23,5) + (7,3,19,9)\big) + Sq^2\big((7,2,23,5)\\
&\quad + (7,5,19,6) + (7,5,15,10) + (7,2,19,9)\big) + Sq^4\big((4,3,23,5)\\
&\quad + (5,2,23,5) + (11,3,15,6) + (5,3,21,6) + (5,3,15,12)\\
&\quad + (3,5,15,12)\big) + Sq^8(7,3,15,6) + (4,3,27,5) + (4,3,23,9)\\
&\quad + (7,2,25,5) + (5,2,27,5) + (5,2,23,9) + (7,3,21,8)\\
&\quad + (7,3,17,12) + (5,3,25,6) + (5,3,21,10) + (5,3,19,12)\\
&\quad + (5,3,15,16) + (7,3,20,9) + (7,2,21,9) \mod \mathcal{L}_4(3;2;2;1;1),\\
(3,15,4,17) &= Sq^1\big((3,19,1,15) + (3,15,1,19)\big) + Sq^2\big((5,15,2,15)\\
&\quad + (2,19,1,15) + (2,15,1,19)\big) + Sq^4(3,15,2,15)\\
&\quad + Sq^8\big((3,12,1,15) + (3,9,4,15)\big) + (2,21,1,15) + (2,15,1,21)\\
&\quad + (3,12,1,23) + (3,9,4,23) + (3,15,1,20) \mod \mathcal{L}_4(3;2;2;1;1),\\
(3,5,14,17) &= Sq^1\big((3,3,17,15) + (3,3,13,19)\big) + Sq^2\big((5,3,14,15)\\
&\quad + (2,3,17,15) + (2,3,13,19)\big) + Sq^4(3,3,14,15) + Sq^8(3,5,8,15)\\
&\quad + (3,4,17,15) + (2,5,17,15) + (3,4,13,19) + (3,3,13,20)\\
&\quad + (2,5,13,19) + (2,3,13,21) + (3,5,8,23) \mod \mathcal{L}_4(3;2;2;1;1),\\
(15,3,5,16) &= Sq^1\big((19,3,3,13) + (15,3,3,17)\big)\\
&\quad + Sq^2\big((15,5,3,14) + (15,2,3,17)\big) + Sq^4(15,3,3,14)\\
&\quad + Sq^8\big((9,3,5,14) + (12,3,3,13) + (11,4,3,13) + (11,3,4,13)\big)\\
&\quad + (9,3,5,22) + (12,3,3,21) + (11,3,4,21) + (11,4,3,21)\\
&\quad + (15,3,4,17) + (15,2,5,17) \mod \mathcal{L}_4(3;2;2;1;1),
\end{aligned}$$

$$\begin{aligned}(7,3,12,17) &= Sq^1\big((7,3,5,23)+(7,3,9,19)\big)+Sq^2\big((7,5,10,15)\\ &\quad+(7,5,6,19)+(7,2,9,19)+(7,2,5,23)\big)+Sq^4\big((11,3,6,15)\\ &\quad+(5,3,12,15)+(5,3,6,21)+(4,3,5,23)+(5,2,5,23)\big)\\ &\quad+Sq^8(7,3,6,15)+(7,3,8,21)+(5,3,16,15)+(5,3,12,19)\\ &\quad+(5,3,10,21)+(5,3,6,25)+(7,3,9,20)+(7,2,9,21)\\ &\quad+(7,3,5,24)+(4,3,9,23)+(4,3,5,27)+(7,2,5,25)\\ &\quad+(5,2,9,23)+(5,2,5,27)\mod \mathcal{L}_4(3;2;2;1;1),\\ (7,11,17,4) &= Sq^1(7,13,13,5)+Sq^2\big((7,7,19,4)+(7,11,13,6)\\ &\quad+(3,11,13,10)+(7,14,11,5)+(3,14,11,9)+(7,11,14,5)\\ &\quad+(3,11,14,9)\big)+Sq^4\big((11,7,13,4)+(5,7,19,4)+(5,11,13,6)\\ &\quad+(3,13,13,6)+(3,7,13,12)+(5,14,11,5)+(3,14,13,5)\\ &\quad+(5,11,14,5)+(3,13,14,5)\big)+Sq^8\big((7,7,13,4)+(7,8,11,5)\big)\\ &\quad+(7,7,17,8)+(7,9,19,4)+(5,11,19,4)+(5,7,19,8)\\ &\quad+(5,11,17,6)+(3,17,13,6)+(3,13,17,6)+(3,7,17,12)\\ &\quad+(3,7,13,16)+(7,8,19,5)+(5,18,11,5)+(3,18,13,5)\\ &\quad+(3,14,17,5)+(7,11,16,5)+(5,11,18,5)\\ &\quad+(3,17,14,5)+(3,13,18,5)\mod \mathcal{L}_4(3;2;2;1;1),\\ (7,11,5,16) &= Sq^1(7,13,5,13)+Sq^2\big((7,11,5,14)+(3,11,9,14)\\ &\quad+(7,14,3,13)+(7,11,6,13)+(3,11,10,13)+(7,7,2,21)\big)\\ &\quad+Sq^4\big((5,11,5,14)+(3,13,5,14)+(5,14,3,13)+(5,11,6,13)\\ &\quad+(3,7,12,13)+(3,13,6,13)+(11,7,4,13)+(5,7,2,21)\big)\\ &\quad+Sq^8\big((7,8,3,13)+(7,7,4,13)\big)+(5,11,5,18)+(3,17,5,14)\\ &\quad+(3,13,5,18)+(7,8,3,21)+(5,18,3,13)+(5,14,3,17)\\ &\quad+(5,11,6,17)+(3,7,16,13)+(3,7,12,17)+(3,17,6,13)\\ &\quad+(3,13,6,17)+(7,11,4,17)+(7,7,8,17)+(7,9,2,21)\\ &\quad+(5,11,2,21)+(5,7,2,25)\mod \mathcal{L}_4(3;2;2;1;1).\end{aligned}$$

The lemma is proved. □

Using the results in Section 7, Lemmas 5.10, 5.12, 5.13, 8.3.4, 8.3.5, 8.4.4 and Theorem 2.5, we obtain Proposition 8.4.3.

Now, we show that the elements listed in Theorem 8.4.2 are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+u+1}+2^{t+1}-1}$.

Proposition 8.4.5. *For any $u \geqslant 2$, the elements $[a_{u,2,1,j}], 1 \leqslant j \leqslant 141$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{u+3}+7}$.*

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{141} \gamma_j [a_{u,2,1,j}] = 0, \tag{8.4.5.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms f_1, f_2, f_3 to (8.4.5.1), we obtain

$$\begin{aligned}
&\gamma_{\{3,129\}} w_{u,2,1,3} + \gamma_{\{4,130\}} w_{u,2,1,4} + \gamma_{13} w_{u,2,1,5} + \gamma_{14} w_{u,2,1,6} + \gamma_{18} w_{u,2,1,8} \\
&\quad + \gamma_{19} w_{u,2,1,9} + \gamma_{25} w_{u,2,1,10} + \gamma_{26} w_{u,2,1,11} + \gamma_{43} w_{u,2,1,12} + \gamma_{29} w_{u,2,1,13} \\
&\quad + \gamma_{30} w_{u,2,1,14} + \gamma_{\{50,136\}} w_{u,2,1,15} + \gamma_{63} w_{u,2,1,16} + \gamma_{64} w_{u,2,1,17} \\
&\quad + \gamma_{74} w_{u,2,1,18} + \gamma_{\{82,98\}} w_{u,2,1,19} + \gamma_{\{96,109\}} w_{u,2,1,20} + \gamma_{\{114,120\}} w_{u,2,1,21} = 0, \\
&\gamma_{\{1,36\}} w_{u,2,1,3} + \gamma_{\{6,131\}} w_{u,2,1,4} + \gamma_{\{10,59\}} w_{u,2,1,5} + \gamma_{\{12,133,137\}} w_{u,2,1,6} \\
&\quad + \gamma_{\{17,68\}} w_{u,2,1,8} + \gamma_{22} w_{u,2,1,9} + \gamma_{33} w_{u,2,1,10} + \gamma_{27} w_{u,2,1,11} + \gamma_{\{42,94\}} w_{u,2,1,12} \\
&\quad + \gamma_{\{31,56,67\}} w_{u,2,1,13} + \gamma_{56} w_{u,2,1,14} + a_1 w_{u,2,1,15} + \gamma_{67} w_{u,2,1,16} + \gamma_{65} w_{u,2,1,17} \\
&\quad + a_2 w_{u,2,1,18} + \gamma_{88} w_{u,2,1,19} + \gamma_{\{99,127\}} w_{u,2,1,20} + a_3 w_{u,2,1,21} = 0, \\
&\gamma_{\{2,37\}} w_{u,2,1,3} + \gamma_{\{5,38\}} w_{u,2,1,4} + \gamma_{\{9,58,118\}} w_{u,2,1,5} + \gamma_{\{11,60,119\}} w_{u,2,1,6} \\
&\quad + \gamma_{\{20,78\}} w_{u,2,1,8} + \gamma_{\{21,79\}} w_{u,2,1,9} + \gamma_{34} w_{u,2,1,10} + \gamma_{35} w_{u,2,1,11} \\
&\quad + \gamma_{\{41,93,95,126\}} w_{u,2,1,12} + a_4 w_{u,2,1,13} + \gamma_{\{55,117,122\}} w_{u,2,1,14} \\
&\quad + a_5 w_{u,2,1,15} + \gamma_{76} w_{u,2,1,16} + \gamma_{77} w_{u,2,1,17} + a_6 w_{u,2,1,18} \\
&\quad + \gamma_{\{57,77,92,106,125\}} w_{u,2,1,19} + \gamma_{101} w_{u,2,1,20} + a_7 w_{u,2,1,21} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{48,56,67,90,140\}}, \quad a_2 = \gamma_{\{73,104,128,140\}}, \quad a_3 = \gamma_{\{56,67,88,99,115,127,138,140\}} \\
&a_4 = \gamma_{\{55,76,77,85,117,122\}}, \quad a_5 = \gamma_{\{49,55,76,91,106,112,122,123,141\}}, \\
&a_6 = \gamma_{\{69,105,106,111,112,141\}}, \quad a_7 = \gamma_{\{55,57,76,77,101,107,112,122,125,141\}}.
\end{aligned}$$

Computing from these equalities, we obtain

$$\begin{cases}
a_i = 0, i = 1,2,3,4,5,6,7, \\
\gamma_j = 0, \ j = 13,14,18,19,22,25,26,27,29,30,31, \\
33,34,35,43,56,63,64,65,67,74,76,77,88,101, \\
\gamma_{\{3,129\}} = \gamma_{\{4,130\}} = \gamma_{\{50,136\}} = \gamma_{\{82,98\}} = \gamma_{\{114,120\}} = 0, \\
\gamma_{\{96,109\}} = \gamma_{\{1,36\}} = \gamma_{\{6,131\}} = \gamma_{\{10,59\}} = \gamma_{\{17,68\}} = 0, \\
\gamma_{\{12,133,137\}} = \gamma_{\{42,94\}} = \gamma_{\{99,127\}} = \gamma_{\{2,37\}} = \gamma_{\{5,38\}} = 0, \\
\gamma_{\{9,58,118\}} = \gamma_{\{11,60,119\}} = \gamma_{\{20,78\}} = \gamma_{\{41,93,95,126\}} = 0, \\
\gamma_{\{21,79\}} = \gamma_{\{55,117,122\}} = \gamma_{\{57,92,106,125\}} = 0.
\end{cases} \tag{8.4.5.2}$$

With the aid of (8.4.5.2), the homomorphisms f_4, f_5, f_6 send (8.4.5.1) to

$$\begin{aligned}
&\gamma_{\{1,3\}} w_{u,2,1,1} + \gamma_{10} w_{u,2,1,2} + \gamma_{\{16,134,135,137\}} w_{u,2,1,4} + \gamma_{\{8,132\}} w_{u,2,1,6} \\
&\quad + \gamma_{17} w_{u,2,1,7} + \gamma_{24} w_{u,2,1,9} + \gamma_{\{36,129\}} w_{u,2,1,10} + \gamma_{\{45,84,94,98,124\}} w_{u,2,1,11}
\end{aligned}$$

$$\begin{aligned}
&+ \gamma_{28}w_{u,2,1,12} + \gamma_{\{51,83,90,136,140\}}w_{u,2,1,13} + \gamma_{59}w_{u,2,1,14} + \gamma_{32}w_{u,2,1,15}\\
&+ \gamma_{68}w_{u,2,1,16} + a_8w_{u,2,1,17} + \gamma_{66}w_{u,2,1,18} + \gamma_{89}w_{u,2,1,19}\\
&+ \gamma_{\{100,128\}}w_{u,2,1,20} + \gamma_{\{116,139,140\}}w_{u,2,1,21} = 0,\\
\gamma_{\{2,4\}}&w_{u,2,1,1} + \gamma_{\{9,55,114,117\}}w_{u,2,1,2} + \gamma_{\{15,62,87,103,119\}}w_{u,2,1,4} + \gamma_{\{7,40\}}w_{u,2,1,6}\\
&+ \gamma_{20}w_{u,2,1,7} + \gamma_{\{23,81\}}w_{u,2,1,9} + \gamma_{\{37,130\}}w_{u,2,1,10} + \gamma_{\{44,93,125\}}w_{u,2,1,11}\\
&+ \gamma_{39}w_{u,2,1,12} + \gamma_{\{52,91,113,118,141\}}w_{u,2,1,13} + \gamma_{\{58,118,120,122\}}w_{u,2,1,14}\\
&+ \gamma_{\{80,86\}}w_{u,2,1,15} + \gamma_{78}w_{u,2,1,16} + \gamma_{\{70,105,113,141\}}w_{u,2,1,17} + \gamma_{80}w_{u,2,1,18}\\
&+ \gamma_{\{61,80,126\}}w_{u,2,1,19} + \gamma_{102}w_{u,2,1,20} + \gamma_{\{108,113,118,141\}}w_{u,2,1,21} = 0,\\
a_9&w_{u,2,1,1} + \gamma_{\{5,6\}}w_{u,2,1,2} + a_{10}w_{u,2,1,3} + \gamma_{\{7,8,28,39\}}w_{u,2,1,5} + \gamma_{21}w_{u,2,1,7}\\
&+ \gamma_{\{23,24,66,80\}}w_{u,2,1,8} + a_{11}w_{u,2,1,10} + \gamma_{\{38,131\}}w_{u,2,1,11} + \gamma_{\{40,132\}}w_{u,2,1,12}\\
&+ a_{12}w_{u,2,1,13} + \gamma_{\{53,54,84,92\}}w_{u,2,1,14} + a_{13}w_{u,2,1,15} + \gamma_{\{71,72,97,106\}}w_{u,2,1,16}\\
&+ \gamma_{79}w_{u,2,1,17} + \gamma_{81}w_{u,2,1,18} + \gamma_{\{87,135\}}w_{u,2,1,19}\\
&+ \gamma_{103}w_{u,2,1,20} + \gamma_{\{119,137,140,141\}}w_{u,2,1,21} = 0,
\end{aligned}$$

where

$$\begin{aligned}
a_8 &= \gamma_{\{75,97,104,109,110,127,140\}}, \quad a_9 = \gamma_{\{11,12,57,85,99,115,117\}},\\
a_{10} &= \gamma_{\{15,16,32,61,86,89,100,102,116,118\}}, \quad a_{11} = \gamma_{\{46,47,83,95,110,111,121,122,123,124\}},\\
a_{12} &= \gamma_{\{60,107,112,125,127,133,138\}}, \quad a_{13} = \gamma_{\{62,108,113,126,128,134,139\}}.
\end{aligned}$$

From these equalities, we get

$$\text{(8.4.5.3)} \quad \begin{cases}
a_i = 0,\ i = 8, 9, \ldots, 13,\ \gamma_j = 0,\ j = 10, 17, 20, 21, 23,\\
24, 28, 32, 39, 59, 66, 68, 78, 79, 80, 81, 86, 89, 102, 103,\\
\gamma_{\{1,3\}} = \gamma_{\{16,134,135,137\}} = \gamma_{\{8,132\}} = \gamma_{\{36,129\}} = 0,\\
\gamma_{\{45,84,94,98,124\}} = \gamma_{\{51,83,90,136,140\}} = \gamma_{\{116,139,140\}} = 0,\\
\gamma_{\{100,128\}} = \gamma_{\{2,4\}} = \gamma_{\{9,55,114,117\}} = \gamma_{\{15,62,87,119\}} = 0,\\
\gamma_{\{7,40\}} = \gamma_{\{37,130\}} = \gamma_{\{44,93,125\}} = \gamma_{\{52,91,113,118,141\}} = 0,\\
\gamma_{\{58,118,120,122\}} = \gamma_{\{70,105,113,141\}} = \gamma_{\{108,113,118,141\}} = 0,\\
\gamma_{\{61,126\}} = \gamma_{\{5,6\}} = \gamma_{\{7,8\}} = \gamma_{\{38,131\}} = \gamma_{\{53,54,84,92\}} = 0,\\
\gamma_{\{40,132\}} = \gamma_{\{71,72,97,106\}} = \gamma_{\{87,135\}} = \gamma_{\{119,137,140,141\}} = 0.
\end{cases}$$

With the aid of (8.4.5.2) and (8.4.5.3), the homomorphisms g_1, g_2 send (8.4.5.1) to

$$\begin{aligned}
&a_{14}w_{u,2,1,4} + \gamma_{\{54,87\}}w_{u,2,1,6} + \gamma_{72}w_{u,2,1,9} + \gamma_{45}w_{u,2,1,11} + \gamma_{84}w_{u,2,1,12}\\
&+ \gamma_{\{48,51,90,140\}}w_{u,2,1,13} + \gamma_{83}w_{u,2,1,15} + \gamma_{\{73,75,104,140\}}w_{u,2,1,17}\\
&+ \gamma_{97}w_{u,2,1,18} + \gamma_{124}w_{u,2,1,19} + \gamma_{110}w_{u,2,1,20} + a_{15}w_{u,2,1,21} = 0,
\end{aligned}$$

$$\begin{aligned}&\gamma_{\{9,58,118\}}w_{u,2,1,2}+a_{16}w_{u,2,1,4}+\gamma_{\{53,87\}}w_{u,2,1,6}+\gamma_{71}w_{u,2,1,9}\\&\quad+\gamma_{\{41,44,50,93,95\}}w_{u,2,1,11}+\gamma_{92}w_{u,2,1,12}+a_{17}w_{u,2,1,13}\\&\quad+\gamma_{\{55,58,114,117,118\}}w_{u,2,1,14}+\gamma_{\{85,106,123\}}w_{u,2,1,15}+a_{18}w_{u,2,1,17}\\&\quad+\gamma_{106}w_{u,2,1,18}+\gamma_{\{57,92,95,125\}}w_{u,2,1,19}+\gamma_{111}w_{u,2,1,20}+a_{19}w_{u,2,1,21}=0,\end{aligned}$$

where

$$\begin{aligned}&a_{14}=\gamma_{\{12,16,47,133,134,137\}},\quad a_{15}=\gamma_{\{115,116,121,138,139,140\}},\\&a_{16}=\gamma_{\{11,15,46,60,62,119\}},\quad a_{17}=\gamma_{\{49,52,82,91,106,112,113,117,118,122,123,141\}},\\&a_{18}=\gamma_{\{69,70,96,105,106,111,112,113,141\}},\quad a_{19}=\gamma_{\{92,106,107,108,112,113,117,118,122,141\}}.\end{aligned}$$

From these equalities, we get

$$\text{(8.4.5.4)}\qquad\begin{cases}a_i=0,\ i=14,15,\dots,19,\\ \gamma_{\{54,87\}}=\gamma_{\{48,51,90,140\}}=\gamma_{\{73,75,104,140\}}=0,\\ \gamma_{\{9,58,118\}}=\gamma_{\{53,87\}}=\gamma_{\{41,44,50,93,95\}}=0,\\ \gamma_{\{55,58,114,117,118\}}=\gamma_{\{85,123\}}=\gamma_{\{57,95,125\}}=0.\end{cases}$$

With the aid of (8.4.5.2), (8.4.5.3) and (8.4.5.4), the homomorphisms g_3, g_4 send (8.4.5.1) to

$$\begin{aligned}&\gamma_{\{11,60,119\}}w_{u,2,1,2}+a_{20}w_{u,2,1,4}+\gamma_{\{12,133,137\}}w_{u,2,1,5}+\gamma_{\{16,52\}}w_{u,2,1,6}\\&\quad+\gamma_{\{70,75,100\}}w_{u,2,1,9}+a_{21}w_{u,2,1,11}+\gamma_{\{47,91,108,134\}}w_{u,2,1,12}\\&\quad+a_{22}w_{u,2,1,13}+a_{23}w_{u,2,1,14}+a_{24}w_{u,2,1,15}+a_{25}w_{u,2,1,17}\\&\quad+a_{26}w_{u,2,1,18}+a_{27}w_{u,2,1,19}+\gamma_{112}w_{u,2,1,20}+a_{28}w_{u,2,1,21}=0,\\&\gamma_{\{15,62,87,119\}}w_{u,2,1,2}+a_{29}w_{u,2,1,4}+\gamma_{\{16,87,134,137\}}w_{u,2,1,5}\\&\quad+\gamma_{\{12,42,49,82,85\}}w_{u,2,1,6}+\gamma_{\{69,73,96,99\}}w_{u,2,1,9}+a_{30}w_{u,2,1,11}\\&\quad+a_{31}w_{u,2,1,12}+a_{32}w_{u,2,1,13}+a_{33}w_{u,2,1,14}+a_{34}w_{u,2,1,15}+a_{35}w_{u,2,1,17}\\&\quad+a_{36}w_{u,2,1,18}+a_{37}w_{u,2,1,19}+\gamma_{113}w_{u,2,1,20}+a_{38}w_{u,2,1,21}=0,\end{aligned}$$

where

$$\begin{aligned}&a_{20}=\gamma_{\{9,15,44,51,58,61,116,118\}},\quad a_{21}=\gamma_{\{41,46,61,62,90,93,95,139\}},\\&a_{22}=\gamma_{\{53,69,85,100,104,105,112,113,115,117,119,125,133,137,138,140,141\}},\\&a_{23}=\gamma_{\{57,60,85,99,115,117,119,133,137\}},\quad a_{25}=\gamma_{\{69,99,100,104,105,112,113,140,141\}},\\&a_{24}=\gamma_{\{54,73,85,100,104,105,107,113,133,140,141\}},\ a_{26}=\gamma_{\{73,100,104,105,113,140,141\}},\\&a_{27}=\gamma_{\{42,48,49,55,61,62,69,73,90,91,93,108,121,122,123,134,139\}},\\&a_{28}=\gamma_{\{42,49,69,85,91,100,104,105,107,108,112,113,115,117,119,123,125,133,134,138,140,141\}},\\&a_{29}=\gamma_{\{9,11,41,48,50,55,57,114,115,117\}},\quad a_{30}=\gamma_{\{44,46,53,60,90,93,125,138\}},\\&a_{32}=\gamma_{\{61,70,99,104,105,112,113,116,118,119,134,137,139,140,141\}},\\&a_{31}=\gamma_{\{42,47,54,82,91,107,123,133\}},\quad a_{33}=\gamma_{\{61,62,100,116,118,119,134,137\}},\\&a_{34}=\gamma_{\{75,96,99,104,105,108,112,134,140,141\}},\quad a_{35}=\gamma_{\{70,99,100,104,105,112,113,140,141\}},\\&a_{36}=\gamma_{\{75,96,99,104,105,112,140,141\}},\quad a_{38}=\gamma_{\{42,52,53,61,70,87,91,99,104,105\}},\end{aligned}$$

$$a_{37} = \gamma_{\{42,50,51,52,53,54,58,60,70,75,90,91,93,95,96,107,114,121,122,125,133,138\}}.$$

These equalities imply

$$\begin{cases} \gamma_j = 0,\ i = 112, 113,\ a_i = 0,\ i = 20, 21, \ldots, 38, \\ \gamma_{\{11,60,119\}} = \gamma_{\{12,133,137\}} = \gamma_{\{16,52\}} = 0, \\ \gamma_{\{70,75,100\}} = \gamma_{\{47,91,108,134\}} = \gamma_{\{15,62,87,119\}} = 0, \\ \gamma_{\{16,87,134,137\}} = \gamma_{\{12,42,49,82,85\}} = \gamma_{\{69,73,96,99\}} = 0. \end{cases} \tag{8.4.5.5}$$

Substituting (8.4.5.2), (8.4.5.3), (8.4.5.4) and (8.4.5.5) into the relation (8.4.5.1), we obtain

$$\sum_{i=1}^{15} b_i[\theta_i], \tag{8.4.5.6}$$

where $b_1 = \gamma_1, b_2 = \gamma_2, b_3 = \gamma_5, b_4 = \gamma_7, b_5 = \gamma_9, b_6 = \gamma_{11}, b_7 = \gamma_{12}, b_8 = \gamma_{15}, b_9 = \gamma_{16}, b_{10} = \gamma_{41}, b_{11} = \gamma_{42}, b_{12} = \gamma_{44}, b_{13} = \gamma_{46}, b_{14} = \gamma_{48}, b_{15} = \gamma_{53}$ and

$$\begin{aligned}
\theta_1 &= a_{u,2,1,1} + a_{u,2,1,3} + a_{u,2,1,36} + a_{u,2,1,129}, \\
\theta_2 &= a_{u,2,1,2} + a_{u,2,1,4} + a_{u,2,1,37} + a_{u,2,1,130}, \\
\theta_3 &= a_{u,2,1,5} + a_{u,2,1,6} + a_{u,2,1,38} + a_{u,2,1,131}, \\
\theta_4 &= a_{u,2,1,7} + a_{u,2,1,8} + a_{u,2,1,40} + a_{u,2,1,132}, \\
\theta_5 &= a_{u,2,1,9} + a_{u,2,1,58} + a_{u,2,1,114} + a_{u,2,1,120}, \\
\theta_6 &= a_{u,2,1,11} + a_{u,2,1,60} + a_{u,2,1,115} + a_{u,2,1,138}, \\
\theta_7 &= a_{u,2,1,12} + a_{u,2,1,49} + a_{u,2,1,55} + a_{u,2,1,107} + a_{u,2,1,117} + a_{u,2,1,133}, \\
\theta_8 &= a_{u,2,1,15} + a_{u,2,1,62} + a_{u,2,1,116} + a_{u,2,1,139}, \\
\theta_9 &= a_{u,2,1,16} + a_{u,2,1,52} + a_{u,2,1,58} + a_{u,2,1,108} + a_{u,2,1,118} + a_{u,2,1,134}, \\
\theta_{10} &= a_{u,2,1,41} + a_{u,2,1,57} + a_{u,2,1,73} + a_{u,2,1,93} \\
&\quad + a_{u,2,1,99} + a_{u,2,1,104} + a_{u,2,1,125} + a_{u,2,1,127}, \\
\theta_{11} &= a_{u,2,1,42} + a_{u,2,1,47} + a_{u,2,1,55} + a_{u,2,1,82} + a_{u,2,1,91} + a_{u,2,1,94} \\
&\quad + a_{u,2,1,98} + a_{u,2,1,105} + a_{u,2,1,107} + a_{u,2,1,108} + a_{u,2,1,114} + a_{u,2,1,120} \\
&\quad + a_{u,2,1,122} + a_{u,2,1,133} + a_{u,2,1,134} + a_{u,2,1,137} + a_{u,2,1,141}, \\
\theta_{12} &= a_{u,2,1,44} + a_{u,2,1,61} + a_{u,2,1,75} + a_{u,2,1,93} \\
\theta_{13} &= a_{u,2,1,46} + a_{u,2,1,60} + a_{u,2,1,62} + a_{u,2,1,90} + a_{u,2,1,104} \\
&\quad + a_{u,2,1,119} + a_{u,2,1,121} + a_{u,2,1,140} + a_{u,2,1,139} + a_{u,2,1,140}, \\
\theta_{14} &= a_{u,2,1,48} + a_{u,2,1,50} + a_{u,2,1,57} + a_{u,2,1,58} + a_{u,2,1,61} + a_{u,2,1,73} \\
&\quad + a_{u,2,1,75} + a_{u,2,1,93} + a_{u,2,1,99} + a_{u,2,1,100} + a_{u,2,1,104} + a_{u,2,1,105} \\
&\quad + a_{u,2,1,114} + a_{u,2,1,115} + a_{u,2,1,116} + a_{u,2,1,117} + a_{u,2,1,118} \\
&\quad + a_{u,2,1,120} + a_{u,2,1,121} + a_{u,2,1,122} + a_{u,2,1,125} + a_{u,2,1,126} \\
&\quad + a_{u,2,1,127} + a_{u,2,1,128} + a_{u,2,1,136} + a_{u,2,1,140} + a_{u,2,1,141},
\end{aligned}$$

$$\theta_{15} = a_{u,2,1,53} + a_{u,2,1,54} + a_{u,2,1,60} + a_{u,2,1,87} + a_{u,2,1,119} + a_{u,2,1,133} + a_{u,2,1,135} + a_{u,2,1,137}.$$

Now, we prove $b_i = 0$ for $i = 1, 2, \ldots, 15$. The proof is divided into 6 steps.

Step 1. First, we prove $b_1 = 0$ by showing that the polynomial $\theta = \theta_1 + \sum_{j=2}^{15} c_j\theta_j$ is non-hit for all $c_j \in \mathbb{F}_2, j = 2, 3, \ldots, 15$. Suppose the contrary that θ is hit. Then we have

$$\theta = \sum_{m=0}^{u+2} Sq^{2^m}(A_m),$$

for some polynomials $A_m, m = 0, 1, \ldots, u+2$. Let $(Sq^2)^3$ act on the both sides of this equality. Since $(Sq^2)^3Sq^1 = 0, (Sq^2)^3Sq^2 = 0$, we get

$$(Sq^2)^3(\theta) = \sum_{m=2}^{u+2} (Sq^2)^3 Sq^{2^m}(A_m).$$

It is not difficult to check that the monomial $(8, 4, 2, 2^{u+3} - 1)$ is a term of $(Sq^2)^3(\theta)$. This monomial is not a term of $\sum_{m=2}^{u+2}(Sq^2)^3Sq^{2^m}(A_m)$ for all polynomials $A_m, m = 2, 3, \ldots, u+2$. So, we have a contradiction.

By an argument analogous to the previous one, we get $b_2 = b_3 = b_4 = 0$. Then, the relation (8.4.5.6) becomes

$$\sum_{i=5}^{15} b_i[\theta_i] = 0. \tag{8.4.5.7}$$

Step 2. Under the homomorphism φ_4, the linear relation (8.4.5.7) is sent to

$$b_{15}[\theta_3] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.4.5.8}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.4.5.8) and by a same argument as given in Step 1, we get $b_{15} = \gamma_{53} = 0$.

Consider the homomorphism induced by the linear map $x_1 \mapsto x_1 + x_3, x_2 \mapsto x_2, x_3 \mapsto x_3, x_4 \mapsto x_4$. Under this homomorphism, the linear relation (8.4.5.7) is sent to

$$b_9[\theta_2] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.4.5.9}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.4.5.9) and by an analogous argument as given in Step 1, we get $b_9 = \gamma_{16} = 0$.

The homomorphism induced by the linear transformation $x_1 \mapsto x_1 + x_4, x_2 \mapsto x_2, x_3 \mapsto x_3, x_4 \mapsto x_4$ sends the linear relation (8.4.5.7) to

$$b_8[\theta_1] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.4.5.10}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.4.5.10) and by a same argument as given in Step 1, we get $b_8 = \gamma_{15} = 0$.

Applying the homomorphism induced by the linear transformation $x_1 \mapsto x_1, x_2 \mapsto x_2 + x_3, x_3 \mapsto x_3, x_4 \mapsto x_4$ to the linear relation (8.4.5.7), we get

$$b_7[\theta_2] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.4.5.11}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.4.5.11) and by a same argument as given in Step 1, we get $b_7 = \gamma_{12} = 0$.

Consider the homomorphism induced by the linear map $x_1 \mapsto x_1, x_2 \mapsto x_2 + x_4, x_3 \mapsto x_3, x_4 \mapsto x_4$. This homomorphism sends the linear relation (8.4.5.7) to

$$b_6[\theta_1] + \sum_{i=5}^{15} d_i[\theta_i] = 0, \tag{8.4.5.12}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.4.5.12) and by a same argument as given in Step 1, we get $b_6 = \gamma_{11} = 0$.

The homomorphism induced by the linear transformation $x_1 \mapsto x_1, x_2 \mapsto x_2, x_3 \mapsto x_3 + x_4, x_4 \mapsto x_4$ sends the linear relation (8.4.5.7) to

$$b_5[\theta_1] + \sum_{i=5}^{15} d_i[\theta_i] = 0. \tag{8.4.5.13}$$

for some $d_i \in \mathbb{F}_2, i = 5, 6, \ldots, 15$. Using (8.4.5.13) and by a same argument as given in Step 1, we get $b_5 = \gamma_9 = 0$.

Now, the relation (8.4.5.7) becomes

$$\sum_{i=10}^{14} b_i[\theta_i] = 0. \tag{8.4.5.14}$$

Step 3. Applying the homomorphism φ_4 to (8.4.5.14), we get

$$b_{13}[\theta_6] + b_{11}[\theta_7] + (b_{10} + b_{12} + b_{14})[\theta_{10}] + \sum_{i=11}^{14} b_i[\theta_i] = 0.$$

Using this equality and by a same argument as given in Step 2, we obtain $b_{11} = \gamma_{42} = 0, b_{13} = \gamma_{46} = 0$. Then, the relation (8.4.5.14) becomes

$$b_{10}[\theta_{10}] + b_{12}[\theta_{12}] + b_{14}[\theta_{14}] = 0. \tag{8.4.5.15}$$

Step 4. Applying the homomorphism φ_2 to the relation (8.4.5.15) we obtain

$$b_{10}[\theta_{10}] + b_{12}[\theta_{13}] + b_{14}[\theta_{14}] = 0.$$

Using this equality and by an argument analogous to Step 3, we get $b_{12} = \gamma_{44} = 0$. So, the relation (8.4.5.15) becomes

$$b_{10}[\theta_{10}] + b_{14}[\theta_{14}] = 0. \tag{8.4.5.16}$$

Step 5. Applying the homomorphism φ_1 to the relation (8.4.5.16) we obtain

$$b_{10}[\theta_{12}] + b_{14}[\theta_{14}] = 0.$$

Using this equality and by an argument analogous to Step 4, we get $b_{10} = \gamma_{41} = 0$. So, the relation (8.4.5.16) becomes

$$b_{14}[\theta_{14}] = 0. \tag{8.4.5.17}$$

Step 6. Applying the homomorphism φ_4 to the relation (8.4.5.17) we obtain

$$b_{14}[\theta_{10}] + b_{14}[\theta_{14}] = 0.$$

Using this equality and by an argument analogous to Step 5, we get $b_{14} = \gamma_{48} = 0$.

The proposition is completely proved. □

Proposition 8.4.6. *For any $t \geqslant 3$ and $u \geqslant 2$, the elements $[a_{u,t,1,j}], 1 \leqslant j \leqslant 126$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{u+t+1}+2^{t+1}-1}$.*

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{126} \gamma_j [a_{u,t,1,j}] = 0, \tag{8.4.6.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms f_1, f_2, f_3 to (8.4.6.1) and we have

$$\begin{aligned}
&\gamma_3 w_{u,t,1,3} + \gamma_4 w_{u,t,1,4} + \gamma_{13} w_{u,t,1,5} + \gamma_{14} w_{u,t,1,6} + \gamma_{18} w_{u,t,1,8} + \gamma_{19} w_{u,t,1,9} \\
&\quad + \gamma_{25} w_{u,t,1,10} + \gamma_{26} w_{u,t,1,11} + \gamma_{43} w_{u,t,1,12} + \gamma_{29} w_{u,t,1,13} + \gamma_{30} w_{u,t,1,14} \\
&\quad + \gamma_{50} w_{u,t,1,15} + \gamma_{63} w_{u,t,1,16} + \gamma_{64} w_{u,t,1,17} + \gamma_{74} w_{u,t,1,18} \\
&\quad + \gamma_{\{82,98\}} w_{u,t,1,19} + \gamma_{\{96,109\}} w_{u,t,1,20} + \gamma_{\{114,120\}} w_{u,t,1,21} = 0, \\
&\gamma_{\{1,36\}} w_{u,t,1,3} + \gamma_6 w_{u,t,1,4} + \gamma_{\{10,59\}} w_{u,t,1,5} + \gamma_{12} w_{u,t,1,6} \\
&\quad + \gamma_{\{17,68\}} w_{u,t,1,8} + \gamma_{22} w_{u,t,1,9} + \gamma_{33} w_{u,t,1,10} + \gamma_{27} w_{u,t,1,11} \\
&\quad + \gamma_{\{42,94\}} w_{u,t,1,12} + \gamma_{31} w_{u,t,1,13} + \gamma_{56} w_{u,t,1,14} + \gamma_{\{48,90\}} w_{u,t,1,15} \\
&\quad + \gamma_{67} w_{u,t,1,16} + \gamma_{65} w_{u,t,1,17} + \gamma_{\{73,104\}} w_{u,t,1,18} + \gamma_{88} w_{u,t,1,19} \\
&\quad + \gamma_{99} w_{u,t,1,20} + \gamma_{\{56,67,88,99,115\}} w_{u,t,1,21} = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_{\{2,37\}} w_{u,t,1,3} + \gamma_{\{5,38\}} w_{u,t,1,4} + \gamma_{\{9,58\}} w_{u,t,1,5} + \gamma_{\{11,60\}} w_{u,t,1,6} \\
&\quad + \gamma_{\{20,78\}} w_{u,t,1,8} + \gamma_{\{21,79\}} w_{u,t,1,9} + \gamma_{34} w_{u,t,1,10} + \gamma_{35} w_{u,t,1,11} \\
&\quad + \gamma_{\{41,93,95,122,125\}} w_{u,t,1,12} + \gamma_{57} w_{u,t,1,13} + \gamma_{55} w_{u,t,1,14} \\
&\quad + \gamma_{\{49,91,92,107,123\}} w_{u,t,1,15} + \gamma_{76} w_{u,t,1,16} + \gamma_{77} w_{u,t,1,17} \\
&\quad + \gamma_{\{69,105,106,111,112\}} w_{u,t,1,18} + \gamma_{\{57,77,92,106,117\}} w_{u,t,1,19} \\
&\quad + \gamma_{101} w_{u,t,1,20} + \gamma_{\{55,76,85,92,101,106,117\}} w_{u,t,1,21} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$(8.4.6.2)\qquad \begin{cases} \gamma_j = 0,\ j = 3,4,6,12,13,14,18,19,22,\\ 25,26,27,29,30,31,33,34,35,43,50,55,56,\\ 57,63,64,65,67,74,76,77,88,99,101,115,\\ \gamma_{\{114,120\}} = \gamma_{\{82,98\}} = \gamma_{\{96,109\}} = \gamma_{\{1,36\}} = 0,\\ \gamma_{\{10,59\}} = \gamma_{\{17,68\}} = \gamma_{\{42,94\}} = \gamma_{\{48,90\}} = 0,\\ \gamma_{\{73,104\}} = \gamma_{\{2,37\}} = \gamma_{\{5,38\}} = \gamma_{\{9,58\}} = 0,\\ \gamma_{\{11,60\}} = \gamma_{\{20,78\}} = \gamma_{\{41,93,95,122,125\}} = 0,\\ \gamma_{\{55,76,85,92,101,106,117\}} = \gamma_{\{49,91,92,107,123\}} = 0,\\ \gamma_{\{21,79\}} = \gamma_{\{69,105,106,111,112\}} = \gamma_{\{92,106,117\}} = 0. \end{cases}$$

With the aid of (8.4.6.2), the homomorphisms f_4, f_5, f_6 send (8.4.6.1) to

$$\begin{aligned}
&\gamma_1 w_{u,t,1,1} + \gamma_{10} w_{u,t,1,2} + \gamma_{16} w_{u,t,1,4} + \gamma_8 w_{u,t,1,6} + \gamma_{17} w_{u,t,1,7}\\
&\quad + \gamma_{24} w_{u,t,1,9} + \gamma_{36} w_{u,t,1,10} + \gamma_{\{45,84,94,98,124\}} w_{u,t,1,11} + \gamma_{28} w_{u,t,1,12}\\
&\quad + \gamma_{\{51,83,90,120,121\}} w_{u,t,1,13} + \gamma_{59} w_{u,t,1,14} + \gamma_{32} w_{u,t,1,15}\\
&\quad + \gamma_{68} w_{u,t,1,16} + \gamma_{\{75,97,104,109,110\}} w_{u,t,1,17} + \gamma_{66} w_{u,t,1,18}\\
&\quad + \gamma_{89} w_{u,t,1,19} + \gamma_{100} w_{u,t,1,20} + \gamma_{116} w_{u,t,1,21} = 0,\\
&\gamma_2 w_{u,t,1,1} + \gamma_9 w_{u,t,1,2} + \gamma_{\{15,62\}} w_{u,t,1,4} + \gamma_{\{7,40\}} w_{u,t,1,6} + \gamma_{20} w_{u,t,1,7}\\
&\quad + \gamma_{\{23,81\}} w_{u,t,1,9} + \gamma_{37} w_{u,t,1,10} + \gamma_{\{44,93,126\}} w_{u,t,1,11}\\
&\quad + \gamma_{39} w_{u,t,1,12} + \gamma_{\{52,91,108\}} w_{u,t,1,13} + \gamma_{58} w_{u,t,1,14} + \gamma_{61} w_{u,t,1,15}\\
&\quad + \gamma_{78} w_{u,t,1,16} + \gamma_{\{70,105,113\}} w_{u,t,1,17} + \gamma_{80} w_{u,t,1,18}\\
&\quad + \gamma_{\{61,80,118\}} w_{u,t,1,19} + \gamma_{102} w_{u,t,1,20} + \gamma_{\{61,80,86\}} w_{u,t,1,21} = 0,\\
&\gamma_{11} w_{u,t,1,1} + \gamma_5 w_{u,t,1,2} + \gamma_{\{15,16,32,61\}} w_{u,t,1,3} + \gamma_{38} w_{u,t,1,11}\\
&\quad + \gamma_{\{7,8,28,39\}} w_{u,t,1,5} + \gamma_{\{23,24,66,80\}} w_{u,t,1,8} + \gamma_{\{46,47,83,95\}} w_{u,t,1,10}\\
&\quad + \gamma_{21} w_{u,t,1,7} + \gamma_{40} w_{u,t,1,12} + \gamma_{60} w_{u,t,1,13} + \gamma_{\{53,54,84,92\}} w_{u,t,1,14}\\
&\quad + \gamma_{\{71,72,97,106\}} w_{u,t,1,16} + \gamma_{62} w_{u,t,1,15} + \gamma_{79} w_{u,t,1,17} + \gamma_{81} w_{u,t,1,18}\\
&\quad + \gamma_{87} w_{u,t,1,19} + \gamma_{103} w_{u,t,1,20} + \gamma_{119} w_{u,t,1,21} = 0.
\end{aligned}$$

From these equalities, we get

$$(8.4.6.3)\qquad \begin{cases} \gamma_j = 0,\ j = 1,2,5,7,8,9,10,11,15,16,17,20,21,\\ 23,24,28,32,36,37,38,39,40,58,59,60,61,62,66,68,\\ 78,79,80,81,86,87,89,100,102,103,116,118,119,\\ \gamma_{\{45,84,94,98,124\}} = \gamma_{\{44,93,126\}} = \gamma_{\{51,83,90,120,121\}} = 0,\\ \gamma_{\{52,91,108\}} = \gamma_{\{75,97,104,109,110\}} = \gamma_{\{70,105,113\}} = 0,\\ \gamma_{\{46,47,83,95\}} = \gamma_{\{53,54,84,92\}} = \gamma_{\{71,72,97,106\}} = 0. \end{cases}$$

With the aid of (8.4.6.2) and (8.4.6.3), the homomorphisms g_1, g_2 send (8.4.6.1) to

$$\begin{aligned}
&\gamma_{47}w_{u,t,1,4} + \gamma_{54}w_{u,t,1,6} + \gamma_{72}w_{u,t,1,9} + \gamma_{45}w_{u,t,1,11}\\
&\quad + \gamma_{84}w_{u,t,1,12} + \gamma_{51}w_{u,t,1,13} + \gamma_{83}w_{u,t,1,15} + \gamma_{75}w_{u,t,1,17}\\
&\quad + \gamma_{97}w_{u,t,1,18} + \gamma_{124}w_{u,t,1,19} + \gamma_{110}w_{u,t,1,20} + \gamma_{121}w_{u,t,1,21} = 0,\\
&\gamma_{46}w_{u,t,1,4} + \gamma_{53}w_{u,t,1,6} + \gamma_{71}w_{u,t,1,9} + a_1w_{u,t,1,11} + \gamma_{92}w_{u,t,1,12}\\
&\quad + a_2w_{u,t,1,13} + \gamma_{95}w_{u,t,1,15} + a_3w_{u,t,1,17} + \gamma_{106}w_{u,t,1,18}\\
&\quad + \gamma_{\{92,95,117,122\}}w_{u,t,1,19} + \gamma_{111}w_{u,t,1,20} + \gamma_{\{85,95,106,123\}}w_{u,t,1,21} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{41,44,93,95,114,122,125,126\}}, \quad a_2 = \gamma_{\{49,52,82,91,92,107,108,123\}},\\
&a_3 = \gamma_{\{69,70,96,105,106,111,112,113\}}.
\end{aligned}$$

From these equalities, we get

$$\text{(8.4.6.4)} \qquad \begin{cases} a_i = 0,\ i = 1,2,3,\ \gamma_j = 0,\ j = 45,46,47,51,53,54,\\ 71,72,75,83,84,92,95,97,106,110,111,121,124,\\ \gamma_{\{117,122\}} = 0. \end{cases}$$

With the aid of (8.4.6.2), (8.4.6.3) and (8.4.6.4), the homomorphisms g_3, g_4 send (8.4.6.1) to

$$\begin{aligned}
&\gamma_{44}w_{u,t,1,4} + \gamma_{52}w_{u,t,1,6} + \gamma_{70}w_{u,t,1,9} + a_4w_{u,t,1,11} + \gamma_{\{91,108\}}w_{u,t,1,12}\\
&\quad + \gamma_{\{93,126\}}w_{u,t,1,15} + \gamma_{\{69,73,105,112,113\}}w_{u,t,1,17} + \gamma_{\{105,113\}}w_{u,t,1,18}\\
&\quad + a_5w_{u,t,1,13} + a_6w_{u,t,1,19} + \gamma_{112}w_{u,t,1,20} + a_7w_{u,t,1,21} = 0,\\
&\gamma_{\{41,48,114,117\}}w_{u,t,1,4} + \gamma_{\{42,49,82,85\}}w_{u,t,1,6} + \gamma_{\{69,73,96\}}w_{u,t,1,9}\\
&\quad + \gamma_{\{44,48,93,125,126\}}w_{u,t,1,11} + \gamma_{\{42,82,91,107,123\}}w_{u,t,1,12}\\
&\quad + \gamma_{\{42,52,91,107,108\}}w_{u,t,1,13} + \gamma_{\{48,93,114,122,125\}}w_{u,t,1,15}\\
&\quad + \gamma_{\{70,73,105,112,113\}}w_{u,t,1,17} + \gamma_{\{73,96,105,112\}}w_{u,t,1,18}\\
&\quad + a_8w_{u,t,1,19} + \gamma_{113}w_{u,t,1,20} + a_9w_{u,t,1,21} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_4 = \gamma_{\{41,48,93,122,125,126\}}, \quad a_5 = \gamma_{\{42,49,91,107,108,123\}},\\
&a_6 = \gamma_{\{42,49,69,73,85,91,93,108,117,123,125,126\}}, \quad a_7 = \gamma_{\{85,93,105,107,113,126\}},\\
&a_8 = \gamma_{\{42,48,52,70,91,93,96,107,114,122,126\}}, \quad a_9 = \gamma_{\{48,73,93,96,105,108,112,114,122,125\}}.
\end{aligned}$$

These equalities imply

$$\text{(8.4.6.5)} \qquad \begin{cases} a_i = 0,\ i = 4,5,\dots,9,\\ \gamma_j = 0,\ j = 44,52,69,70,73,96,105,112,113,\\ \gamma_{\{91,108\}} = \gamma_{\{93,126\}} = \gamma_{\{41,48,114,117\}} = 0,\\ \gamma_{\{42,49,82,85\}} = \gamma_{\{48,93,125,126\}} = \gamma_{\{42,82,91,107,123\}} = 0,\\ \gamma_{\{42,91,107,108\}} = \gamma_{\{48,93,114,122,125\}} = 0. \end{cases}$$

Combining (8.4.6.2), (8.4.6.3), (8.4.6.4) and (8.4.6.5), we get $\gamma_j = 0$ for all j. So, the proposition is proved. □

Remark 8.4.7. By a direct computation, we see that the $\mathbb{F}_2$-subspace of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{2^{u+3}+7}$ generated by $[\theta_i], i = 1, 2, \ldots, 15$, which are defined as in the proof of Proposition 8.4.5, is an $GL_4(\mathbb{F}_2)$-submodule of $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_4)_{2^{u+3}+7}$.

8.5. **The case $s \geqslant 2, t = 1$ and $u = 1$.**

According to Kameko [8], for $s \geqslant 2$, $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+2}+2^{s+1}+2^s-3} = 8$ with a basis given by the classes $w_{1,1,s,j}, 1 \leqslant j \leqslant 8$, which are determined as follows:

1. $[2^s - 1, 2^{s+1} - 1, 2^{s+2} - 1]$,
2. $[2^s - 1, 2^{s+2} - 1, 2^{s+1} - 1]$,
3. $[2^{s+1} - 1, 2^s - 1, 2^{s+2} - 1]$,
4. $[2^{s+1} - 1, 2^{s+2} - 1, 2^s - 1]$,
5. $[2^{s+2} - 1, 2^s - 1, 2^{s+1} - 1]$,
6. $[2^{s+2} - 1, 2^{s+1} - 1, 2^s - 1]$,
7. $[2^{s+1} - 1, 2^{s+1} - 1, 2^{s+1} + 2^s - 1]$,
8. $[2^{s+1} - 1, 2^{s+1} + 2^s - 1, 2^{s+1} - 1]$.

So, we easily obtain

Proposition 8.5.1. *For any positive integer $s \geqslant 2$, we have*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+2}+2^{s+1}+2^s-3} = 32.$$

By Proposition 2.8, we need only to determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}+2^s-3}$. The main result of this subsection is the following.

Theorem 8.5.2. *For $s \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}+2^s-3}$ is an $\mathbb{F}_2$-vector space of dimension 88 with a basis consisting of all the classes represented by the monomials $a_{1,1,s,j}, 1 \leqslant j \leqslant 88$, which are determined as follows:*

For $s \geqslant 2$,

1. $(1, 2^s-2, 2.2^s-1, 4.2^s-1)$, 2. $(1, 2^s-2, 4.2^s-1, 2.2^s-1)$,
3. $(1, 2.2^s-1, 2^s-2, 4.2^s-1)$, 4. $(1, 2.2^s-1, 4.2^s-1, 2^s-2)$,
5. $(1, 4.2^s-1, 2^s-2, 2.2^s-1)$, 6. $(1, 4.2^s-1, 2.2^s-1, 2^s-2)$,
7. $(2.2^s-1, 1, 2^s-2, 4.2^s-1)$, 8. $(2.2^s-1, 1, 4.2^s-1, 2^s-2)$,
9. $(2.2^s-1, 4.2^s-1, 1, 2^s-2)$, 10. $(4.2^s-1, 1, 2^s-2, 2.2^s-1)$,
11. $(4.2^s-1, 1, 2.2^s-1, 2^s-2)$, 12. $(4.2^s-1, 2.2^s-1, 1, 2^s-2)$,
13. $(1, 2^s-1, 2.2^s-2, 4.2^s-1)$, 14. $(1, 2^s-1, 4.2^s-1, 2.2^s-2)$,
15. $(1, 2.2^s-2, 2^s-1, 4.2^s-1)$, 16. $(1, 2.2^s-2, 4.2^s-1, 2^s-1)$,
17. $(1, 4.2^s-1, 2^s-1, 2.2^s-2)$, 18. $(1, 4.2^s-1, 2.2^s-2, 2^s-1)$,
19. $(2^s-1, 1, 2.2^s-2, 4.2^s-1)$, 20. $(2^s-1, 1, 4.2^s-1, 2.2^s-2)$,
21. $(2^s-1, 4.2^s-1, 1, 2.2^s-2)$, 22. $(4.2^s-1, 1, 2^s-1, 2.2^s-2)$,
23. $(4.2^s-1, 1, 2.2^s-2, 2^s-1)$, 24. $(4.2^s-1, 2^s-1, 1, 2.2^s-2)$,
25. $(1, 2^s-1, 2.2^s-1, 4.2^s-2)$, 26. $(1, 2^s-1, 4.2^s-2, 2.2^s-1)$,
27. $(1, 2.2^s-1, 2^s-1, 4.2^s-2)$, 28. $(1, 2.2^s-1, 4.2^s-2, 2^s-1)$,
29. $(1, 4.2^s-2, 2^s-1, 2.2^s-1)$, 30. $(1, 4.2^s-2, 2.2^s-1, 2^s-1)$,
31. $(2^s-1, 1, 2.2^s-1, 4.2^s-2)$, 32. $(2^s-1, 1, 4.2^s-2, 2.2^s-1)$,
33. $(2^s-1, 2.2^s-1, 1, 4.2^s-2)$, 34. $(2.2^s-1, 1, 2^s-1, 4.2^s-2)$,
35. $(2.2^s-1, 1, 4.2^s-2, 2^s-1)$, 36. $(2.2^s-1, 2^s-1, 1, 4.2^s-2)$,
37. $(1, 2.2^s-2, 2.2^s-1, 3.2^s-1)$, 38. $(1, 2.2^s-2, 3.2^s-1, 2.2^s-1)$,
39. $(1, 2.2^s-1, 2.2^s-2, 3.2^s-1)$, 40. $(1, 2.2^s-1, 3.2^s-1, 2.2^s-2)$,
41. $(2.2^s-1, 1, 2.2^s-2, 3.2^s-1)$, 42. $(2.2^s-1, 1, 3.2^s-1, 2.2^s-2)$,
43. $(2.2^s-1, 3.2^s-1, 1, 2.2^s-2)$, 44. $(1, 2.2^s-1, 2.2^s-1, 3.2^s-2)$,
45. $(1, 2.2^s-1, 3.2^s-2, 2.2^s-1)$, 46. $(2.2^s-1, 1, 2.2^s-1, 3.2^s-2)$,
47. $(2.2^s-1, 1, 3.2^s-2, 2.2^s-1)$, 48. $(2.2^s-1, 2.2^s-1, 1, 3.2^s-2)$,
49. $(3, 2.2^s-3, 2^s-2, 4.2^s-1)$, 50. $(3, 2.2^s-3, 4.2^s-1, 2^s-2)$,
51. $(3, 4.2^s-1, 2.2^s-3, 2^s-2)$, 52. $(4.2^s-1, 3, 2.2^s-3, 2^s-2)$,
53. $(3, 2.2^s-1, 4.2^s-3, 2^s-2)$, 54. $(3, 4.2^s-3, 2^s-2, 2.2^s-1)$,
55. $(3, 4.2^s-3, 2.2^s-1, 2^s-2)$, 56. $(2.2^s-1, 3, 4.2^s-3, 2^s-2)$,
57. $(3, 2^s-1, 2.2^s-3, 4.2^s-2)$, 58. $(3, 2.2^s-3, 2^s-1, 4.2^s-2)$,
59. $(3, 2.2^s-3, 4.2^s-2, 2^s-1)$, 60. $(3, 2^s-1, 4.2^s-3, 2.2^s-2)$,
61. $(3, 4.2^s-3, 2^s-1, 2.2^s-2)$, 62. $(3, 4.2^s-3, 2.2^s-2, 2^s-1)$,
63. $(3, 2.2^s-3, 2.2^s-2, 3.2^s-1)$, 64. $(3, 2.2^s-3, 3.2^s-1, 2.2^s-2)$,
65. $(3, 2.2^s-3, 2.2^s-1, 3.2^s-2)$, 66. $(3, 2.2^s-3, 3.2^s-2, 2.2^s-1)$,
67. $(3, 2.2^s-1, 2.2^s-3, 3.2^s-2)$, 68. $(2.2^s-1, 3, 2.2^s-3, 3.2^s-2)$,
69. $(3, 2.2^s-1, 3.2^s-3, 2.2^s-2)$, 70. $(2.2^s-1, 3, 3.2^s-3, 2.2^s-2)$,
71. $(7, 4.2^s-5, 2.2^s-3, 2^s-2)$.

For $s = 2$,

72. $(3, 3, 4, 15)$, 73. $(3, 3, 15, 4)$, 74. $(3, 15, 3, 4)$,
75. $(15, 3, 3, 4)$, 76. $(3, 3, 7, 12)$, 77. $(3, 3, 12, 7)$,
78. $(3, 7, 3, 12)$, 79. $(7, 3, 3, 12)$, 80. $(7, 7, 9, 2)$,
81. $(3, 7, 11, 4)$, 82. $(7, 3, 11, 4)$, 83. $(7, 11, 3, 4)$,
84. $(7, 9, 3, 6)$, 85. $(3, 7, 7, 8)$, 86. $(7, 3, 7, 8)$,
87. $(7, 7, 3, 8)$, 88. $(7, 7, 8, 3)$.

For $s \geqslant 3$,

72. $(3, 2^s-3, 2.2^s-2, 4.2^s-1)$, 73. $(3, 2^s-3, 4.2^s-1, 2.2^s-2)$,
74. $(3, 4.2^s-1, 2^s-3, 2.2^s-2)$, 75. $(4.2^s-1, 3, 2^s-3, 2.2^s-2)$,
76. $(3, 2^s-3, 2.2^s-1, 4.2^s-2)$, 77. $(3, 2^s-3, 4.2^s-2, 2.2^s-1)$,
78. $(3, 2.2^s-1, 2^s-3, 4.2^s-2)$, 79. $(2.2^s-1, 3, 2^s-3, 4.2^s-2)$,
80. $(2^s-1, 3, 2.2^s-3, 4.2^s-2)$, 81. $(2^s-1, 3, 4.2^s-3, 2.2^s-2)$,
82. $(7, 2.2^s-5, 2^s-3, 4.2^s-2)$, 83. $(7, 4.2^s-5, 2^s-3, 2.2^s-2)$,
84. $(7, 2.2^s-5, 4.2^s-3, 2^s-2)$, 85. $(7, 2.2^s-5, 2.2^s-3, 3.2^s-2)$,
86. $(7, 2.2^s-5, 3.2^s-3, 2.2^s-2)$.

For $s = 3$,

87. $(7, 7, 9, 30)$, 88. $(7, 7, 25, 14)$.

For $s \geqslant 4$,

87. $(7, 2^s-5, 2.2^s-3, 4.2^s-2)$, 88. $(7, 2^s-5, 4.2^s-3, 2.2^s-2)$.

We prove the theorem by proving the following.

Proposition 8.5.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}+2^s-3}$ is generated by the elements listed in Theorem 8.5.2.*

The proof of this proposition is based on the following lemmas.

Lemma 8.5.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 1\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(7, 9, 2, 7), (7, 9, 7, 2), (7, 9, 6, 3), (3, 7, 8, 7),$$
$$(7, 3, 8, 7), (7, 8, 3, 7), (7, 8, 7, 3).$$

We prove the lemma for the matrices associated with the monomials

$$(7, 9, 2, 7), (7, 9, 6, 3), (7, 3, 8, 7), (7, 8, 3, 7).$$

By a direct computation, we have

$$\begin{aligned}(7, 9, 2, 7) &= Sq^1(7, 7, 3, 7) + Sq^2(7, 7, 2, 7) + Sq^4\big((5, 7, 2, 7) + (4, 7, 3, 7)\big)\\ &\quad + (5, 11, 2, 7) + (5, 7, 2, 11) + (7, 7, 2, 9) + (4, 11, 3, 7)\\ &\quad + (4, 7, 3, 11) + (7, 8, 3, 7) + (7, 7, 3, 8) \mod \mathcal{L}_4(3; 3; 2; 1),\end{aligned}$$

$$\begin{aligned}(7,9,6,3) &= Sq^1(7,7,5,5) + Sq^2\big((7,7,6,3) + (7,7,3,6)\big)\\ &\quad + Sq^4\big((5,7,6,3) + (5,7,3,6)\big) + (7,7,8,3)\\ &\quad + (5,11,6,3) + (5,7,10,3) + (5,11,3,6) + (5,7,3,10)\\ &\quad + (7,9,3,6) + (7,7,3,8) \bmod \mathcal{L}_4(3;3;2;1),\\ (7,3,8,7) &= Sq^1(7,3,7,7) + Sq^2(7,2,7,7) + Sq^4\big((4,3,7,7)\\ &\quad + (5,2,7,7)\big) + (5,2,11,7) + (5,2,7,11) + (7,2,9,7) + (7,2,7,9)\\ &\quad + (7,3,7,8) + (4,3,11,7) + (4,3,7,11) \quad \bmod \mathcal{L}_4(3;3;2;1),\\ (7,8,3,7) &= Sq^1(7,5,5,7) + Sq^2\big((7,6,3,7) + (7,3,6,7)\big)\\ &\quad + Sq^4\big((5,6,3,7) + (5,3,6,7)\big) + (5,10,3,7)\\ &\quad + (5,6,3,11) + (7,6,3,9) + (7,3,8,7) + (7,3,6,9)\\ &\quad + (5,3,10,7) + (5,3,6,11) \quad \bmod \mathcal{L}_4(3;3;2;1).\end{aligned}$$

The lemma is proved. □

Lemma 8.5.5. *The following matrices are strictly inadmissible*

$$\begin{pmatrix}1&1&1&0\\1&1&0&1\\1&1&0&1\\1&1&0&0\\0&0&1&0\end{pmatrix} \quad \begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&1&0\\0&1&1&0\\0&0&0&1\end{pmatrix} \quad \begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&1&0\\1&0&1&0\\0&0&0&1\end{pmatrix}$$

$$\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&1&0\\1&1&0&0\\0&0&0&1\end{pmatrix} \quad \begin{pmatrix}1&1&0&1\\1&1&0&1\\1&1&0&1\\1&1&0&0\\0&0&1&0\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(15,15,17,6), (7,15,15,16), (15,7,15,16), (15,15,7,16), (15,15,16,7).$$

By a direct computation, we have

$$\begin{aligned}(15,15,17,6) &= Sq^1\big((15,15,15,7) + (15,11,15,11)\big) + Sq^2\big((15,15,15,6)\\ &\quad + (15,11,15,10)\big) + Sq^4\big((15,13,15,6) + (15,12,15,7)\big)\\ &\quad + Sq^8\big((9,15,15,6) + (11,13,15,6) + (8,15,15,7) + (11,12,15,7)\big)\\ &\quad + (9,23,15,6) + (9,15,23,6) + (15,13,19,6) + (11,21,15,6)\\ &\quad + (11,13,23,6) + (8,23,15,7) + (8,15,23,7) + (15,12,19,7)\\ &\quad + (15,15,16,7) + (11,20,15,7) + (11,12,23,7) \quad \bmod \mathcal{L}_4(3;3;3;2;1),\end{aligned}$$

$$\begin{aligned}(15,7,15,16) &= Sq^1(15,7,15,15) + Sq^2(15,7,14,15) + Sq^4\big((15,4,15,15)\\ &\quad + (15,5,14,15)\big) + Sq^8\big((8,7,15,15) + (11,4,15,15) + (9,7,14,15)\\ &\quad + (11,5,14,15)\big) + (8,7,23,15) + (8,7,15,23) + (15,4,19,15)\\ &\quad + (15,4,15,19) + (11,4,23,15) + (11,4,15,23) + (9,7,22,15)\\ &\quad + (9,7,14,23) + (15,7,14,17) + (11,5,22,15) + (11,5,14,23)\\ &\quad + (15,5,18,15) + (15,5,14,19) \mod \mathcal{L}_4(3;3;3;2;1),\\ (7,15,15,16) &= \overline{\varphi}_1(15,7,15,16),\\ (15,15,7,16) &= Sq^1\big((15,15,7,15) + (15,11,11,15)\big) + Sq^2\big((15,15,6,15)\\ &\quad + (15,11,10,15)\big) + Sq^4\big((15,12,7,15) + (15,13,6,15)\big)\\ &\quad + Sq^8\big((8,15,7,15) + (11,12,7,15) + (9,15,6,15)\\ &\quad + (11,13,6,15)\big) + (8,23,7,15) + (8,15,7,23)\\ &\quad + (15,12,7,19) + (11,20,7,15) + (11,12,7,23)\\ &\quad + (15,15,6,17) + (9,23,6,15) + (9,15,6,23) + (15,13,6,19)\\ &\quad + (11,21,6,15) + (11,13,6,23) \mod \mathcal{L}_4(3;3;3;2;1),\\ (15,15,16,7) &= Sq^1\big((15,15,13,9) + (15,15,11,11) + (15,15,9,13)\\ &\quad + (15,11,13,13)\big) + Sq^2\big((15,15,14,7) + (15,15,11,10)\\ &\quad + (15,15,10,11) + (15,15,7,14) + (15,11,14,11)\\ &\quad + (15,11,11,14)\big) + Sq^4\big((15,13,14,7) + (15,13,7,14)\big)\\ &\quad + Sq^8\big((9,15,14,7) + (11,13,14,7) + (9,15,7,14)\\ &\quad + (11,13,7,14)\big) + (9,23,14,7) + (9,15,22,7)\\ &\quad + (15,13,18,7) + (11,21,14,7) + (11,13,22,7) + (15,15,7,16)\\ &\quad + (9,23,7,14) + (9,15,7,22) + (15,13,7,18)\\ &\quad + (11,21,7,14) + (11,13,7,22) \mod \mathcal{L}_4(3;3;3;2;1).\end{aligned}$$

The lemma is proved. □

Combining Lemmas 3.5, 3.6, 5.10, 6.3.4, 6.3.5, 8.5.4, 8.5.5, Theorem 2.5 and the results in Section 7, we get Proposition 8.5.3.

Now, we prove that the classes listed in Theorem 8.5.2 are linearly independent.

Proposition 8.5.6. *The elements* $[a_{1,1,2,j}], 1 \leqslant j \leqslant 88$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{25}$.

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{88} \gamma_j [a_{1,1,2,j}] = 0, \tag{8.5.6.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.5.6.1), we obtain

$$\begin{aligned}
&\gamma_1[3,7,15] + \gamma_2[3,15,7] + \gamma_{15}[7,3,15] + \gamma_{16}[7,15,3]\\
&\quad + \gamma_{29}[15,3,7] + \gamma_{30}[15,7,3] + \gamma_{37}[7,7,11] + \gamma_{38}[7,11,7] = 0,\\
&\gamma_3[3,7,15] + \gamma_5[3,15,7] + \gamma_{\{13,72\}}[7,3,15] + \gamma_{18}[7,15,3] + \gamma_{45}[7,11,7]\\
&\quad + \gamma_{\{26,77\}}[15,3,7] + \gamma_{\{28,88\}}[15,7,3] + \gamma_{\{39,45\}}[7,7,11] = 0,\\
&\gamma_4[3,7,15] + \gamma_6[3,15,7] + \gamma_{\{14,73\}}[7,3,15] + \gamma_{\{17,74\}}[7,15,3]\\
&\quad + \gamma_{\{25,76,86\}}[15,3,7] + \gamma_{\{27,78,87\}}[15,7,3]\\
&\quad + \gamma_{\{40,44,81,85\}}[7,7,11] + \gamma_{\{44,85\}}[7,11,7] = 0,\\
&\gamma_{\{19,49,72\}}[3,7,15] + \gamma_7[7,3,15] + \gamma_{\{35,88\}}[7,15,3] + \gamma_{10}[15,3,7]\\
&\quad + \gamma_{\{32,54,66,77\}}[3,15,7] + \gamma_{23}[15,7,3] + \gamma_{41}[7,7,11] + \gamma_{47}[7,11,7] = 0,\\
&\gamma_{\{20,50,73\}}[3,7,15] + \gamma_{\{31,55,65,76,85\}}[3,15,7]\\
&\quad + \gamma_8[7,3,15] + \gamma_{\{34,79,83,84,87\}}[7,15,3] + \gamma_{11}[15,3,7]\\
&\quad + \gamma_{\{22,75\}}[15,7,3] + \gamma_{\{42,82\}}[7,7,11] + \gamma_{\{46,86\}}[7,11,7] = 0,\\
&\gamma_{\{33,53,67,69,78,81,85\}}[3,7,15] + \gamma_{\{36,56,68,70,79,82,86\}}[7,3,15]\\
&\quad + \gamma_9[7,15,3] + \gamma_{\{21,51,74\}}[3,15,7] + \gamma_{\{24,52,75\}}[15,3,7]\\
&\quad + \gamma_{12}[15,7,3] + \gamma_{\{48,80,87,88\}}[7,7,11] + \gamma_{\{43,71,83\}}[7,11,7] = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\text{(8.5.6.2)}\qquad \begin{cases}
\gamma_j = 0,\ j = 1,2,\ldots,12,\\
\quad 15,16,18,23,29,30,37,38,39,41,45,47,\\
\gamma_{\{13,72\}} = \gamma_{\{26,77\}} = \gamma_{\{28,88\}} = \gamma_{\{24,52,75\}} = \gamma_{\{17,74\}} = 0,\\
\gamma_{\{25,76,86\}} = \gamma_{\{27,78,87\}} = \gamma_{\{40,44,81,85\}} = \gamma_{\{44,85\}} = 0,\\
\gamma_{\{19,49,72\}} = \gamma_{\{32,54,66,77\}} = \gamma_{\{35,88\}} = \gamma_{\{20,50,73\}} = 0,\\
\gamma_{\{31,55,65,76,85\}} = \gamma_{\{22,75\}} = \gamma_{\{43,71,83\}} = \gamma_{\{42,82\}} = 0,\\
\gamma_{\{34,79,83,84,87\}} = \gamma_{\{46,86\}} = \gamma_{\{48,80,87,88\}} = \gamma_{\{81,85\}} = 0,\\
\gamma_{\{33,53,67,69,78,21,51,74\}} = \gamma_{\{36,56,68,70,79,82,86\}} = \gamma_{\{14,73\}} = 0.
\end{cases}$$

With the aid of (8.5.6.2), the homomorphisms g_1, g_2 send (8.5.6.1) to

$$\begin{aligned}
&\gamma_{19}[3,7,15] + \gamma_{32}[3,15,7] + \gamma_{49}[7,3,15] + \gamma_{\{28,59\}}[7,15,3]\\
&\quad + \gamma_{54}[15,3,7] + \gamma_{62}[15,7,3] + \gamma_{63}[7,7,11] + \gamma_{66}[7,11,7] = 0,\\
&\gamma_{20}[3,7,15] + \gamma_{\{25,31,76\}}[3,15,7] + \gamma_{50}[7,3,15] + \gamma_{\{27,34,58,78,79,87\}}[7,15,3]\\
&\quad + \gamma_{55}[15,3,7] + \gamma_{\{61,83,84\}}[15,7,3] + \gamma_{64}[7,7,11] + \gamma_{65}[7,11,7] = 0.
\end{aligned}$$

These equalities imply

$$\text{(8.5.6.3)}\qquad \begin{cases}
\gamma_j = 0,\ j = 19,20,32,49,50,54,55,62,63,64,65,66,\\
\gamma_{\{28,59\}} = \gamma_{\{25,31,76\}} = \gamma_{\{27,34,58,78,79,87\}} = \gamma_{\{61,83,84\}} = 0.
\end{cases}$$

With the aid of (8.5.6.2) and (8.5.6.3), the homomorphisms g_3, g_4 send (8.5.6.1) to

$$\begin{aligned}
&\gamma_{21}[3,7,15] + \gamma_{\{27,33,48,78,87\}}[3,15,7] + \gamma_{\{53,80\}}[15,3,7]\\
&\quad + \gamma_{51}[7,3,15] + \gamma_{\{28,43,69,71,80,83\}}[7,7,11] + a_1[7,15,3]\\
&\quad + \gamma_{\{28,48,67,69,80,87\}}[7,11,7] + a_2[15,7,3] = 0,\\
&\gamma_{24}[3,7,15] + a_3[3,15,7] + \gamma_{52}[7,3,15] + a_4[7,15,3] + \gamma_{\{56,71,80\}}[15,3,7]\\
&\quad + a_5[15,7,3] + \gamma_{\{28,43,70,80,83,84\}}[7,7,11] + a_6[7,11,7] = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_1 = \gamma_{\{22,24,25,36,46,52,57,68,76,79\}}, \quad a_2 = \gamma_{\{22,24,42,52,56,60,70\}},\\
&a_3 = \gamma_{\{34,36,43,48,79,83,84,87\}}, \quad a_4 = \gamma_{\{13,17,21,31,33,44,51,57,58,61,67,76,78\}},\\
&a_5 = \gamma_{\{14,17,21,26,40,51,53,59,60,61,69\}}, \quad a_6 = \gamma_{\{28,43,48,68,70,71,80,83,84,87\}}.
\end{aligned}$$

From the above equalities, we obtain

$$\begin{cases}
a_i = 0,\ i = 1,2\ldots,6,\ \gamma_j = 0,\ j = 21,24,51,52,\\
\gamma_{\{27,33,48,78,87\}} = \gamma_{\{53,80\}} = \gamma_{\{28,43,69,71,80,83\}} = 0,\\
\gamma_{\{28,48,67,69,80,87\}} = \gamma_{\{56,71,80\}} = \gamma_{\{28,43,70,80,83,84\}} = 0.
\end{cases} \tag{8.5.6.4}$$

With the aid of (8.5.6.2), (8.5.6.3) and (8.5.6.4), the homomorphism h sends (8.5.6.1) to

$$\begin{aligned}
&a_7[3,7,15] + \gamma_{\{28,56,60,70\}}[3,15,7] + a_8[7,3,15] + \gamma_{56}[7,15,3]\\
&\quad + \gamma_{\{43,61,83,84\}}[15,3,7] + \gamma_{71}[15,7,3] + a_9[7,7,11] + a_{10}[7,11,7] = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_7 = \gamma_{\{25,31,34,36,57,68,76,79\}}, \quad a_8 = \gamma_{\{27,33,34,36,48,58,78,79,87\}},\\
&a_9 = \gamma_{\{43,48,67,68,71,83,84,87\}}, \quad a_{10} = \gamma_{\{28,43,53,69,70,71,83,84\}}.
\end{aligned}$$

From the above equalities, it implies

$$\begin{cases}
a_i = 0,\ i = 7,8,9,10,\ \gamma_j = 0,\ j = 56,71,\\
\gamma_{\{28,60,70\}} = \gamma_{\{43,61,83,84\}} = 0.
\end{cases} \tag{8.5.6.5}$$

Combining (8.5.6.2), (8.5.6.3), (8.5.6.4) and (8.5.6.5), we get $\gamma_j = 0$ for $1 \leqslant j \leqslant 88$. The proposition is proved. □

Proposition 8.5.7. *For $s \geqslant 3$, the elements $[a_{1,1,s,j}], 1 \leqslant j \leqslant 88$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+2}+2^{s+1}+2^s-3}$.*

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{88} \gamma_j [a_{1,1,s,j}] = 0, \tag{8.5.7.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.5.7.1), we have

$$\begin{aligned}
&\gamma_1 w_{1,1,s,1} + \gamma_2 w_{1,1,s,2} + \gamma_{15} w_{1,1,s,3} + \gamma_{16} w_{1,1,s,4}\\
&\quad + \gamma_{29} w_{1,1,s,5} + \gamma_{30} w_{1,1,s,6} + \gamma_{37} w_{1,1,s,7} + \gamma_{38} w_{1,1,s,8} = 0,\\
&\gamma_3 w_{1,1,s,1} + \gamma_5 w_{1,1,s,2} + \gamma_{13} w_{1,1,s,3} + \gamma_{18} w_{1,1,s,4}\\
&\quad + \gamma_{26} w_{1,1,s,5} + \gamma_{28} w_{1,1,s,6} + \gamma_{\{39,45\}} w_{1,1,s,7} + \gamma_{45} w_{1,1,s,8} = 0,\\
&\gamma_4 w_{1,1,s,1} + \gamma_6 w_{1,1,s,2} + \gamma_{14} w_{1,1,s,3} + \gamma_{17} w_{1,1,s,4}\\
&\quad + \gamma_{25} w_{1,1,s,5} + \gamma_{27} w_{1,1,s,6} + \gamma_{\{40,44\}} w_{1,1,s,7} + \gamma_{44} w_{1,1,s,8} = 0,\\
&\gamma_{19} w_{1,1,s,1} + \gamma_{32} w_{1,1,s,2} + \gamma_7 w_{1,1,s,3} + \gamma_{35} w_{1,1,s,4}\\
&\quad + \gamma_{10} w_{1,1,s,5} + \gamma_{23} w_{1,1,s,6} + \gamma_{41} w_{1,1,s,7} + \gamma_{47} w_{1,1,s,8} = 0,\\
&\gamma_{20} w_{1,1,s,1} + \gamma_{31} w_{1,1,s,2} + \gamma_8 w_{1,1,s,3} + \gamma_{34} w_{1,1,s,4}\\
&\quad + \gamma_{11} w_{1,1,s,5} + \gamma_{22} w_{1,1,s,6} + \gamma_{42} w_{1,1,s,7} + \gamma_{46} w_{1,1,s,8} = 0,\\
&\gamma_{33} w_{1,1,s,1} + \gamma_{21} w_{1,1,s,2} + \gamma_{36} w_{1,1,s,3} + \gamma_9 w_{1,1,s,4}\\
&\quad + \gamma_{24} w_{1,1,s,5} + \gamma_{12} w_{1,1,s,6} + \gamma_{48} w_{1,1,s,7} + \gamma_{43} w_{1,1,s,8} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\gamma_j = 0,\ j = 1, 2, \ldots, 48. \tag{8.5.7.2}$$

With the aid of (8.5.7.2), the homomorphisms g_1, g_2, g_3, g_4 send (8.5.7.1) to

$$\begin{aligned}
&\gamma_{72} w_{1,1,s,1} + \gamma_{77} w_{1,1,s,2} + \gamma_{49} w_{1,1,s,3} + \gamma_{59} w_{1,1,s,4}\\
&\quad + \gamma_{54} w_{1,1,s,5} + \gamma_{62} w_{1,1,s,6} + \gamma_{63} w_{1,1,s,7} + \gamma_{66} w_{1,1,s,8} = 0,\\
&\gamma_{73} w_{1,1,s,1} + \gamma_{76} w_{1,1,s,2} + \gamma_{50} w_{1,1,s,3} + \gamma_{58} w_{1,1,s,4}\\
&\quad + \gamma_{55} w_{1,1,s,5} + \gamma_{61} w_{1,1,s,6} + \gamma_{64} w_{1,1,s,7} + \gamma_{65} w_{1,1,s,8} = 0,\\
&\gamma_{74} w_{1,1,s,1} + \gamma_{78} w_{1,1,s,2} + \gamma_{51} w_{1,1,s,3} + a_1 w_{1,1,s,4}\\
&\quad + \gamma_{53} w_{1,1,s,5} + a_2 w_{1,1,s,6} + \gamma_{69} w_{1,1,s,7} + \gamma_{\{67,69\}} w_{1,1,s,8} = 0,\\
&\gamma_{75} w_{1,1,s,1} + \gamma_{79} w_{1,1,s,2} + \gamma_{52} w_{1,1,s,3} + a_3 w_{1,1,s,4}\\
&\quad + \gamma_{56} w_{1,1,s,5} + a_4 w_{1,1,s,6} + \gamma_{70} w_{1,1,s,7} + \gamma_{\{68,70\}} w_{1,1,s,8} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases} \gamma_{\{57,87\}}, & s = 3,\\ \gamma_{57} & s \geqslant 4, \end{cases} \quad a_2 = \begin{cases} \gamma_{\{60,88\}}, & s = 3,\\ \gamma_{60}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{71,80,82,83,85,87\}}, & s = 3,\\ \gamma_{80}, & s \geqslant 4, \end{cases} \quad a_4 = \begin{cases} \gamma_{\{71,81,83,84,86,88\}}, & s = 3,\\ \gamma_{81}, & s \geqslant 4. \end{cases}$$

These equalities imply

$$\begin{cases} a_1 = a_2 = a_3 = a_4 = 0,\\ \gamma_j = 0,\ j = 49, 50, \ldots, 56, 58, 59,\\ 61, 62, \ldots, 66, 69, 70, 72, 73, \ldots, 79. \end{cases} \tag{8.5.7.3}$$

With the aid of (8.5.7.2) and (8.5.7.3), the homomorphism h sends (8.5.7.1) to

$$\begin{aligned}&a_5w_{1,1,s,1} + a_6w_{1,1,s,2} + \gamma_{82}w_{1,1,s,3} + \gamma_{84}w_{1,1,s,4}\\ &\quad + \gamma_{83}w_{1,1,s,5} + \gamma_{71}w_{1,1,s,6} + \gamma_{85}w_{1,1,s,7} + \gamma_{86}w_{1,1,s,8} = 0,\end{aligned}$$

where

$$a_5 = \begin{cases}\gamma_{80}, & s = 3,\\ \gamma_{87}, & s \geqslant 4,\end{cases} \quad a_6 = \begin{cases}\gamma_{81}, & s = 3,\\ \gamma_{88}, & s \geqslant 4.\end{cases}$$

From the above equalities, it implies

$$a_5 = a_6 = 0,\ \gamma_j = 0,\ j = 71, 82, 83, 84, 85, 86. \tag{8.5.7.4}$$

Combining (8.5.7.2), (8.5.7.3) and (8.5.7.4), we get $\gamma_j = 0$ for $1 \leqslant j \leqslant 88$. The proposition is proved. □

8.6. **The case $s \geqslant 2, t = 1$ and $u = 2$.**

According to Kameko [8], for $s \geqslant 2$, $\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+3}+2^{s+1}+2^s-3} = 15$ with a basis given by the classes $w_{2,1,s,j}, 1 \leqslant j \leqslant 15$, which are determined as follows:

1. $[2^s - 1, 2.2^s - 1, 8.2^s - 1]$,
2. $[2^s - 1, 8.2^s - 1, 2.2^s - 1]$,
3. $[2.2^s - 1, 2^s - 1, 8.2^s - 1]$,
4. $[2.2^s - 1, 8.2^s - 1, 2^s - 1]$,
5. $[8.2^s - 1, 2^s - 1, 2.2^s - 1]$,
6. $[8.2^s - 1, 2.2^s - 1, 2^s - 1]$,
7. $[2^s - 1, 4.2^s - 1, 6.2^s - 1]$,
8. $[4.2^s - 1, 2^s - 1, 6.2^s - 1]$,
9. $[4.2^s - 1, 6.2^s - 1, 2^s - 1]$,
10. $[2.2^s - 1, 2.2^s - 1, 7.2^s - 1]$,
11. $[2.2^s - 1, 7.2^s - 1, 2.2^s - 1]$,
12. $[2.2^s - 1, 3.2^s - 1, 6.2^s - 1]$,
13. $[2.2^s - 1, 4.2^s - 1, 5.2^s - 1]$,
14. $[4.2^s - 1, 2.2^s - 1, 5.2^s - 1]$,
15. $[4.2^s - 1, 5.2^s - 1, 2.2^s - 1]$.

So, we easily obtain

Proposition 8.6.1. *For any positive integer $s \geqslant 2$, we have*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+3}+2^{s+1}+2^s-3} = 60.$$

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+1}+2^s-3}$. The main result in this subsection is

Theorem 8.6.2. *For $s \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+1}+2^s-3}$ is an $\mathbb{F}_2$-vector space with of dimension 165 a basis consisting of all the classes represented by the monomials $a_{2,1,s,j}, 1 \leqslant j \leqslant 165$, which are determined as follows:*

For $s \geqslant 2$,

1. $(1, 2^s - 2, 2.2^s - 1, 8.2^s - 1)$,
2. $(1, 2^s - 2, 8.2^s - 1, 2.2^s - 1)$,
3. $(1, 2.2^s - 1, 2^s - 2, 8.2^s - 1)$,
4. $(1, 2.2^s - 1, 8.2^s - 1, 2^s - 2)$,
5. $(1, 8.2^s - 1, 2^s - 2, 2.2^s - 1)$,
6. $(1, 8.2^s - 1, 2.2^s - 1, 2^s - 2)$,
7. $(2.2^s - 1, 1, 2^s - 2, 8.2^s - 1)$,
8. $(2.2^s - 1, 1, 8.2^s - 1, 2^s - 2)$,

9. $(2.2^s-1, 8.2^s-1, 1, 2^s-2)$,
10. $(8.2^s-1, 1, 2^s-2, 2.2^s-1)$,
11. $(8.2^s-1, 1, 2.2^s-1, 2^s-2)$,
12. $(8.2^s-1, 2.2^s-1, 1, 2^s-2)$,
13. $(1, 2^s-1, 2.2^s-2, 8.2^s-1)$,
14. $(1, 2^s-1, 8.2^s-1, 2.2^s-2)$,
15. $(1, 2.2^s-2, 2^s-1, 8.2^s-1)$,
16. $(1, 2.2^s-2, 8.2^s-1, 2^s-1)$,
17. $(1, 8.2^s-1, 2^s-1, 2.2^s-2)$,
18. $(1, 8.2^s-1, 2.2^s-2, 2^s-1)$,
19. $(2^s-1, 1, 2.2^s-2, 8.2^s-1)$,
20. $(2^s-1, 1, 8.2^s-1, 2.2^s-2)$,
21. $(2^s-1, 8.2^s-1, 1, 2.2^s-2)$,
22. $(8.2^s-1, 1, 2^s-1, 2.2^s-2)$,
23. $(8.2^s-1, 1, 2.2^s-2, 2^s-1)$,
24. $(8.2^s-1, 2^s-1, 1, 2.2^s-2)$,
25. $(1, 2^s-1, 2.2^s-1, 8.2^s-2)$,
26. $(1, 2^s-1, 8.2^s-2, 2.2^s-1)$,
27. $(1, 2.2^s-1, 2^s-1, 8.2^s-2)$,
28. $(1, 2.2^s-1, 8.2^s-2, 2^s-1)$,
29. $(1, 8.2^s-2, 2^s-1, 2.2^s-1)$,
30. $(1, 8.2^s-2, 2.2^s-1, 2^s-1)$,
31. $(2^s-1, 1, 2.2^s-1, 8.2^s-2)$,
32. $(2^s-1, 1, 8.2^s-2, 2.2^s-1)$,
33. $(2^s-1, 2.2^s-1, 1, 8.2^s-2)$,
34. $(2.2^s-1, 1, 2^s-1, 8.2^s-2)$,
35. $(2.2^s-1, 1, 8.2^s-2, 2^s-1)$,
36. $(2.2^s-1, 2^s-1, 1, 8.2^s-2)$,
37. $(1, 2^s-2, 4.2^s-1, 6.2^s-1)$,
38. $(1, 4.2^s-1, 2^s-2, 6.2^s-1)$,
39. $(1, 4.2^s-1, 6.2^s-1, 2^s-2)$,
40. $(4.2^s-1, 1, 2^s-2, 6.2^s-1)$,
41. $(4.2^s-1, 1, 6.2^s-1, 2^s-2)$,
42. $(4.2^s-1, 6.2^s-1, 1, 2^s-2)$,
43. $(1, 2^s-1, 4.2^s-2, 6.2^s-1)$,
44. $(1, 4.2^s-2, 2^s-1, 6.2^s-1)$,
45. $(1, 4.2^s-2, 6.2^s-1, 2^s-1)$,
46. $(2^s-1, 1, 4.2^s-2, 6.2^s-1)$,
47. $(1, 2^s-1, 4.2^s-1, 6.2^s-2)$,
48. $(1, 4.2^s-1, 2^s-1, 6.2^s-2)$,
49. $(1, 4.2^s-1, 6.2^s-2, 2^s-1)$,
50. $(2^s-1, 1, 4.2^s-1, 6.2^s-2)$,
51. $(2^s-1, 4.2^s-1, 1, 6.2^s-2)$,
52. $(4.2^s-1, 1, 2^s-1, 6.2^s-2)$,
53. $(4.2^s-1, 1, 6.2^s-2, 2^s-1)$,
54. $(4.2^s-1, 2^s-1, 1, 6.2^s-2)$,
55. $(1, 2.2^s-2, 2.2^s-1, 7.2^s-1)$,
56. $(1, 2.2^s-2, 7.2^s-1, 2.2^s-1)$,
57. $(1, 2.2^s-1, 2.2^s-2, 7.2^s-1)$,
58. $(1, 2.2^s-1, 7.2^s-1, 2.2^s-2)$,
59. $(2.2^s-1, 1, 2.2^s-2, 7.2^s-1)$,
60. $(2.2^s-1, 1, 7.2^s-1, 2.2^s-2)$,
61. $(2.2^s-1, 7.2^s-1, 1, 2.2^s-2)$,
62. $(1, 2.2^s-1, 2.2^s-1, 7.2^s-2)$,
63. $(1, 2.2^s-1, 7.2^s-2, 2.2^s-1)$,
64. $(2.2^s-1, 1, 2.2^s-1, 7.2^s-2)$,
65. $(2.2^s-1, 1, 7.2^s-2, 2.2^s-1)$,
66. $(2.2^s-1, 2.2^s-1, 1, 7.2^s-2)$,
67. $(3, 2.2^s-3, 2^s-2, 8.2^s-1)$,
68. $(3, 2.2^s-3, 8.2^s-1, 2^s-2)$,
69. $(3, 8.2^s-1, 2.2^s-3, 2^s-2)$,
70. $(8.2^s-1, 3, 2.2^s-3, 2^s-2)$,
71. $(1, 2.2^s-2, 3.2^s-1, 6.2^s-1)$,
72. $(1, 2.2^s-1, 3.2^s-2, 6.2^s-1)$,
73. $(2.2^s-1, 1, 3.2^s-2, 6.2^s-1)$,
74. $(1, 2.2^s-1, 3.2^s-1, 6.2^s-2)$,
75. $(2.2^s-1, 1, 3.2^s-1, 6.2^s-2)$,
76. $(2.2^s-1, 3.2^s-1, 1, 6.2^s-2)$,
77. $(3, 2.2^s-1, 8.2^s-3, 2^s-2)$,
78. $(3, 8.2^s-3, 2^s-2, 2.2^s-1)$,
79. $(3, 8.2^s-3, 2.2^s-1, 2^s-2)$,
80. $(2.2^s-1, 3, 8.2^s-3, 2^s-2)$,
81. $(1, 2.2^s-2, 4.2^s-1, 5.2^s-1)$,
82. $(1, 4.2^s-1, 2.2^s-2, 5.2^s-1)$,
83. $(1, 4.2^s-1, 5.2^s-1, 2.2^s-2)$,
84. $(3, 2^s-1, 2.2^s-3, 8.2^s-2)$,
85. $(3, 2.2^s-3, 2^s-1, 8.2^s-2)$,
86. $(3, 2.2^s-3, 8.2^s-2, 2^s-1)$,
87. $(4.2^s-1, 1, 2.2^s-2, 5.2^s-1)$,
88. $(4.2^s-1, 1, 5.2^s-1, 2.2^s-2)$,
89. $(4.2^s-1, 5.2^s-1, 1, 2.2^s-2)$,
90. $(1, 2.2^s-1, 4.2^s-2, 5.2^s-1)$,
91. $(1, 4.2^s-2, 2.2^s-1, 5.2^s-1)$,
92. $(1, 4.2^s-2, 5.2^s-1, 2.2^s-1)$,
93. $(2.2^s-1, 1, 4.2^s-2, 5.2^s-1)$,
94. $(1, 2.2^s-1, 4.2^s-1, 5.2^s-2)$,
95. $(1, 4.2^s-1, 2.2^s-1, 5.2^s-2)$,
96. $(1, 4.2^s-1, 5.2^s-2, 2.2^s-1)$,
97. $(2.2^s-1, 1, 4.2^s-1, 5.2^s-2)$,
98. $(2.2^s-1, 4.2^s-1, 1, 5.2^s-2)$,

99. $(4.2^s-1, 1, 2.2^s-1, 5.2^s-2)$,
100. $(4.2^s-1, 1, 5.2^s-2, 2.2^s-1)$,
101. $(4.2^s-1, 2.2^s-1, 1, 5.2^s-2)$,
102. $(3, 2^s-1, 8.2^s-3, 2.2^s-2)$,
103. $(3, 8.2^s-3, 2^s-1, 2.2^s-2)$,
104. $(3, 8.2^s-3, 2.2^s-2, 2^s-1)$,
105. $(3, 4.2^s-3, 2^s-2, 6.2^s-1)$,
106. $(3, 4.2^s-3, 6.2^s-1, 2^s-2)$,
107. $(3, 4.2^s-1, 6.2^s-3, 2^s-2)$,
108. $(4.2^s-1, 3, 6.2^s-3, 2^s-2)$,
109. $(3, 2^s-1, 4.2^s-3, 6.2^s-2)$,
110. $(3, 4.2^s-3, 2^s-1, 6.2^s-2)$,
111. $(3, 4.2^s-3, 6.2^s-2, 2^s-1)$,
112. $(3, 2.2^s-3, 2.2^s-2, 7.2^s-1)$,
113. $(3, 2.2^s-3, 7.2^s-1, 2.2^s-2)$,
114. $(3, 2.2^s-3, 2.2^s-1, 7.2^s-2)$,
115. $(3, 2.2^s-3, 7.2^s-2, 2.2^s-1)$,
116. $(3, 2.2^s-1, 2.2^s-3, 7.2^s-2)$,
117. $(2.2^s-1, 3, 2.2^s-3, 7.2^s-2)$,
118. $(3, 2.2^s-1, 7.2^s-3, 2.2^s-2)$,
119. $(2.2^s-1, 3, 7.2^s-3, 2.2^s-2)$,
120. $(7, 8.2^s-5, 2.2^s-3, 2^s-2)$,
121. $(3, 2.2^s-3, 3.2^s-2, 6.2^s-1)$,
122. $(3, 2.2^s-3, 3.2^s-1, 6.2^s-2)$,
123. $(3, 2.2^s-1, 3.2^s-3, 6.2^s-2)$,
124. $(2.2^s-1, 3, 3.2^s-3, 6.2^s-2)$,
125. $(3, 2.2^s-3, 4.2^s-2, 5.2^s-1)$,
126. $(3, 2.2^s-3, 4.2^s-1, 5.2^s-2)$,
127. $(3, 4.2^s-1, 2.2^s-3, 5.2^s-2)$,
128. $(4.2^s-1, 3, 2.2^s-3, 5.2^s-2)$,
129. $(3, 4.2^s-3, 2.2^s-2, 5.2^s-1)$,
130. $(3, 4.2^s-3, 5.2^s-1, 2.2^s-2)$,
131. $(3, 4.2^s-1, 5.2^s-3, 2.2^s-2)$,
132. $(4.2^s-1, 3, 5.2^s-3, 2.2^s-2)$,
133. $(3, 2.2^s-1, 4.2^s-3, 5.2^s-2)$,
134. $(3, 4.2^s-3, 2.2^s-1, 5.2^s-2)$,
135. $(3, 4.2^s-3, 5.2^s-2, 2.2^s-1)$,
136. $(2.2^s-1, 3, 4.2^s-3, 5.2^s-2)$,
137. $(7, 4.2^s-5, 6.2^s-3, 2^s-2)$,
138. $(7, 4.2^s-5, 2.2^s-3, 5.2^s-2)$,
139. $(7, 4.2^s-5, 5.2^s-3, 2.2^s-2)$.

For $s = 2$,

140. $(3, 3, 4, 31)$,
141. $(3, 3, 31, 4)$,
142. $(3, 31, 3, 4)$,
143. $(31, 3, 3, 4)$,
144. $(3, 3, 7, 28)$,
145. $(3, 3, 28, 7)$,
146. $(3, 7, 3, 28)$,
147. $(7, 3, 3, 28)$,
148. $(3, 7, 27, 4)$,
149. $(7, 3, 27, 4)$,
150. $(7, 27, 3, 4)$,
151. $(7, 7, 25, 2)$,
152. $(3, 3, 12, 23)$,
153. $(3, 3, 15, 20)$,
154. $(3, 15, 3, 20)$,
155. $(15, 3, 3, 20)$,
156. $(3, 7, 7, 24)$,
157. $(7, 3, 7, 24)$,
158. $(7, 7, 3, 24)$,
159. $(7, 7, 24, 3)$,
160. $(3, 7, 11, 20)$,
161. $(7, 3, 11, 20)$,
162. $(7, 11, 3, 20)$,
163. $(7, 7, 8, 19)$,
164. $(7, 7, 9, 18)$,
165. $(7, 7, 11, 16)$.

For $s \geqslant 3$,

140. $(3, 2^s-3, 2.2^s-2, 8.2^s-1)$,
141. $(3, 2^s-3, 8.2^s-1, 2.2^s-2)$,
142. $(3, 8.2^s-1, 2^s-3, 2.2^s-2)$,
143. $(8.2^s-1, 3, 2^s-3, 2.2^s-2)$,
144. $(3, 2^s-3, 2.2^s-1, 8.2^s-2)$,
145. $(3, 2^s-3, 8.2^s-2, 2.2^s-1)$,
146. $(3, 2.2^s-1, 2^s-3, 8.2^s-2)$,
147. $(2.2^s-1, 3, 2^s-3, 8.2^s-2)$,
148. $(2^s-1, 3, 2.2^s-3, 8.2^s-2)$,
149. $(2^s-1, 3, 8.2^s-3, 2.2^s-2)$,
150. $(3, 2^s-3, 4.2^s-2, 6.2^s-1)$,
151. $(3, 2^s-3, 4.2^s-1, 6.2^s-2)$,
152. $(3, 4.2^s-1, 2^s-3, 6.2^s-2)$,
153. $(4.2^s-1, 3, 2^s-3, 6.2^s-2)$,
154. $(2^s-1, 3, 4.2^s-3, 6.2^s-2)$,
155. $(7, 2.2^s-5, 2^s-3, 8.2^s-2)$,
156. $(7, 2.2^s-5, 8.2^s-3, 2^s-2)$,
157. $(7, 8.2^s-5, 2^s-3, 2.2^s-2)$,
158. $(7, 4.2^s-5, 2^s-3, 6.2^s-2)$,
159. $(7, 2.2^s-5, 2.2^s-3, 7.2^s-2)$,

160. $(7, 2.2^s - 5, 7.2^s - 3, 2.2^s - 2)$, 161. $(7, 2.2^s - 5, 3.2^s - 3, 6.2^s - 2)$,
162. $(7, 2.2^s - 5, 4.2^s - 3, 5.2^s - 2)$.

For $s = 3$,

163. $(7, 7, 9, 62)$, 164. $(7, 7, 57, 14)$, 165. $(7, 7, 25, 46)$.

For $s \geqslant 4$,

163. $(7, 2^s - 5, 2.2^s - 3, 8.2^s - 2)$, 164. $(7, 2^s - 5, 8.2^s - 3, 2.2^s - 2)$,
165. $(7, 2^s - 5, 4.2^s - 3, 6.2^s - 2)$.

We prove the theorem by proving the following.

Proposition 8.6.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+1}+2^s-3}$ is generated by the elements listed in Theorem 8.6.2.*

The proof of this proposition is based on the following lemmas.

Lemma 8.6.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 1\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 0 & 1 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 0 & 1 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 1\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 & 1\\ 1 & 1 & 0 & 1\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$\begin{aligned}&(15,19,5,2),(7,25,3,6),(3,15,19,4),(15,3,19,4),(15,19,3,4),\\ &(7,15,17,2),(15,7,17,2),(15,17,2,7),(15,17,7,2),(7,9,3,22),\\ &(15,17,3,6),(15,17,6,3),(3,7,15,16),(3,15,7,16),(3,15,16,7),\\ &(7,3,15,16),(7,15,3,16),(7,15,16,3),(15,3,7,16),(15,3,16,7),\\ &(15,7,3,16),(15,7,16,3),(15,16,3,7),(15,16,7,3),(7,11,19,4),\\ &(7,9,19,6),(7,11,7,16),(7,11,16,7).\end{aligned}$$

We prove the lemma for the matrices associated with the monomials

$$\begin{aligned}&(15,19,5,2),(7,25,3,6),(3,15,19,4),(15,19,3,4),(15,7,17,2),\\ &(15,17,7,2),(7,9,3,22),(15,17,3,6),(7,3,15,16),(7,15,3,16),\\ &(15,3,7,16),(15,16,3,7),(7,11,19,4),(7,9,19,6),(7,11,7,16).\end{aligned}$$

By a direct computation, we have

$$\begin{aligned}(15,19,5,2) &= Sq^1(15,15,7,3) + Sq^2(15,15,7,2) + Sq^4\big((15,15,5,2)\\ &\quad + (15,15,4,3)\big) + Sq^8\big((11,15,5,2) + (9,15,7,2)\\ &\quad + (8,15,7,3) + (11,15,4,3)\big) + (11,23,5,2)\\ &\quad + (9,23,7,2) + (15,17,7,2) + (8,23,7,3) + (15,16,7,3)\\ &\quad + (11,23,4,3) + (15,19,4,3) \mod \mathcal{L}_4(3;3;2;1;1),\\ (7,25,3,6) &= Sq^1\big((7,15,7,11) + (7,15,5,13)\big) + Sq^2\big((7,23,3,6)\\ &\quad + (7,15,7,10) + (3,15,11,10) + (2,15,11,11) + (7,15,6,11)\\ &\quad + (3,15,10,11) + (7,15,3,14)\big) + Sq^4\big((5,23,3,6) + (11,15,5,6)\\ &\quad + (5,15,7,10) + (3,15,13,6) + (3,15,7,12) + (4,15,7,11)\\ &\quad + (2,15,7,13) + (2,15,13,7) + (5,15,6,11) + (3,15,6,13)\\ &\quad + (3,15,12,7) + (5,15,3,14)\big) + Sq^8(7,15,5,6) + (7,23,3,8)\\ &\quad + (5,27,3,6) + (5,23,3,10) + (7,19,9,6) + (7,19,5,10)\end{aligned}$$

$$
\begin{aligned}
&+ (5,19,7,10) + (7,17,7,10) + (3,19,13,6) + (3,15,17,6)\\
&+ (3,19,7,12) + (3,15,7,16) + (4,19,7,11) + (2,19,13,7)\\
&+ (2,15,17,7) + (2,19,7,13) + (2,15,7,17) + (7,16,7,11)\\
&+ (5,19,6,11) + (3,19,6,13) + (3,15,6,17) + (3,19,12,7)\\
&+ (3,15,16,7) + (7,17,6,11) + (7,17,3,14) + (7,15,3,16)\\
&+ (5,19,3,14) + (5,15,3,18) \mod \mathcal{L}_4(3;3;2;1;1),\\
(3,15,19,4) &= Sq^1(3,15,15,7) + Sq^2(2,15,15,7) + Sq^4\big((3,15,15,4)\\
&+ (2,15,15,5)\big) + Sq^8\big((3,11,15,4) + (3,8,15,7)\big) + (3,11,23,4)\\
&+ (2,17,15,7) + (2,15,17,7) + (2,19,15,5) + (2,15,19,5)\\
&+ (3,8,23,7) + (3,15,16,7) \mod \mathcal{L}_4(3;3;2;1;1),\\
(15,19,3,4) &= Sq^1(15,15,3,7) + Sq^2(15,15,2,7)\\
&+ Sq^4\big((15,15,3,4) + (15,15,2,5)\big) + Sq^8\big((11,15,3,4)\\
&+ (8,15,3,7) + (9,15,2,7) + (11,15,2,5)\big) + (11,23,3,4)\\
&+ (9,23,2,7) + (15,17,2,7) + (11,23,2,5) + (15,19,2,5)\\
&+ (8,23,3,7) + (15,16,3,7) \mod \mathcal{L}_4(3;3;2;1;1),\\
(15,7,17,2) &= Sq^1(15,7,15,3) + Sq^2(15,7,15,2)\\
&+ Sq^4\big((15,5,15,2) + (15,4,15,3)\big) + Sq^8\big((9,7,15,2)\\
&+ (11,5,15,2) + (8,7,15,3) + (11,4,15,3)\big) + (9,7,23,2)\\
&+ (11,5,23,2) + (15,5,19,2) + (8,7,23,3) + (15,7,16,3)\\
&+ (11,4,23,3) + (15,4,19,3) \mod \mathcal{L}_4(3;3;2;1;1),\\
(15,17,7,2) &= Sq^1\big((15,7,11,7) + (15,7,9,9) + (15,7,7,11)\\
&+ (15,7,5,13)\big) + Sq^2\big((15,11,11,2) + (15,7,10,7) + (15,7,7,10)\\
&+ (15,7,6,11) + (15,7,3,14) + (15,3,11,10)\big) + Sq^4\big((15,13,7,2)\\
&+ (15,7,13,2) + (15,7,11,4) + (15,4,11,7) + (15,5,10,7)\\
&+ (15,5,7,10) + (15,4,7,11) + (15,5,6,11) + (15,5,3,14)\\
&+ (15,3,13,6) + (15,3,7,12)\big) + Sq^8\big((11,13,7,2) + (11,7,13,2)\\
&+ (11,7,11,4) + (8,7,11,7) + (11,4,11,7) + (9,7,10,7)\\
&+ (11,5,10,7) + (9,7,7,10) + (11,5,7,10) + (8,7,7,11)\\
&+ (11,4,7,11) + (9,7,6,11) + (11,5,6,11) + (9,7,3,14)\\
&+ (11,5,3,14) + (11,3,13,6) + (11,3,7,12)\big) + (11,21,7,2)\\
&+ (11,7,21,2) + (15,7,17,2) + (11,7,19,4) + (8,7,19,7)\\
&+ (11,4,19,7) + (9,7,18,7) + (11,5,18,7) + (9,7,7,18)\\
&+ (11,5,7,18) + (8,7,7,19) + (11,4,7,19) + (9,7,6,19)\\
&+ (11,5,6,19) + (9,7,3,22) + (11,5,3,22) + (15,5,3,18)
\end{aligned}
$$

$$\begin{aligned}
&+ (15,7,3,16) + (11,3,21,6) + (15,3,17,6)\\
&+ (11,3,7,20) + (15,3,7,16) \mod \mathcal{L}_4(3;3;2;1;1),\\
(7,9,3,22) &= Sq^1\big((7,7,5,21) + (7,7,11,15)\big) + Sq^2\big((7,7,3,22)\\
&+ (7,7,6,19) + (7,3,6,23) + (7,7,10,15) + (3,11,10,15)\\
&+ (2,11,11,5) + (7,2,7,23)\big) + Sq^4\big((5,7,3,22)\\
&+ (5,7,6,19) + (11,5,6,15) + (5,3,6,23) + (5,7,10,15)\\
&+ (3,13,6,15) + (3,7,12,15) + (4,7,11,15) + (2,13,7,15)\\
&+ (2,7,13,15) + (11,4,7,15) + (5,2,7,23)\big)\\
&+ Sq^8\big((7,5,6,15) + (7,4,7,15)\big) + (7,7,3,24) + (5,11,3,22)\\
&+ (5,7,3,26) + (7,7,8,19) + (5,11,6,19) + (5,7,10,19)\\
&+ (7,5,10,19) + (7,3,8,23) + (5,3,10,23) + (5,3,6,27)\\
&+ (7,7,10,17) + (5,7,10,19) + (3,17,6,15) + (3,13,6,19)\\
&+ (3,7,16,15) + (3,7,12,19) + (7,7,11,16) + (4,7,11,19)\\
&+ (2,17,7,15) + (2,13,7,19) + (2,7,17,15) + (2,7,13,19)\\
&+ (7,4,11,19) + (7,2,9,23) + (7,2,7,25) + (7,8,7,19)\\
&+ (5,2,11,23) + (5,2,7,27) + (7,3,6,25) \mod \mathcal{L}_4(3;3;2;1;1),\\
(15,17,3,6) &= Sq^1\big((15,7,7,11) + (15,7,5,13) + (15,3,11,11)\big)\\
&+ Sq^2\big((15,11,3,10) + (15,7,7,10) + (15,7,6,11) + (15,7,3,14)\\
&+ (15,3,11,10) + (15,3,10,11)\big) + Sq^4\big(15,13,3,6) + (15,7,3,12)\\
&+ (15,7,5,10) + (15,5,7,10) + (15,4,7,11) + (15,5,6,11)\\
&+ (15,5,3,14) + (15,3,13,6) + (15,3,6,13)\big) + Sq^8\big((11,13,3,6)\\
&+ (11,7,3,12) + (11,7,5,10) + (9,7,7,10) + (11,5,7,10)\\
&+ (8,7,7,11) + (11,4,7,11) + (9,7,6,11) + (11,5,6,11)\\
&+ (9,7,3,14) + (11,5,3,14) + (11,3,13,6) + (11,3,6,13)\big)\\
&+ (11,21,3,6) + (11,7,3,20) + (15,7,3,16) + (11,7,5,18)\\
&+ (9,7,7,18) + (11,5,7,18) + (8,7,7,19) + (11,4,7,19)\\
&+ (9,7,6,19) + (11,5,6,19) + (9,7,3,22) + (11,5,3,22)\\
&+ (15,5,3,18) + (15,7,3,16) + (15,3,17,6) + (15,3,6,17)\\
&+ (11,3,21,6) + (11,3,6,21) \mod \mathcal{L}_4(3;3;2;1;1),\\
(7,3,15,16) &= Sq^1(7,3,15,15) + Sq^2(7,2,15,15) + Sq^4\big((4,3,15,15)\\
&+ (5,2,15,15)\big) + Sq^8(7,3,8,15) + (4,3,19,15) + (4,3,15,19)\\
&+ (7,2,17,15) + (7,2,15,17) + (5,2,19,15)\\
&+ (5,2,15,19) + (7,3,8,23) \mod \mathcal{L}_4(3;3;2;1;1),
\end{aligned}$$

$$\begin{aligned}(15,3,7,16) &= Sq^1(15,3,7,15) + Sq^2(15,2,7,15) + Sq^4\big((15,2,5,15)\\ &\quad + (15,3,4,15)\big) + Sq^8\big((8,3,7,15) + (9,2,7,15) + (11,2,5,15)\\ &\quad + (11,3,4,15)\big) + (8,3,7,23) + (9,2,7,23)\\ &\quad + (15,2,7,17) + (11,2,5,23) + (15,2,5,19)\\ &\quad + (11,3,4,23) + (15,3,4,19) \mod \mathcal{L}_4(3;3;2;1;1),\\ (15,16,3,7) &= Sq^1(15,9,5,11) + Sq^2\big((15,10,3,11) + (15,3,10,11)\big)\\ &\quad + Sq^4\big((15,12,3,7) + (15,6,3,13) + (15,5,6,11) + (15,3,6,13)\\ &\quad + (15,3,12,7)\big) + Sq^8\big((11,12,3,7) + (11,6,3,13) + (11,5,6,11)\\ &\quad + (11,3,6,13) + (11,3,12,7)\big) + (11,20,3,7) + (15,6,3,17)\\ &\quad + (11,6,3,21) + (11,5,6,19) + (11,3,6,21) + (15,3,6,17)\\ &\quad + (11,3,20,7) + (15,3,16,7) \mod \mathcal{L}_4(3;3;2;1;1),\\ (7,11,19,4) &= Sq^1\big((7,7,23,3) + (7,11,15,7) + (7,5,23,5)\big)\\ &\quad + Sq^2\big((2,11,15,11) + (7,10,15,7) + (3,10,15,11) + (7,3,23,6)\big)\\ &\quad + Sq^4\big((11,7,15,4) + (4,7,23,3) + (4,11,15,7) + (2,7,15,13)\\ &\quad + (2,13,15,7) + (5,10,15,7) + (3,6,15,13) + (3,12,15,7)\\ &\quad + (11,6,15,5) + (5,3,23,6)\big) + Sq^8\big((7,7,15,4) + (7,6,15,5)\big)\\ &\quad + (7,7,19,8) + (4,11,23,3) + (4,7,27,3) + (7,8,23,3)\\ &\quad + (7,7,24,3) + (4,11,19,7) + (2,7,19,13) + (2,7,15,17)\\ &\quad + (2,17,15,7) + (2,13,19,7) + (7,11,16,7) + (5,10,19,7)\\ &\quad + (3,6,19,13) + (3,6,15,17) + (3,16,15,7) + (3,12,19,7)\\ &\quad + (7,10,17,7) + (7,10,19,5) + (7,6,19,9) + (7,3,25,6)\\ &\quad + (7,3,23,8) + (5,3,27,6) + (5,3,23,10) \mod \mathcal{L}_4(3;3;2;1;1),\\ (7,9,19,6) &= Sq^1\big((7,7,15,11) + (7,3,23,7)\big) + Sq^2\big((7,3,23,6)\\ &\quad + (7,7,15,10) + (3,11,15,10) + (2,11,15,11)\big)\\ &\quad + Sq^4\big((11,5,15,6) + (5,3,23,6) + (5,7,15,10) + (3,13,15,6)\\ &\quad + (3,7,15,12) + (4,7,15,11) + (2,13,15,7) + (2,7,15,13)\\ &\quad + (11,4,15,7) + (4,3,23,7)\big) + Sq^8\big((7,5,15,6) + (7,4,15,7)\big)\\ &\quad + (7,5,19,10) + (7,3,25,6) + (7,3,23,8) + (5,3,27,6)\\ &\quad + (5,3,23,10) + (7,7,17,10) + (5,7,19,10) + (3,17,15,6)\\ &\quad + (3,13,19,6) + (3,7,19,12) + (3,7,15,16) + (7,7,16,11)\\ &\quad + (4,7,19,11) + (2,17,15,7) + (2,13,19,7) + (2,7,19,13)\\ &\quad + (2,7,15,17) + (7,4,19,11) + (7,8,19,7) + (7,3,24,7)\\ &\quad + (7,3,23,8) + (4,3,27,7) + (4,3,23,11) \mod \mathcal{L}_4(3;3;2;1;1),\end{aligned}$$

$$\begin{aligned}(7,11,7,16) &= Sq^1\big((7,11,7,15)+(7,7,3,23)\big)+Sq^2\big((2,11,11,15)\\
&\quad+(7,10,7,15)+(3,10,11,15)+(7,6,3,23)\big)\\
&\quad+Sq^4\big((4,11,7,15)+(2,7,13,15)+(2,13,7,15)\\
&\quad+(5,10,7,15)+(3,6,13,15)+(3,12,7,15)+(11,6,5,15)\\
&\quad+(5,6,3,23)+(11,7,4,15)+(4,7,3,23)\big)+Sq^8\big((7,6,5,15)\\
&\quad+(7,7,4,15)\big)+(4,11,7,19)+(2,7,17,15)+(2,7,13,19)\\
&\quad+(2,17,7,15)+(2,13,7,19)+(7,10,7,17)+(5,10,7,19)\\
&\quad+(3,6,17,15)+(3,6,13,19)+(3,16,7,15)+(3,12,7,19)\\
&\quad+(7,10,5,19)+(7,6,9,19)+(7,6,3,25)+(5,10,3,23)\\
&\quad+(5,6,3,27)+(7,11,4,19)+(7,7,8,19)+(7,7,3,24)\\
&\quad+(4,11,3,23)+(4,7,3,27) \mod \mathcal{L}_4(3;3;2;1;1).\end{aligned}$$

The lemma is proved. □

Lemma 8.6.5. *The following matrix is strictly inadmissible*

$$\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&1&0\\1&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}.$$

Proof. The monomial corresponding to the above matrix is (15,15,23,32). By a direct computation, we have

$$\begin{aligned}(15,15,23,32) &= Sq^1(15,15,23,31)+Sq^2\big((15,10,27,31)+(15,15,22,31)\\
&\quad+(15,11,26,31)\big)+Sq^4\big((4,23,23,31)+(15,12,23,31)+(7,20,23,31)\\
&\quad+(15,6,29,31)+(5,23,22,31)+(15,13,22,31)+(7,21,22,31)\\
&\quad+(15,7,28,31)\big)+Sq^8\big((8,15,23,31)+(4,27,15,31)+(4,15,27,31)\\
&\quad+(11,12,23,31)+(7,24,15,31)+(7,12,27,31)+(11,6,29,31)\\
&\quad+(9,15,22,31)+(5,15,26,31)+(5,27,14,31)+(11,13,22,31)\\
&\quad+(7,13,26,31)+(7,25,14,31)+(11,7,28,31)\big)+(8,15,23,39)\\
&\quad+(4,27,15,39)+(4,35,15,31)+(4,15,35,31)+(4,15,27,39)\\
&\quad+(15,12,23,35)+(11,12,23,39)+(7,24,15,39)+(7,32,15,31)\\
&\quad+(7,12,35,31)+(7,12,27,39)+(11,6,37,31)+(11,6,29,39)\\
&\quad+(15,6,33,31)+(15,6,29,35)+(15,15,22,33)+(9,15,22,39)\\
&\quad+(5,15,34,31)+(5,15,26,39)+(5,35,14,31)+(5,27,14,39)\\
&\quad+(15,13,22,35)+(11,13,22,39)+(7,13,34,31)+(7,13,26,39)\end{aligned}$$

$$+ (7,33,14,31) + (7,25,14,39) + (15,7,32,31) + (15,7,28,35)$$
$$+ (11,7,36,31) + (11,7,28,39) \mod \mathcal{L}_4(3;3;3;2;1;1).$$

The lemma follows. □

Using the results in Section 7, Lemmas 3.5, 3.6, 5.10, 6.3.4, 6.3.5, 8.5.4, 8.5.5, 8.6.4, 8.6.5 and Theorem 2.5, we obtain Proposition 8.6.3.

Now, we prove that the classes listed in Theorem 8.6.2 are linearly independent.

Proposition 8.6.6. *The elements* $[a_{2,1,2,j}], 1 \leqslant j \leqslant 165$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{41}$.

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{165} \gamma_j [a_{2,1,2,j}] = 0, \tag{8.6.6.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.6.6.1), we obtain

$$\begin{aligned}
&\gamma_1[3,7,31] + \gamma_2[3,31,7] + \gamma_{15}[7,3,31] + \gamma_{16}[7,31,3]\\
&\quad + \gamma_{29}[31,3,7] + \gamma_{30}[31,7,3] + \gamma_{37}[3,15,23] + \gamma_{44}[15,3,23]\\
&\quad + \gamma_{45}[15,23,3] + \gamma_{55}[7,7,27] + \gamma_{56}[7,27,7] + \gamma_{71}[7,11,23]\\
&\quad + \gamma_{81}[7,15,19] + \gamma_{91}[15,7,19] + \gamma_{92}[15,19,7] = 0,\\
&\gamma_3[3,7,31] + \gamma_5[3,31,7] + \gamma_{\{13,140\}}[7,3,31] + \gamma_{18}[7,31,3]\\
&\quad + \gamma_{\{26,145\}}[31,3,7] + \gamma_{\{28,159\}}[31,7,3] + \gamma_{38}[3,15,23] + \gamma_{\{43,152\}}[15,3,23]\\
&\quad + \gamma_{49}[15,23,3] + \gamma_{\{57,72\}}[7,7,27] + \gamma_{96}[7,27,7] + \gamma_{72}[7,11,23]\\
&\quad + \gamma_{\{82,96\}}[7,15,19] + \gamma_{\{63,90,163\}}[15,7,19] + \gamma_{63}[15,19,7] = 0,\\
&\gamma_4[3,7,31] + \gamma_6[3,31,7] + \gamma_{\{14,141\}}[7,3,31] + \gamma_{\{17,142\}}[7,31,3]\\
&\quad + \gamma_{\{25,144,157\}}[31,3,7] + \gamma_{\{27,146,158\}}[31,7,3] + \gamma_{95}[7,27,7] + \gamma_{94}[7,11,23]\\
&\quad + \gamma_{39}[3,15,23] + \gamma_{\{47,153\}}[15,3,23] + \gamma_{\{48,154\}}[15,23,3] + \gamma_{\{83,95\}}[7,15,19]\\
&\quad + \gamma_{\{58,94,148\}}[7,7,27] + \gamma_{\{62,74,156,160,165\}}[15,7,19] + \gamma_{\{62,156\}}[15,19,7] = 0,\\
&\gamma_{\{19,67,140\}}[3,7,31] + \gamma_{\{32,78,115,135,145\}}[3,31,7] + \gamma_7[7,3,31]\\
&\quad + \gamma_{\{35,159\}}[7,31,3] + \gamma_{10}[31,3,7] + \gamma_{23}[31,7,3] + \gamma_{40}[15,3,23]\\
&\quad + \gamma_{\{46,105,121,152\}}[3,15,23] + \gamma_{53}[15,23,3] + \gamma_{59}[7,7,27] + \gamma_{65}[7,27,7]\\
&\quad + \gamma_{73}[7,11,23] + \gamma_{\{93,163\}}[7,15,19] + \gamma_{87}[15,7,19] + \gamma_{100}[15,19,7] = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_{\{20,68,141\}}[3,7,31] + a_1[3,31,7] + \gamma_8[7,3,31] + \gamma_{11}[31,3,7] + \gamma_{41}[15,3,23]\\
&\quad + \gamma_{\{34,147,150,158,162\}}[7,31,3] + \gamma_{\{22,143\}}[31,7,3] + \gamma_{\{50,106,126,153\}}[3,15,23]\\
&\quad + \gamma_{\{52,155\}}[15,23,3] + \gamma_{\{64,157\}}[7,27,7] + \gamma_{\{75,161,165\}}[7,15,19]\\
&\quad + \gamma_{\{60,149\}}[7,7,27] + \gamma_{97}[7,11,23] + \gamma_{88}[15,7,19] + \gamma_{99}[15,19,7] = 0,\\
&a_2[3,7,31] + \gamma_{\{21,69,142\}}[3,31,7] + a_3[7,3,31] + \gamma_9[7,31,3] + \gamma_{101}[15,7,19]\\
&\quad + \gamma_{\{24,70,143\}}[31,3,7] + \gamma_{12}[31,7,3] + \gamma_{42}[15,23,3] + \gamma_{98}[7,15,19]\\
&\quad + \gamma_{\{51,107,127,131,154\}}[3,15,23] + a_4[7,7,27] + \gamma_{\{54,108,128,132,155\}}[15,3,23]\\
&\quad + \gamma_{\{61,120,150\}}[7,27,7] + \gamma_{\{76,137,138,139,162\}}[7,11,23] + \gamma_{89}[15,19,7] = 0,
\end{aligned}$$

where

$$a_1 = \gamma_{\{31,79,114,134,144,156\}},\quad a_2 = \gamma_{\{33,77,116,118,123,133,146,148,156,160\}},$$
$$a_3 = \gamma_{\{36,80,117,119,124,136,147,149,157,161\}},\quad a_4 = \gamma_{\{66,151,158,159,163,164,165\}}.$$

Computing from these equalities, we obtain

$$\begin{cases}
a_i = 0,\ i = 1,2,3,4,\\
\gamma_j = 0,\ j = 1,2,\ldots,12,15,16,18,23,29,30,\\
37,\ldots,42,44,45,49,53,55,56,57,59,63,65,\\
71,72,73,81,82,83,87,88,89,91,92,94,\ldots,101,\\
\gamma_{\{13,140\}} = \gamma_{\{26,145\}} = \gamma_{\{28,159\}} = \gamma_{\{43,152\}} = \gamma_{\{90,163\}} = 0,\\
\gamma_{\{14,141\}} = \gamma_{\{17,142\}} = \gamma_{\{47,153\}} = \gamma_{\{62,156\}} = \gamma_{\{48,154\}} = 0,\\
\gamma_{\{25,144,157\}} = \gamma_{\{27,146,158\}} = \gamma_{\{58,148\}} = \gamma_{\{19,67,140\}} = 0,\\
\gamma_{\{62,74,156,160,165\}} = \gamma_{\{32,78,115,135,145\}} = \gamma_{\{35,159\}} = 0,\\
\gamma_{\{46,105,121,152\}} = \gamma_{\{20,68,141\}} = \gamma_{\{34,147,150,158,162\}} = 0,\\
\gamma_{\{93,163\}} = \gamma_{\{22,143\}} = \gamma_{\{50,106,126,153\}} = \gamma_{\{52,155\}} = 0,\\
\gamma_{\{60,149\}} = \gamma_{\{75,161,165\}} = \gamma_{\{21,69,142\}} = \gamma_{\{24,70,143\}} = 0,\\
\gamma_{\{51,107,127,131,154\}} = \gamma_{\{54,108,128,132,155\}} = \gamma_{\{61,120,150\}} = 0,\\
\gamma_{\{64,157\}} = \gamma_{\{76,137,138,139,162\}} = 0.
\end{cases} \tag{8.6.6.2}$$

With the aid of (8.6.6.2), the homomorphisms g_1, g_2 send (8.6.6.1) to

$$\begin{aligned}
&\gamma_{19}[3,7,31] + \gamma_{32}[3,31,7] + \gamma_{67}[7,3,31] + \gamma_{\{28,86\}}[7,31,3]\\
&\quad + \gamma_{78}[31,3,7] + \gamma_{104}[31,7,3] + \gamma_{46}[3,15,23] + \gamma_{105}[15,3,23]\\
&\quad + \gamma_{111}[15,23,3] + \gamma_{112}[7,7,27] + \gamma_{115}[7,27,7] + \gamma_{121}[7,11,23]\\
&\quad + \gamma_{\{90,125\}}[7,15,19] + \gamma_{129}[15,7,19] + \gamma_{135}[15,19,7] = 0,\\
&\gamma_{20}[3,7,31] + \gamma_{\{25,31,144\}}[3,31,7] + \gamma_{\{27,34,85,146,147,158\}}[7,31,3]\\
&\quad + \gamma_{68}[7,3,31] + \gamma_{79}[31,3,7] + \gamma_{\{103,150\}}[31,7,3] + \gamma_{50}[3,15,23]\\
&\quad + \gamma_{106}[15,3,23] + \gamma_{\{110,162\}}[15,23,3] + \gamma_{113}[7,7,27]\\
&\quad + \gamma_{114}[7,27,7] + \gamma_{126}[7,11,23] + \gamma_{\{74,75,122,160,161,165\}}[7,15,19]\\
&\quad + \gamma_{130}[15,7,19] + \gamma_{134}[15,19,7] = 0.
\end{aligned}$$

These equalities imply

$$(8.6.6.3)\quad \begin{cases} \gamma_j = 0,\ j = 19, 20, 32, 46, 50, 67, 68, 78, 79, 104, 105, \\ 106, 111, 112, 113, 114, 115, 121, 126, 129, 130, 134, 135, \\ \gamma_{\{28,86\}} = \gamma_{\{90,125\}} = \gamma_{\{25,31,144\}} = \gamma_{\{27,34,85,146,147,158\}} = 0, \\ \gamma_{\{103,150\}} = \gamma_{\{110,162\}} = \gamma_{\{74,75,122,160,161,165\}} = 0. \end{cases}$$

With the aid of (8.6.6.2) and (8.6.6.3), the homomorphisms g_3, g_4 send (8.6.6.1) to

$$\begin{aligned} &\gamma_{21}[3,7,31] + \gamma_{\{27,33,66,146,158\}}[3,31,7] + \gamma_{69}[7,3,31] + a_5[7,31,3] \\ &\quad + \gamma_{\{77,151\}}[31,3,7] + a_6[31,7,3] + \gamma_{51}[3,15,23] + \gamma_{107}[15,3,23] \\ &\quad + a_7[15,23,3] + \gamma_{\{61,120,131,150\}}[7,7,27] + a_8[7,27,7] + a_9[7,15,19] \\ &\quad + \gamma_{\{127,131\}}[7,11,23] + \gamma_{\{28,118,137,139,151\}}[15,7,19] + a_{10}[15,19,7] = 0, \\ &\gamma_{24}[3,7,31] + a_{11}[3,31,7] + \gamma_{70}[7,3,31] + a_{12}[7,31,3] + a_{13}[31,7,3] \\ &\quad + \gamma_{\{80,120,137,151\}}[31,3,7] + \gamma_{54}[3,15,23] + \gamma_{108}[15,3,23] \\ &\quad + a_{14}[15,23,3] + \gamma_{132}[7,7,27] + a_{15}[7,27,7] + \gamma_{\{128,132\}}[7,11,23] \\ &\quad + a_{16}[7,15,19] + a_{17}[15,7,19] + a_{18}[15,19,7] = 0, \end{aligned}$$

where

$$\begin{aligned} a_5 &= \gamma_{\{22,24,25,36,52,54,64,70,84,117,128,144,147\}}, \quad a_6 = \gamma_{\{22,24,60,70,80,102,108,119,132\}}, \\ a_7 &= \gamma_{\{52,54,108,109,124,128,132,136,161\}}, \quad a_8 = \gamma_{\{66,74,90,116,123,158,160,164,165\}}, \\ a_9 &= \gamma_{\{74,76,90,123,138,160,162,164,165\}}, \quad a_{10} = \gamma_{\{28,118,133,151,164,165\}}, \\ a_{11} &= \gamma_{\{34,36,61,66,76,147,150,158,162\}}, \\ a_{12} &= \gamma_{\{13,17,21,31,33,48,51,62,69,84,85,103,110,116,127,144,146\}}, \\ a_{13} &= \gamma_{\{14,17,21,26,58,69,77,86,102,103,107,118,131\}}, \\ a_{14} &= \gamma_{\{43,47,48,51,107,109,110,122,123,125,127,131,133,160\}}, \\ a_{15} &= \gamma_{\{66,75,76,90,117,120,124,138,139,158,161,162,164,165\}}, \\ a_{16} &= \gamma_{\{75,76,90,124,139,161,162,164,165\}}, \quad a_{17} = \gamma_{\{28,61,119,139,150,151\}}, \\ a_{18} &= \gamma_{\{28,61,119,136,137,138,139,150,151,164,165\}}. \end{aligned}$$

From the above equalities, we obtain

$$(8.6.6.4)\quad \begin{cases} a_i = 0,\ i = 5, 6, \ldots, 18, \\ \gamma_j = 0,\ j = 21, 24, 51, 54, 69, 70, 107, 108, 128, 132, \\ \gamma_{\{27,33,66,146,158\}} = \gamma_{\{77,151\}} = \gamma_{\{127,131\}} = 0, \\ \gamma_{\{28,118,137,139,151\}} = \gamma_{\{61,120,131,150\}} = \gamma_{\{80,120,137,151\}} = 0. \end{cases}$$

With the aid of (8.6.6.2), (8.6.6.3) and (8.6.6.4), the homomorphism h sends (8.6.6.1) to

$$\begin{aligned}
&a_{19}[3,7,31] + \gamma_{\{28,80,102,119\}}[3,31,7] + a_{20}[7,3,31] + \gamma_{80}[7,31,3]\\
&\quad + \gamma_{61}[31,3,7] + \gamma_{120}[31,7,3] + a_{21}[3,15,23] + \gamma_{76}[15,3,23]\\
&\quad + \gamma_{137}[15,23,3] + a_{22}[7,7,27] + a_{23}[7,27,7] + \gamma_{\{61,103,120,127,138\}}[15,7,19]\\
&\quad + a_{24}[7,11,23] + a_{25}[7,15,19] + \gamma_{\{61,103,120,127,139\}}[15,19,7] = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_{19} = \gamma_{\{25,31,34,36,84,117,144,147\}}, \quad a_{20} = \gamma_{\{27,33,34,36,66,85,146,147,158\}},\\
&a_{21} = \gamma_{\{75,90,109,124,136,161\}}, \quad a_{22} = \gamma_{\{66,76,110,116,117,138,158\}},\\
&a_{23} = \gamma_{\{28,77,118,119,137,139\}}, \quad a_{25} = \gamma_{\{133,136,137,139,164,165\}},\\
&a_{24} = \gamma_{\{74,75,76,90,110,122,123,124,138,160,161,164,165\}}.
\end{aligned}$$

From the above equalities, it implies

$$\text{(8.6.6.5)} \qquad \begin{cases} a_i = 0, \ i = 19, 20, \ldots, 25,\\ \gamma_j = 0, \ j = 61, 76, 80, 120, 137,\\ \gamma_{\{28,102,119\}} = \gamma_{\{103,127,138\}} = \gamma_{\{103,127,139\}} = 0. \end{cases}$$

Combining (8.6.6.2), (8.6.6.3), (8.6.6.4) and (8.6.6.5), we get $\gamma_j = 0$ for $1 \leqslant j \leqslant 165$. The proposition is proved. □

Proposition 8.6.7. *For $s \geqslant 3$, the elements $[a_{2,1,s,j}], 1 \leqslant j \leqslant 165$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+1}+2^s-3}$.*

Proof. Suppose that there is a linear relation

$$\text{(8.6.7.1)} \qquad \sum_{j=1}^{165} \gamma_j [a_{2,1,s,j}] = 0,$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.6.7.1), we obtain

$$\begin{aligned}
&\gamma_1 w_{2,1,s,1} + \gamma_2 w_{2,1,s,2} + \gamma_{15} w_{2,1,s,3} + \gamma_{16} w_{2,1,s,4} + \gamma_{29} w_{2,1,s,5} + \gamma_{30} w_{2,1,s,6}\\
&\quad + \gamma_{37} w_{2,1,s,7} + \gamma_{44} w_{2,1,s,8} + \gamma_{45} w_{2,1,s,9} + \gamma_{55} w_{2,1,s,10} + \gamma_{56} w_{2,1,s,11}\\
&\quad + \gamma_{71} w_{2,1,s,12} + \gamma_{81} w_{2,1,s,13} + \gamma_{91} w_{2,1,s,14} + \gamma_{92} w_{2,1,s,15} = 0,\\
&\gamma_3 w_{2,1,s,1} + \gamma_5 w_{2,1,s,2} + \gamma_{13} w_{2,1,s,3} + \gamma_{18} w_{2,1,s,4} + \gamma_{26} w_{2,1,s,5} + \gamma_{28} w_{2,1,s,6}\\
&\quad + \gamma_{38} w_{2,1,s,7} + \gamma_{43} w_{2,1,s,8} + \gamma_{49} w_{2,1,s,9} + \gamma_{\{57,72\}} w_{2,1,s,10} + \gamma_{96} w_{2,1,s,11}\\
&\quad + \gamma_{72} w_{2,1,s,12} + \gamma_{\{82,96\}} w_{2,1,s,13} + \gamma_{\{63,90\}} w_{2,1,s,14} + \gamma_{63} w_{2,1,s,15} = 0,\\
&\gamma_4 w_{2,1,s,1} + \gamma_6 w_{2,1,s,2} + \gamma_{14} w_{2,1,s,3} + \gamma_{17} w_{2,1,s,4} + \gamma_{25} w_{2,1,s,5} + \gamma_{27} w_{2,1,s,6}\\
&\quad + \gamma_{39} w_{2,1,s,7} + \gamma_{47} w_{2,1,s,8} + \gamma_{48} w_{2,1,s,9} + \gamma_{\{58,94\}} w_{2,1,s,10} + \gamma_{95} w_{2,1,s,11}\\
&\quad + \gamma_{94} w_{2,1,s,12} + \gamma_{\{83,95\}} w_{2,1,s,13} + \gamma_{\{62,74\}} w_{2,1,s,14} + \gamma_{62} w_{2,1,s,15} = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_{19}w_{2,1,s,1} + \gamma_{32}w_{2,1,s,2} + \gamma_{7}w_{2,1,s,3} + \gamma_{35}w_{2,1,s,4} + \gamma_{10}w_{2,1,s,5} + \gamma_{23}w_{2,1,s,6}\\
&\quad + \gamma_{46}w_{2,1,s,7} + \gamma_{40}w_{2,1,s,8} + \gamma_{53}w_{2,1,s,9} + \gamma_{59}w_{2,1,s,10} + \gamma_{65}w_{2,1,s,11}\\
&\quad + \gamma_{73}w_{2,1,s,12} + \gamma_{93}w_{2,1,s,13} + \gamma_{87}w_{2,1,s,14} + \gamma_{100}w_{2,1,s,15} = 0,\\
&\gamma_{20}w_{2,1,s,1} + \gamma_{31}w_{2,1,s,2} + \gamma_{8}w_{2,1,s,3} + \gamma_{34}w_{2,1,s,4} + \gamma_{11}w_{2,1,s,5} + \gamma_{22}w_{2,1,s,6}\\
&\quad + \gamma_{50}w_{2,1,s,7} + \gamma_{41}w_{2,1,s,8} + \gamma_{52}w_{2,1,s,9} + \gamma_{60}w_{2,1,s,10} + \gamma_{64}w_{2,1,s,11}\\
&\quad + \gamma_{97}w_{2,1,s,12} + \gamma_{75}w_{2,1,s,13} + \gamma_{88}w_{2,1,s,14} + \gamma_{99}w_{2,1,s,15} = 0,\\
&\gamma_{33}w_{2,1,s,1} + \gamma_{21}w_{2,1,s,2} + \gamma_{36}w_{2,1,s,3} + \gamma_{9}w_{2,1,s,4} + \gamma_{24}w_{2,1,s,5} + \gamma_{12}w_{2,1,s,6}\\
&\quad + \gamma_{51}w_{2,1,s,7} + \gamma_{54}w_{2,1,s,8} + \gamma_{42}w_{2,1,s,9} + \gamma_{66}w_{2,1,s,10} + \gamma_{61}w_{2,1,s,11}\\
&\quad + \gamma_{76}w_{2,1,s,12} + \gamma_{98}w_{2,1,s,13} + \gamma_{101}w_{2,1,s,14} + \gamma_{89}w_{2,1,s,15} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\gamma_j = 0, j = 1, \dots, 66, 71, \dots, 76, 81, 82, 83, 87, \dots, 101. \tag{8.6.7.2}$$

With the aid of (8.6.7.2), the homomorphisms g_1, g_2, g_3, g_4 send (8.6.7.1) to

$$\begin{aligned}
&\gamma_{140}w_{2,1,s,1} + \gamma_{145}w_{2,1,s,2} + \gamma_{67}w_{2,1,s,3} + \gamma_{86}w_{2,1,s,4}\\
&\quad + \gamma_{78}w_{2,1,s,5} + \gamma_{104}w_{2,1,s,6} + \gamma_{150}w_{2,1,s,7} + \gamma_{105}w_{2,1,s,8}\\
&\quad + \gamma_{111}w_{2,1,s,9} + \gamma_{112}w_{2,1,s,10} + \gamma_{115}w_{2,1,s,11} + \gamma_{121}w_{2,1,s,12}\\
&\quad + \gamma_{125}w_{2,1,s,13} + \gamma_{129}w_{2,1,s,14} + \gamma_{135}w_{2,1,s,15} = 0,\\
&\gamma_{141}w_{2,1,s,1} + \gamma_{144}w_{2,1,s,2} + \gamma_{68}w_{2,1,s,3} + \gamma_{85}w_{2,1,s,4}\\
&\quad + \gamma_{79}w_{2,1,s,5} + \gamma_{103}w_{2,1,s,6} + \gamma_{151}w_{2,1,s,7} + \gamma_{106}w_{2,1,s,8}\\
&\quad + \gamma_{110}w_{2,1,s,9} + \gamma_{113}w_{2,1,s,10} + \gamma_{114}w_{2,1,s,11} + \gamma_{126}w_{2,1,s,12}\\
&\quad + \gamma_{122}w_{2,1,s,13} + \gamma_{130}w_{2,1,s,14} + \gamma_{134}w_{2,1,s,15} = 0,\\
&\gamma_{142}w_{2,1,s,1} + \gamma_{146}w_{2,1,s,2} + \gamma_{69}w_{2,1,s,3} + a_1w_{2,1,s,4} + \gamma_{77}w_{2,1,s,5}\\
&\quad + a_2w_{2,1,s,6} + \gamma_{152}w_{2,1,s,7} + \gamma_{107}w_{2,1,s,8} + a_3w_{2,1,s,9}\\
&\quad + \gamma_{131}w_{2,1,s,10} + \gamma_{\{116,123\}}w_{2,1,s,11} + \gamma_{\{127,131\}}w_{2,1,s,12}\\
&\quad + \gamma_{123}w_{2,1,s,13} + \gamma_{118}w_{2,1,s,14} + \gamma_{\{118,133\}}w_{2,1,s,15} = 0,\\
&\gamma_{143}w_{2,1,s,1} + \gamma_{147}w_{2,1,s,2} + \gamma_{70}w_{2,1,s,3} + a_4w_{2,1,s,4} + \gamma_{80}w_{2,1,s,5}\\
&\quad + a_5w_{2,1,s,6} + \gamma_{153}w_{2,1,s,7} + \gamma_{108}w_{2,1,s,8} + a_6w_{2,1,s,9}\\
&\quad + \gamma_{132}w_{2,1,s,10} + \gamma_{\{117,124\}}w_{2,1,s,11} + \gamma_{\{128,132\}}w_{2,1,s,12}\\
&\quad + \gamma_{124}w_{2,1,s,13} + \gamma_{119}w_{2,1,s,14} + \gamma_{\{119,136\}}w_{2,1,s,15} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases} \gamma_{\{84,163\}}, & s = 3,\\ \gamma_{84}, & s \geqslant 4, \end{cases} \quad a_4 = \begin{cases} \gamma_{\{120,138,148,155,157,158,159,163\}}, & s = 3,\\ \gamma_{148}, & s \geqslant 4, \end{cases}$$

$$a_2 = \begin{cases} \gamma_{\{102,164\}}, & s = 3,\\ \gamma_{102}, & s \geqslant 4, \end{cases} \quad a_5 = \begin{cases} \gamma_{\{120,137,139,149,156,157,160,164\}}, & s = 3,\\ \gamma_{149}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{109,165\}}, & s = 3, \\ \gamma_{109}, & s \geqslant 4, \end{cases} \quad a_6 = \begin{cases} \gamma_{\{137,138,139,154,158,161,162,165\}}, & s = 3, \\ \gamma_{154}, & s \geqslant 4. \end{cases}$$

These equalities imply

$$\begin{cases} a_i = 0, i = 1, 2, \ldots, 6, \\ \gamma_j = 0,\ j = 67, \ldots, 70, 77, \ldots, 80, 85, 86, 103, \ldots, 108, \\ 110, \ldots, 119, 121, \ldots, 136, 140, \ldots, 147, 150, \ldots, 153. \end{cases} \tag{8.6.7.3}$$

With the aid of (8.6.7.2), (8.6.7.3) and (8.6.7.4), the homomorphism h sends (8.6.7.1) to

$$\begin{aligned} &a_7 w_{2,1,s,1} + a_8 w_{2,1,s,2} + \gamma_{155} w_{2,1,s,3} + \gamma_{156} w_{2,1,s,4} + \gamma_{157} w_{2,1,s,5} + \gamma_{120} w_{2,1,s,6} \\ &\quad + a_9 w_{2,1,s,7} + \gamma_{158} w_{2,1,s,8} + \gamma_{137} w_{2,1,s,9} + \gamma_{159} w_{2,1,s,10} + \gamma_{160} w_{2,1,s,11} \\ &\quad + \gamma_{161} w_{2,1,s,12} + \gamma_{162} w_{2,1,s,13} + \gamma_{138} w_{2,1,s,14} + \gamma_{139} w_{2,1,s,15} = 0, \end{aligned}$$

where

$$a_7 = \begin{cases} \gamma_{148}, & s = 3, \\ \gamma_{163}, & s \geqslant 4, \end{cases} \quad a_8 = \begin{cases} \gamma_{149}, & s = 3, \\ \gamma_{164}, & s \geqslant 4, \end{cases} \quad a_9 = \begin{cases} \gamma_{154}, & s = 3, \\ \gamma_{165}, & s \geqslant 4. \end{cases}$$

From the above equalities, it implies

$$a_7 = a_8 = a_9 = 0, \gamma_j = 0,\ j = 120, 137, 138, 139, 155, \ldots, 162. \tag{8.6.7.4}$$

Combining (8.6.7.2), (8.6.7.3) and (8.6.7.4), we get $\gamma_j = 0$ for $1 \leqslant j \leqslant 165$. The proposition is proved. □

8.7. **The case** $s \geqslant 2, t = 1$ **and** $u \geqslant 3$**.**

According to Kameko [8], for $s \geqslant 2$ and $u \geqslant 3$, the dimension of the space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+u+1}+2^{s+1}+2^s-3}$ is 14 with a basis given by the following classes:

$$\begin{aligned} w_{u,1,s,1} &= [2^s - 1, 2^{s+1} - 1, 2^{s+u+1} - 1], \\ w_{u,1,s,2} &= [2^s - 1, 2^{s+u+1} - 1, 2^{s+1} - 1], \\ w_{u,1,s,3} &= [2^{s+1} - 1, 2^s - 1, 2^{s+u+1} - 1], \\ w_{u,1,s,4} &= [2^{s+1} - 1, 2^{s+u+1} - 1, 2^s - 1], \\ w_{u,1,s,5} &= [2^{s+u+1} - 1, 2^s - 1, 2^{s+1} - 1], \\ w_{u,1,s,6} &= [2^{s+u+1} - 1, 2^{s+1} - 1, 2^s - 1], \\ w_{u,1,s,7} &= [2^s - 1, 2^{s+2} - 1, 2^{s+u+1} - 2^{s+1} - 1], \\ w_{u,1,s,8} &= [2^{s+2} - 1, 2^s - 1, 2^{s+u+1} - 2^{s+1} - 1], \\ w_{u,1,s,9} &= [2^{s+2} - 1, 2^{s+u+1} - 2^{s+1} - 1, 2^s - 1], \\ w_{u,1,s,10} &= [2^{s+1} - 1, 2^{s+1} - 1, 2^{s+u+1} - 2^s - 1], \\ w_{u,1,s,11} &= [2^{s+1} - 1, 2^{s+u+1} - 2^s - 1, 2^{s+1} - 1], \\ w_{u,1,s,12} &= [2^{s+1} - 1, 2^{s+1} + 2^s - 1, 2^{s+u+1} - 2^{s+1} - 1], \\ w_{u,1,s,13} &= [2^{s+1} - 1, 2^{s+2} - 1, 2^{s+u+1} - 2^{s+1} - 2^s - 1], \\ w_{u,1,s,14} &= [2^{s+2} - 1, 2^{s+1} - 1, 2^{s+u+1} - 2^{s+1} - 2^s - 1]. \end{aligned}$$

So, we easily obtain

Proposition 8.7.1. *For any positive integers $s \geqslant 2$ and $u \geqslant 3$, we have*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+u+1}+2^{s+1}+2^s-3} = 56.$$

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+u+1}+2^{s+1}+2^s-3}$. We have

Theorem 8.7.2. *For $s \geqslant 2, u \geqslant 3$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+u+1}+2^{s+1}+2^s-3}$ is an $\mathbb{F}_2$-vector space of dimension 154 with a basis consisting of all the classes represented by the monomials $a_{u,1,s,j}, 1 \leqslant j \leqslant 154$, which are determined as follows:*

For $s \geqslant 2$,

1. $(1, 2^s-2, 2^{s+1}-1, 2^{s+u+1}-1)$,
2. $(1, 2^s-2, 2^{s+u+1}-1, 2^{s+1}-1)$,
3. $(1, 2^{s+1}-1, 2^s-2, 2^{s+u+1}-1)$,
4. $(1, 2^{s+1}-1, 2^{s+u+1}-1, 2^s-2)$,
5. $(1, 2^{s+u+1}-1, 2^s-2, 2^{s+1}-1)$,
6. $(1, 2^{s+u+1}-1, 2^{s+1}-1, 2^s-2)$,
7. $(2^{s+1}-1, 1, 2^s-2, 2^{s+u+1}-1)$,
8. $(2^{s+1}-1, 1, 2^{s+u+1}-1, 2^s-2)$,
9. $(2^{s+1}-1, 2^{s+u+1}-1, 1, 2^s-2)$,
10. $(2^{s+u+1}-1, 1, 2^s-2, 2^{s+1}-1)$,
11. $(2^{s+u+1}-1, 1, 2^{s+1}-1, 2^s-2)$,
12. $(2^{s+u+1}-1, 2^{s+1}-1, 1, 2^s-2)$,
13. $(1, 2^s-1, 2^{s+1}-2, 2^{s+u+1}-1)$,
14. $(1, 2^s-1, 2^{s+u+1}-1, 2^{s+1}-2)$,
15. $(1, 2^{s+1}-2, 2^s-1, 2^{s+u+1}-1)$,
16. $(1, 2^{s+1}-2, 2^{s+u+1}-1, 2^s-1)$,
17. $(1, 2^{s+u+1}-1, 2^s-1, 2^{s+1}-2)$,
18. $(1, 2^{s+u+1}-1, 2^{s+1}-2, 2^s-1)$,
19. $(2^s-1, 1, 2^{s+1}-2, 2^{s+u+1}-1)$,
20. $(2^s-1, 1, 2^{s+u+1}-1, 2^{s+1}-2)$,
21. $(2^s-1, 2^{s+u+1}-1, 1, 2^{s+1}-2)$,
22. $(2^{s+u+1}-1, 1, 2^s-1, 2^{s+1}-2)$,
23. $(2^{s+u+1}-1, 1, 2^{s+1}-2, 2^s-1)$,
24. $(2^{s+u+1}-1, 2^s-1, 1, 2^{s+1}-2)$,
25. $(1, 2^s-1, 2^{s+1}-1, 2^{s+u+1}-2)$,
26. $(1, 2^s-1, 2^{s+u+1}-2, 2^{s+1}-1)$,
27. $(1, 2^{s+1}-1, 2^s-1, 2^{s+u+1}-2)$,
28. $(1, 2^{s+1}-1, 2^{s+u+1}-2, 2^s-1)$,
29. $(1, 2^{s+u+1}-2, 2^s-1, 2^{s+1}-1)$,
30. $(1, 2^{s+u+1}-2, 2^{s+1}-1, 2^s-1)$,
31. $(2^s-1, 1, 2^{s+1}-1, 2^{s+u+1}-2)$,
32. $(2^s-1, 1, 2^{s+u+1}-2, 2^{s+1}-1)$,
33. $(2^s-1, 2^{s+1}-1, 1, 2^{s+u+1}-2)$,
34. $(2^{s+1}-1, 1, 2^s-1, 2^{s+u+1}-2)$,

35. $(2^{s+1}-1, 1, 2^{s+u+1}-2, 2^s-1)$,
36. $(2^{s+1}-1, 2^s-1, 1, 2^{s+u+1}-2)$,
37. $(1, 2^s-2, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-1)$,
38. $(1, 2^{s+2}-1, 2^s-2, 2^{s+u+1}-2^{s+1}-1)$,
39. $(1, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-1, 2^s-2)$,
40. $(2^{s+2}-1, 1, 2^s-2, 2^{s+u+1}-2^{s+1}-1)$,
41. $(2^{s+2}-1, 1, 2^{s+u+1}-2^{s+1}-1, 2^s-2)$,
42. $(2^{s+2}-1, 2^{s+u+1}-2^{s+1}-1, 1, 2^s-2)$,
43. $(1, 2^s-1, 2^{s+2}-2, 2^{s+u+1}-2^{s+1}-1)$,
44. $(1, 2^{s+2}-2, 2^s-1, 2^{s+u+1}-2^{s+1}-1)$,
45. $(1, 2^{s+2}-2, 2^{s+u+1}-2^{s+1}-1, 2^s-1)$,
46. $(2^s-1, 1, 2^{s+2}-2, 2^{s+u+1}-2^{s+1}-1)$,
47. $(1, 2^s-1, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-2)$,
48. $(1, 2^{s+2}-1, 2^s-1, 2^{s+u+1}-2^{s+1}-2)$,
49. $(1, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-2, 2^s-1)$,
50. $(2^s-1, 1, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-2)$,
51. $(2^s-1, 2^{s+2}-1, 1, 2^{s+u+1}-2^{s+1}-2)$,
52. $(2^{s+2}-1, 1, 2^s-1, 2^{s+u+1}-2^{s+1}-2)$,
53. $(2^{s+2}-1, 1, 2^{s+u+1}-2^{s+1}-2, 2^s-1)$,
54. $(2^{s+2}-1, 2^s-1, 1, 2^{s+u+1}-2^{s+1}-2)$,
55. $(1, 2^{s+1}-2, 2^{s+1}-1, 2^{s+u+1}-2^s-1)$,
56. $(1, 2^{s+1}-2, 2^{s+u+1}-2^s-1, 2^{s+1}-1)$,
57. $(1, 2^{s+1}-1, 2^{s+1}-2, 2^{s+u+1}-2^s-1)$,
58. $(1, 2^{s+1}-1, 2^{s+u+1}-2^s-1, 2^{s+1}-2)$,
59. $(2^{s+1}-1, 1, 2^{s+1}-2, 2^{s+u+1}-2^s-1)$,
60. $(2^{s+1}-1, 1, 2^{s+u+1}-2^s-1, 2^{s+1}-2)$,
61. $(2^{s+1}-1, 2^{s+u+1}-2^s-1, 1, 2^{s+1}-2)$,
62. $(1, 2^{s+1}-1, 2^{s+1}-1, 2^{s+u+1}-2^s-2)$,
63. $(1, 2^{s+1}-1, 2^{s+u+1}-2^s-2, 2^{s+1}-1)$,
64. $(2^{s+1}-1, 1, 2^{s+1}-1, 2^{s+u+1}-2^s-2)$,
65. $(2^{s+1}-1, 1, 2^{s+u+1}-2^s-2, 2^{s+1}-1)$,
66. $(2^{s+1}-1, 2^{s+1}-1, 1, 2^{s+u+1}-2^s-2)$,
67. $(3, 2^{s+1}-3, 2^s-2, 2^{s+u+1}-1)$,
68. $(3, 2^{s+1}-3, 2^{s+u+1}-1, 2^s-2)$,
69. $(3, 2^{s+u+1}-1, 2^{s+1}-3, 2^s-2)$,
70. $(2^{s+u+1}-1, 3, 2^{s+1}-3, 2^s-2)$,
71. $(1, 2^{s+1}-2, 2^{s+1}+2^s-1, 2^{s+u+1}-2^{s+1}-1)$,
72. $(1, 2^{s+1}-1, 2^{s+1}+2^s-2, 2^{s+u+1}-2^{s+1}-1)$,
73. $(2^{s+1}-1, 1, 2^{s+1}+2^s-2, 2^{s+u+1}-2^{s+1}-1)$,
74. $(1, 2^{s+1}-1, 2^{s+1}+2^s-1, 2^{s+u+1}-2^{s+1}-2)$,
75. $(2^{s+1}-1, 1, 2^{s+1}+2^s-1, 2^{s+u+1}-2^{s+1}-2)$,
76. $(2^{s+1}-1, 2^{s+1}+2^s-1, 1, 2^{s+u+1}-2^{s+1}-2)$,
77. $(3, 2^{s+1}-1, 2^{s+u+1}-3, 2^s-2)$,
78. $(3, 2^{s+u+1}-3, 2^s-2, 2^{s+1}-1)$,
79. $(3, 2^{s+u+1}-3, 2^{s+1}-1, 2^s-2)$,

80. $(2^{s+1}-1, 3, 2^{s+u+1}-3, 2^s-2)$,
81. $(3, 2^s-1, 2^{s+1}-3, 2^{s+u+1}-2)$,
82. $(3, 2^{s+1}-3, 2^s-1, 2^{s+u+1}-2)$,
83. $(3, 2^{s+1}-3, 2^{s+u+1}-2, 2^s-1)$,
84. $(3, 2^s-1, 2^{s+u+1}-3, 2^{s+1}-2)$,
85. $(3, 2^{s+u+1}-3, 2^s-1, 2^{s+1}-2)$,
86. $(3, 2^{s+u+1}-3, 2^{s+1}-2, 2^s-1)$,
87. $(1, 2^{s+1}-2, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-2^s-1)$,
88. $(1, 2^{s+2}-1, 2^{s+1}-2, 2^{s+u+1}-2^{s+1}-2^s-1)$,
89. $(2^{s+2}-1, 1, 2^{s+1}-2, 2^{s+u+1}-2^{s+1}-2^s-1)$,
90. $(1, 2^{s+1}-1, 2^{s+2}-2, 2^{s+u+1}-2^{s+1}-2^s-1)$,
91. $(1, 2^{s+2}-2, 2^{s+1}-1, 2^{s+u+1}-2^{s+1}-2^s-1)$,
92. $(2^{s+1}-1, 1, 2^{s+2}-2, 2^{s+u+1}-2^{s+1}-2^s-1)$,
93. $(1, 2^{s+1}-1, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-2^s-2)$,
94. $(1, 2^{s+2}-1, 2^{s+1}-1, 2^{s+u+1}-2^{s+1}-2^s-2)$,
95. $(2^{s+1}-1, 1, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-2^s-2)$,
96. $(2^{s+1}-1, 2^{s+2}-1, 1, 2^{s+u+1}-2^{s+1}-2^s-2)$,
97. $(2^{s+2}-1, 1, 2^{s+1}-1, 2^{s+u+1}-2^{s+1}-2^s-2)$,
98. $(2^{s+2}-1, 2^{s+1}-1, 1, 2^{s+u+1}-2^{s+1}-2^s-2)$,
99. $(3, 2^{s+2}-3, 2^s-2, 2^{s+u+1}-2^{s+1}-1)$,
100. $(3, 2^{s+2}-3, 2^{s+u+1}-2^{s+1}-1, 2^s-2)$,
101. $(3, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-3, 2^s-2)$,
102. $(2^{s+2}-1, 3, 2^{s+u+1}-2^{s+1}-3, 2^s-2)$,
103. $(3, 2^s-1, 2^{s+2}-3, 2^{s+u+1}-2^{s+1}-2)$,
104. $(3, 2^{s+2}-3, 2^s-1, 2^{s+u+1}-2^{s+1}-2)$,
105. $(3, 2^{s+2}-3, 2^{s+u+1}-2^{s+1}-2, 2^s-1)$,
106. $(3, 2^{s+1}-3, 2^{s+1}-2, 2^{s+u+1}-2^s-1)$,
107. $(3, 2^{s+1}-3, 2^{s+u+1}-2^s-1, 2^{s+1}-2)$,
108. $(3, 2^{s+1}-3, 2^{s+1}-1, 2^{s+u+1}-2^s-2)$,
109. $(3, 2^{s+1}-3, 2^{s+u+1}-2^s-2, 2^{s+1}-1)$,
110. $(3, 2^{s+1}-1, 2^{s+1}-3, 2^{s+u+1}-2^s-2)$,
111. $(2^{s+1}-1, 3, 2^{s+1}-3, 2^{s+u+1}-2^s-2)$,
112. $(3, 2^{s+1}-1, 2^{s+u+1}-2^s-3, 2^{s+1}-2)$,
113. $(2^{s+1}-1, 3, 2^{s+u+1}-2^s-3, 2^{s+1}-2)$,
114. $(7, 2^{s+u+1}-5, 2^{s+1}-3, 2^s-2)$,
115. $(3, 2^{s+1}-3, 2^{s+1}+2^s-2, 2^{s+u+1}-2^{s+1}-1)$,
116. $(3, 2^{s+1}-3, 2^{s+1}+2^s-1, 2^{s+u+1}-2^{s+1}-2)$,
117. $(3, 2^{s+1}-1, 2^{s+1}+2^s-3, 2^{s+u+1}-2^{s+1}-2)$,
118. $(2^{s+1}-1, 3, 2^{s+1}+2^s-3, 2^{s+u+1}-2^{s+1}-2)$,
119. $(3, 2^{s+1}-3, 2^{s+2}-2, 2^{s+u+1}-2^{s+1}-2^s-1)$,
120. $(3, 2^{s+1}-3, 2^{s+2}-1, 2^{s+u+1}-2^{s+1}-2^s-2)$,
121. $(3, 2^{s+2}-1, 2^{s+1}-3, 2^{s+u+1}-2^{s+1}-2^s-2)$,
122. $(2^{s+2}-1, 3, 2^{s+1}-3, 2^{s+u+1}-2^{s+1}-2^s-2)$,
123. $(3, 2^{s+2}-3, 2^{s+1}-2, 2^{s+u+1}-2^{s+1}-2^s-1)$,
124. $(3, 2^{s+1}-1, 2^{s+2}-3, 2^{s+u+1}-2^{s+1}-2^s-2)$,

125. $(3, 2^{s+2} - 3, 2^{s+1} - 1, 2^{s+u+1} - 2^{s+1} - 2^s - 2)$,
126. $(2^{s+1} - 1, 3, 2^{s+2} - 3, 2^{s+u+1} - 2^{s+1} - 2^s - 2)$,
127. $(7, 2^{s+2} - 5, 2^{s+u+1} - 2^{s+1} - 3, 2^s - 2)$,
128. $(7, 2^{s+2} - 5, 2^{s+1} - 3, 2^{s+u+1} - 2^{s+1} - 2^s - 2)$,

For $s = 2$,

129. $(3, 3, 4, 2^{u+3} - 1)$,
130. $(3, 3, 2^{u+3} - 1, 4)$,
131. $(3, 2^{u+3} - 1, 3, 4)$,
132. $(2^{u+3} - 1, 3, 3, 4)$,
133. $(3, 3, 7, 2^{u+3} - 4)$,
134. $(3, 3, 2^{u+3} - 4, 7)$,
135. $(3, 7, 3, 2^{u+3} - 4)$,
136. $(7, 3, 3, 2^{u+3} - 4)$,
137. $(3, 7, 2^{u+3} - 5, 4)$,
138. $(7, 3, 2^{u+3} - 5, 4)$,
139. $(7, 2^{u+3} - 5, 3, 4)$,
140. $(7, 7, 2^{u+3} - 7, 2)$,
141. $(3, 3, 12, 2^{u+3} - 9)$,
142. $(3, 3, 15, 2^{u+3} - 12)$,
143. $(3, 15, 3, 2^{u+3} - 12)$,
144. $(15, 3, 3, 2^{u+3} - 12)$,
145. $(3, 7, 7, 2^{u+3} - 8)$,
146. $(7, 3, 7, 2^{u+3} - 8)$,
147. $(7, 7, 3, 2^{u+3} - 8)$,
148. $(7, 7, 2^{u+3} - 8, 3)$,
149. $(3, 7, 11, 2^{u+3} - 12)$,
150. $(7, 3, 11, 2^{u+3} - 12)$,
151. $(7, 11, 3, 2^{u+3} - 12)$,
152. $(7, 7, 8, 2^{u+3} - 13)$,
153. $(7, 7, 9, 2^{u+3} - 14)$,
154. $(7, 7, 11, 2^{u+3} - 16)$.

For $s \geqslant 3$

129. $(3, 2^s - 3, 2^{s+1} - 2, 2^{s+u+1} - 1)$,
130. $(3, 2^s - 3, 2^{s+u+1} - 1, 2^{s+1} - 2)$,
131. $(3, 2^{s+u+1} - 1, 2^s - 3, 2^{s+1} - 2)$,
132. $(2^{s+u+1} - 1, 3, 2^s - 3, 2^{s+1} - 2)$,
133. $(3, 2^s - 3, 2^{s+1} - 1, 2^{s+u+1} - 2)$,
134. $(3, 2^s - 3, 2^{s+u+1} - 2, 2^{s+1} - 1)$,
135. $(3, 2^{s+1} - 1, 2^s - 3, 2^{s+u+1} - 2)$,
136. $(2^{s+1} - 1, 3, 2^s - 3, 2^{s+u+1} - 2)$,
137. $(2^s - 1, 3, 2^{s+1} - 3, 2^{s+u+1} - 2)$,
138. $(2^s - 1, 3, 2^{s+u+1} - 3, 2^{s+1} - 2)$,
139. $(3, 2^s - 3, 2^{s+2} - 2, 2^{s+u+1} - 2^{s+1} - 1)$,
140. $(3, 2^s - 3, 2^{s+2} - 1, 2^{s+u+1} - 2^{s+1} - 2)$,
141. $(3, 2^{s+2} - 1, 2^s - 3, 2^{s+u+1} - 2^{s+1} - 2)$,
142. $(2^{s+2} - 1, 3, 2^s - 3, 2^{s+u+1} - 2^{s+1} - 2)$,
143. $(2^s - 1, 3, 2^{s+2} - 3, 2^{s+u+1} - 2^{s+1} - 2)$,
144. $(7, 2^{s+1} - 5, 2^s - 3, 2^{s+u+1} - 2)$,
145. $(7, 2^{s+1} - 5, 2^{s+u+1} - 3, 2^s - 2)$,
146. $(7, 2^{s+u+1} - 5, 2^s - 3, 2^{s+1} - 2)$,
147. $(7, 2^{s+2} - 5, 2^s - 3, 2^{s+u+1} - 2^{s+1} - 2)$,
148. $(7, 2^{s+1} - 5, 2^{s+1} - 3, 2^{s+u+1} - 2^s - 2)$,
149. $(7, 2^{s+1} - 5, 2^{s+u+1} - 2^s - 3, 2^{s+1} - 2)$,
150. $(7, 2^{s+1} - 5, 2^{s+1} + 2^s - 3, 2^{s+u+1} - 2^{s+1} - 2)$,
151. $(7, 2^{s+1} - 5, 2^{s+2} - 3, 2^{s+u+1} - 2^{s+1} - 2^s - 2)$.

For $s = 3$,

152. $(7, 7, 9, 2^{u+4} - 2)$, 153. $(7, 7, 2^{u+4} - 7, 14)$, 154. $(7, 7, 25, 2^{u+4} - 18)$.

For $s \geqslant 4$,

152. $(7, 2^s - 5, 2^{s+1} - 3, 2^{s+u+1} - 2)$,
153. $(7, 2^s - 5, 2^{s+u+1} - 3, 2^{s+1} - 2)$,
154. $(7, 2^s - 5, 2^{s+2} - 3, 2^{s+u+1} - 2^{s+1} - 2)$.

We prove the theorem by proving some propositions.

Combining Lemmas 3.5, 3.6, 5.10, 6.3.4, 6.3.5, 8.2.4, 8.4.4, 8.4.1. 8.5.4, 8.5.5, 8.6.4, 8.6.5, Theorem 2.5 and the results in Section 7, we obtain the following.

Proposition 8.7.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+u+1}+2^{s+1}+2^s-3}$ is generated by the elements listed in Theorem 8.7.2.*

Now, we prove that the elements listed in Theorem 8.7.2 are linearly independent.

Proposition 8.7.4. *For $u \geqslant 3$, the elements $[a_{u,1,2,j}], 1 \leqslant j \leqslant 154$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{u+3}+9}$.*

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{154} \gamma_j [a_{u,1,2,j}] = 0, \tag{8.7.4.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms f_1, f_2, f_3 to the relation (8.7.4.1), we have

$$\begin{aligned}
&\gamma_1 w_{u,1,2,1} + \gamma_2 w_{u,1,2,2} + \gamma_{15} w_{u,1,2,3} + \gamma_{16} w_{u,1,2,4} + \gamma_{29} w_{u,1,2,5} \\
&\quad + \gamma_{30} w_{u,1,2,6} + \gamma_{37} w_{u,1,2,7} + \gamma_{44} w_{u,1,2,8} + \gamma_{45} w_{u,1,2,9} + \gamma_{55} w_{u,1,2,10} \\
&\quad + \gamma_{56} w_{u,1,2,11} + \gamma_{71} w_{u,1,2,12} + \gamma_{87} w_{u,1,2,13} + \gamma_{91} w_{u,1,2,14} = 0, \\
&\gamma_3 w_{u,1,2,1} + \gamma_5 w_{u,1,2,2} + \gamma_{\{13,129\}} w_{u,1,2,3} + \gamma_{18} w_{u,1,2,4} \\
&\quad + \gamma_{\{26,134\}} w_{u,1,2,5} + \gamma_{\{28,148\}} w_{u,1,2,6} + \gamma_{38} w_{u,1,2,7} \\
&\quad + \gamma_{\{43,141\}} w_{u,1,2,8} + \gamma_{49} w_{u,1,2,9} + \gamma_{\{57,63,72\}} w_{u,1,2,10} \\
&\quad + \gamma_{63} w_{u,1,2,11} + \gamma_{72} w_{u,1,2,12} + \gamma_{88} w_{u,1,2,13} + \gamma_{\{90,152\}} w_{u,1,2,14} = 0, \\
&\gamma_4 w_{u,1,2,1} + \gamma_6 w_{u,1,2,2} + \gamma_{\{14,130\}} w_{u,1,2,3} + \gamma_{\{17,131\}} w_{u,1,2,4} \\
&\quad + \gamma_{\{25,133,146\}} w_{u,1,2,5} + \gamma_{\{27,135,147\}} w_{u,1,2,6} + \gamma_{39} w_{u,1,2,7} + \gamma_{\{47,142\}} w_{u,1,2,8} \\
&\quad + \gamma_{\{48,143\}} w_{u,1,2,9} + \gamma_{\{58,62,93,137,145\}} w_{u,1,2,10} + \gamma_{\{62,94,145\}} w_{u,1,2,11} \\
&\quad + \gamma_{93} w_{u,1,2,12} + \gamma_{94} w_{u,1,2,13} + \gamma_{\{74,149,154\}} w_{u,1,2,14} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$(8.7.4.2)\qquad \begin{cases} \gamma_j = 0,\ j = 1,2,3,4,5,6,15,16,18,29,30,37,38,\\ 39,44,45,49,55,56,57,63,71,72,87,88,91,93,94,\\ \gamma_{\{13,129\}} = \gamma_{\{26,134\}} = \gamma_{\{28,148\}} = \gamma_{\{43,141\}} = 0,\\ \gamma_{\{90,152\}} = \gamma_{\{14,130\}} = \gamma_{\{17,131\}} = \gamma_{\{47,142\}} = 0,\\ \gamma_{\{48,143\}} = \gamma_{\{25,133,146\}} = \gamma_{\{27,135,147\}} = 0,\\ \gamma_{\{62,145\}} = \gamma_{\{58,62,137,145\}} = \gamma_{\{74,149,154\}} = 0. \end{cases}$$

With the aid of (8.7.4.2), the homomorphisms f_4, f_5, f_6 send (8.7.4.1) to

$$\begin{aligned} &\gamma_{\{19,67,129\}}w_{u,1,2,1} + \gamma_{\{32,78,109,134\}}w_{u,1,2,2} + \gamma_7 w_{u,1,2,3}\\ &\quad + \gamma_{\{35,148\}}w_{u,1,2,4} + \gamma_{10}w_{u,1,2,5} + \gamma_{23}w_{u,1,2,6} + \gamma_{40}w_{u,1,2,8}\\ &\quad + \gamma_{\{46,99,115,141\}}w_{u,1,2,7} + \gamma_{53}w_{u,1,2,9} + \gamma_{59}w_{u,1,2,10}\\ &\quad + \gamma_{65}w_{u,1,2,11} + \gamma_{73}w_{u,1,2,12} + \gamma_{\{92,152\}}w_{u,1,2,13} + \gamma_{89}w_{u,1,2,14} = 0,\\ &\gamma_{\{20,68,130\}}w_{u,1,2,1} + a_1 w_{u,1,2,2} + \gamma_8 w_{u,1,2,3} + \gamma_{11}w_{u,1,2,5}\\ &\quad + \gamma_{\{34,136,139,147,151\}}w_{u,1,2,4} + \gamma_{\{22,132\}}w_{u,1,2,6} + \gamma_{\{50,100,120,142\}}w_{u,1,2,7}\\ &\quad + \gamma_{41}w_{u,1,2,8} + \gamma_{\{52,144\}}w_{u,1,2,9} + \gamma_{\{60,97,138\}}w_{u,1,2,10} + \gamma_{\{64,97,146\}}w_{u,1,2,11}\\ &\quad + \gamma_{95}w_{u,1,2,12} + \gamma_{\{75,150,154\}}w_{u,1,2,13} + \gamma_{97}w_{u,1,2,14} = 0,\\ &a_2 w_{u,1,2,1} + \gamma_{\{21,69,131\}}w_{u,1,2,2} + a_3 w_{u,1,2,3} + \gamma_9 w_{u,1,2,4}\\ &\quad + \gamma_{\{24,70,132\}}w_{u,1,2,5} + \gamma_{\{51,101,121,143\}}w_{u,1,2,7} + \gamma_{12}w_{u,1,2,6}\\ &\quad + \gamma_{\{54,102,122,144\}}w_{u,1,2,8} + \gamma_{42}w_{u,1,2,9} + a_4 w_{u,1,2,10} + \gamma_{\{61,114,139\}}w_{u,1,2,11}\\ &\quad + \gamma_{96}w_{u,1,2,13} + \gamma_{\{76,127,128,151\}}w_{u,1,2,12} + \gamma_{98}w_{u,1,2,14} = 0, \end{aligned}$$

where

$$\begin{aligned} &a_1 = \gamma_{\{31,79,108,125,133,145\}},\quad a_2 = \gamma_{\{33,77,110,112,117,124,135,137,145,149\}},\\ &a_3 = \gamma_{\{36,80,111,113,118,126,136,138,146,150\}},\quad a_4 = \gamma_{\{66,140,147,148,152,153,154\}}. \end{aligned}$$

Computing from these equalities, we obtain

$$(8.7.4.3)\qquad \begin{cases} a_i = 0,\ i = 1,2,3,4,\\ \gamma_j = 0,\ j = 7,\ldots,12,23,40,41,42,\\ \qquad\qquad 53,59,65,73,89,95,96,97,98,\\ \gamma_{\{32,78,109,134\}} = \gamma_{\{35,148\}} = \gamma_{\{92,152\}} = \gamma_{\{46,99,115,141\}} = 0,\\ \gamma_{\{22,132\}} = \gamma_{\{52,144\}} = \gamma_{\{20,68,130\}} = \gamma_{\{34,136,139,147,151\}} = 0,\\ \gamma_{\{50,100,120,142\}} = \gamma_{\{76,127,128,151\}} = \gamma_{\{60,138\}} = \gamma_{\{64,146\}} = 0\\ \gamma_{\{75,150,154\}} = \gamma_{\{21,69,131\}} = \gamma_{\{24,70,132\}} = \gamma_{\{61,114,139\}} = 0,\\ \gamma_{\{51,101,121,143\}} = \gamma_{\{54,102,122,144\}} = 0. \end{cases}$$

With the aid of (8.7.4.2) and (8.7.4.3), the homomorphisms g_1, g_2 send (8.7.4.1) to

$$\begin{aligned}
&\gamma_{19}w_{u,1,2,1} + \gamma_{32}w_{u,1,2,2} + \gamma_{67}w_{u,1,2,3} + \gamma_{\{28,83\}}w_{u,1,2,4} + \gamma_{78}w_{u,1,2,5}\\
&\quad + \gamma_{86}w_{u,1,2,6} + \gamma_{46}w_{u,1,2,7} + \gamma_{99}w_{u,1,2,8} + \gamma_{105}w_{u,1,2,9} + \gamma_{106}w_{u,1,2,10}\\
&\quad + \gamma_{109}w_{u,1,2,11} + \gamma_{115}w_{u,1,2,12} + \gamma_{\{90,119\}}w_{u,1,2,13} + \gamma_{123}w_{u,1,2,14} = 0,\\
&\gamma_{20}w_{u,1,2,1} + \gamma_{\{25,31,133\}}w_{u,1,2,2} + \gamma_{68}w_{u,1,2,3} + \gamma_{\{27,34,82,135,136,147\}}w_{u,1,2,4}\\
&\quad + \gamma_{79}w_{u,1,2,5} + \gamma_{\{85,139\}}w_{u,1,2,6} + \gamma_{50}w_{u,1,2,7} + \gamma_{100}w_{u,1,2,8}\\
&\quad + \gamma_{\{104,151\}}w_{u,1,2,9} + \gamma_{\{107,125\}}w_{u,1,2,10} + \gamma_{\{108,125\}}w_{u,1,2,11}\\
&\quad + \gamma_{120}w_{u,1,2,12} + \gamma_{\{74,75,116,149,150,154\}}w_{u,1,2,13} + \gamma_{125}w_{u,1,2,14} = 0.
\end{aligned}$$

These equalities imply

$$\text{(8.7.4.4)}\quad \begin{cases}
\gamma_j = 0,\ j = 19, 20, 32, 46, 50, 67, 68, 78, 79, 86,\\
99, 100, 105, 106, 107, 108, 109, 115, 120, 123, 125,\\
\gamma_{\{28,83\}} = \gamma_{\{90,119\}} = \gamma_{\{25,31,133\}} = \gamma_{\{85,139\}} = 0,\\
\gamma_{\{27,34,82,135,136,147\}} = \gamma_{\{104,151\}} = \gamma_{\{74,75,116,149,150,154\}} = 0.
\end{cases}$$

With the aid of (8.7.4.2), (8.7.4.3) and (8.7.4.4), the homomorphisms g_3, g_4 send (8.7.4.1) to

$$\begin{aligned}
&\gamma_{21}w_{u,1,2,1} + \gamma_{\{27,33,66,135,147\}}w_{u,1,2,2} + \gamma_{69}w_{u,1,2,3} + a_5w_{u,1,2,4} + a_6w_{u,1,2,6}\\
&\quad + \gamma_{\{77,140\}}w_{u,1,2,5} + \gamma_{51}w_{u,1,2,7} + \gamma_{101}w_{u,1,2,8} + a_7w_{u,1,2,9} + a_8w_{u,1,2,10}\\
&\quad + a_9w_{u,1,2,11} + \gamma_{121}w_{u,1,2,12} + a_{10}w_{u,1,2,13} + \gamma_{\{124,127,153,154\}}w_{u,1,2,14} = 0,\\
&\gamma_{24}w_{u,1,2,1} + a_{11}w_{u,1,2,2} + \gamma_{70}w_{u,1,2,3} + a_{12}w_{u,1,2,4} + \gamma_{\{80,114,127,140\}}w_{u,1,2,5}\\
&\quad + a_{13}w_{u,1,2,6} + \gamma_{54}w_{u,1,2,7} + \gamma_{102}w_{u,1,2,8} + a_{14}w_{u,1,2,9} + a_{15}w_{u,1,2,10}\\
&\quad + a_{16}w_{u,1,2,11} + \gamma_{122}w_{u,1,2,12} + a_{17}w_{u,1,2,13} + \gamma_{\{126,127,128,153,154\}}w_{u,1,2,14} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_5 = \gamma_{\{22,24,25,36,52,54,64,70,81,111,122,133,136\}},\quad a_6 = \gamma_{\{22,24,60,70,80,84,102,113\}},\\
&a_7 = \gamma_{\{52,54,102,103,118,122,126,150\}},\quad a_8 = \gamma_{\{28,61,112,114,124,139,140,153,154\}},\\
&a_9 = \gamma_{\{28,66,74,90,110,112,117,124,140,147,149\}}, a_{10} = \gamma_{\{74,76,90,117,128,149,151,153,154\}},\\
&a_{12} = \gamma_{\{13,17,21,31,33,48,51,62,69,81,82,85,104,110,121,133,135\}},\\
&a_{11} = \gamma_{\{34,36,61,66,76,136,139,147,151\}},\quad a_{13} = \gamma_{\{14,17,21,26,58,69,77,83,84,85,101,112\}},\\
&a_{14} = \gamma_{\{43,47,48,51,101,103,104,116,117,119,121,124,149\}},\\
&a_{15} = \gamma_{\{28,61,113,126,127,128,139,140,153,154\}},\quad a_{17} = \gamma_{\{75,76,90,118,150,151,153,154\}},\\
&a_{16} = \gamma_{\{28,61,66,75,76,90,111,113,114,118,126,127,139,140,147,150,151\}}.
\end{aligned}$$

From the above equalities, we obtain

$$(8.7.4.5)\qquad \begin{cases} a_i = 0,\ i = 5, 6, \ldots, 17, \\ \gamma_j = 0,\ j = 21, 24, 51, 54, 69, 70, 101, 102, 121, 122, \\ \gamma_{\{27,33,66,135,147\}} = \gamma_{\{77,140\}} = \gamma_{\{124,127,153,154\}} = 0, \\ \gamma_{\{80,114,127,140\}} = \gamma_{\{126,127,128,153,154\}} = 0. \end{cases}$$

With the aid of (8.7.4.2), (8.7.4.3), (8.7.4.4) and (8.7.4.5), the homomorphism h sends (8.7.4.1) to

$$\begin{aligned} &a_{18}w_{u,1,2,1} + \gamma_{\{28,80,84,113\}}w_{u,1,2,2} + a_{19}w_{u,1,2,3} + \gamma_{80}w_{u,1,2,4} \\ &\quad + \gamma_{61}w_{u,1,2,5} + \gamma_{114}w_{u,1,2,6} + \gamma_{\{75,90,103,118,126,150\}}w_{u,1,2,7} \\ &\quad + \gamma_{76}w_{u,1,2,8} + \gamma_{127}w_{u,1,2,9} + a_{20}w_{u,1,2,10} + a_{21}w_{u,1,2,11} \\ &\quad + a_{22}w_{u,1,2,12} + \gamma_{\{124,126,127,153,154\}}w_{u,1,2,13} + \gamma_{128}w_{u,1,2,14} = 0, \end{aligned}$$

where

$$\begin{aligned} &a_{18} = \gamma_{\{25,31,34,36,81,111,133,136\}},\quad a_{19} = \gamma_{\{27,33,34,36,66,82,135,136,147\}}, \\ &a_{20} = \gamma_{\{61,66,76,85,104,110,111,114,128,147\}},\quad a_{21} = \gamma_{\{28,61,77,85,112,113,114,127\}}, \\ &a_{22} = \gamma_{\{74,75,76,90,104,116,117,118,128,149,150,153,154\}}. \end{aligned}$$

From the above equalities, it implies

$$(8.7.4.6)\qquad \begin{cases} a_i = 0,\ i = 18, 19, \ldots, 22, \\ \gamma_j = 0,\ j = 61, 76, 80, 114, 127, 128, \\ \gamma_{\{75,90,103,118,126,150\}} = \gamma_{\{28,84,113\}} = \gamma_{\{124,126,153,154\}} = 0. \end{cases}$$

Combining (8.7.4.2), (8.7.4.3), (8.7.4.4), (8.7.4.5) and (8.7.4.6), we get $\gamma_j = 0$ for $1 \leqslant j \leqslant 154$. The proposition is proved. □

Proposition 8.7.5. *For $s \geqslant 3$ and $u \geqslant 3$, the elements $[a_{u,1,s,j}], 1 \leqslant j \leqslant 154$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+u+1}+2^{s+1}+2^s-3}$.*

Proof. Suppose that there is a linear relation

$$(8.7.5.1)\qquad \sum_{j=1}^{154} \gamma_j [a_{u,1,s,j}] = 0,$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.7.5.1), we obtain

$$\begin{aligned} &\gamma_1 w_{u,1,s,1} + \gamma_2 w_{u,1,s,2} + \gamma_{15} w_{u,1,s,3} + \gamma_{16} w_{u,1,s,4} + \gamma_{29} w_{u,1,s,5} \\ &\quad + \gamma_{30} w_{u,1,s,6} + \gamma_{37} w_{u,1,s,7} + \gamma_{44} w_{u,1,s,8} + \gamma_{45} w_{u,1,s,9} + \gamma_{55} w_{u,1,s,10} \\ &\quad + \gamma_{56} w_{u,1,s,11} + \gamma_{71} w_{u,1,s,12} + \gamma_{87} w_{u,1,s,13} + \gamma_{91} w_{u,1,s,14} = 0, \end{aligned}$$

$$\begin{aligned}
&\gamma_3 w_{u,1,s,1} + \gamma_5 w_{u,1,s,2} + \gamma_{13} w_{u,1,s,3} + \gamma_{18} w_{u,1,s,4} + \gamma_{26} w_{u,1,s,5} + \gamma_{28} w_{u,1,s,6}\\
&\quad + \gamma_{38} w_{u,1,s,7} + \gamma_{43} w_{u,1,s,8} + \gamma_{49} w_{u,1,s,9} + \gamma_{\{57,63,72\}} w_{u,1,s,10}\\
&\quad + \gamma_{63} w_{u,1,s,11} + \gamma_{72} w_{u,1,s,12} + \gamma_{88} w_{u,1,s,13} + \gamma_{90} w_{u,1,s,14} = 0,\\
&\gamma_4 w_{u,1,s,1} + \gamma_6 w_{u,1,s,2} + \gamma_{14} w_{u,1,s,3} + \gamma_{17} w_{u,1,s,4} + \gamma_{25} w_{u,1,s,5} + \gamma_{27} w_{u,1,s,6}\\
&\quad + \gamma_{39} w_{u,1,s,7} + \gamma_{47} w_{u,1,s,8} + \gamma_{48} w_{u,1,s,9} + \gamma_{\{58,62,93\}} w_{u,1,s,10}\\
&\quad + \gamma_{\{62,94\}} w_{u,1,s,11} + \gamma_{93} w_{u,1,s,12} + \gamma_{94} w_{u,1,s,13} + \gamma_{74} w_{u,1,s,14} = 0,\\
&\gamma_{19} w_{u,1,s,1} + \gamma_{32} w_{u,1,s,2} + \gamma_7 w_{u,1,s,3} + \gamma_{35} w_{u,1,s,4} + \gamma_{10} w_{u,1,s,5}\\
&\quad + \gamma_{23} w_{u,1,s,6} + \gamma_{46} w_{u,1,s,7} + \gamma_{40} w_{u,1,s,8} + \gamma_{53} w_{u,1,s,9} + \gamma_{59} w_{u,1,s,10}\\
&\quad + \gamma_{65} w_{u,1,s,11} + \gamma_{73} w_{u,1,s,12} + \gamma_{92} w_{u,1,s,13} + \gamma_{89} w_{u,1,s,14} = 0,\\
&\gamma_{20} w_{u,1,s,1} + \gamma_{31} w_{u,1,s,2} + \gamma_8 w_{u,1,s,3} + \gamma_{34} w_{u,1,s,4} + \gamma_{11} w_{u,1,s,5}\\
&\quad + \gamma_{22} w_{u,1,s,6} + \gamma_{50} w_{u,1,s,7} + \gamma_{41} w_{u,1,s,8} + \gamma_{52} w_{u,1,s,9} + \gamma_{\{60,97\}} w_{u,1,s,10}\\
&\quad + \gamma_{\{64,97\}} w_{u,1,s,11} + \gamma_{95} w_{u,1,s,12} + \gamma_{75} w_{u,1,s,13} + \gamma_{97} w_{u,1,s,14} = 0,\\
&\gamma_{33} w_{u,1,s,1} + \gamma_{21} w_{u,1,s,2} + \gamma_{36} w_{u,1,s,3} + \gamma_9 w_{u,1,s,4} + \gamma_{24} w_{u,1,s,5}\\
&\quad + \gamma_{12} w_{u,1,s,6} + \gamma_{51} w_{u,1,s,7} + \gamma_{54} w_{u,1,s,8} + \gamma_{42} w_{u,1,s,9} + \gamma_{66} w_{u,1,s,10}\\
&\quad + \gamma_{61} w_{u,1,s,11} + \gamma_{76} w_{u,1,s,12} + \gamma_{96} w_{u,1,s,13} + \gamma_{98} w_{u,1,s,14} = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\gamma_j = 0, j = 1, \dots, 66, 71, \dots, 76, 87, \dots, 98. \tag{8.7.5.2}$$

With the aid of (8.7.5.2), the homomorphisms g_1, g_2, g_3, g_4 send (8.7.5.1) to

$$\begin{aligned}
&\gamma_{129} w_{u,1,s,1} + \gamma_{134} w_{u,1,s,2} + \gamma_{67} w_{u,1,s,3} + \gamma_{83} w_{u,1,s,4} + \gamma_{78} w_{u,1,s,5}\\
&\quad + \gamma_{86} w_{u,1,s,6} + \gamma_{139} w_{u,1,s,7} + \gamma_{99} w_{u,1,s,8} + \gamma_{105} w_{u,1,s,9} + \gamma_{106} w_{u,1,s,10}\\
&\quad + \gamma_{109} w_{u,1,s,11} + \gamma_{115} w_{u,1,s,12} + \gamma_{119} w_{u,1,s,13} + \gamma_{123} w_{u,1,s,14} = 0,\\
&\gamma_{130} w_{u,1,s,1} + \gamma_{133} w_{u,1,s,2} + \gamma_{68} w_{u,1,s,3} + \gamma_{82} w_{u,1,s,4} + \gamma_{79} w_{u,1,s,5} + \gamma_{85} w_{u,1,s,6}\\
&\quad + \gamma_{140} w_{u,1,s,7} + \gamma_{100} w_{u,1,s,8} + \gamma_{104} w_{u,1,s,9} + \gamma_{\{107,125\}} w_{u,1,s,10}\\
&\quad + \gamma_{\{108,125\}} w_{u,1,s,11} + \gamma_{120} w_{u,1,s,12} + \gamma_{116} w_{u,1,s,13} + \gamma_{125} w_{u,1,s,14} = 0,\\
&\gamma_{131} w_{u,1,s,1} + \gamma_{135} w_{u,1,s,2} + \gamma_{69} w_{u,1,s,3} + a_1 w_{u,1,s,4,77} w_{u,1,s,5} + a_2 w_{u,1,s,6}\\
&\quad + \gamma_{141} w_{u,1,s,7} + \gamma_{101} w_{u,1,s,8} + a_3 w_{u,1,s,9} + \gamma_{\{110,112,117,124\}} w_{u,1,s,11}\\
&\quad + \gamma_{\{112,124\}} w_{u,1,s,10} + \gamma_{121} w_{u,1,s,12} + \gamma_{117} w_{u,1,s,13} + \gamma_{124} w_{u,1,s,14} = 0,\\
&\gamma_{132} w_{u,1,s,1} + \gamma_{136} w_{u,1,s,2} + \gamma_{70} w_{u,1,s,3} + a_4 w_{u,1,s,4} + \gamma_{80} w_{u,1,s,5} + a_5 w_{u,1,s,6}\\
&\quad + \gamma_{142} w_{u,1,s,7} + \gamma_{102} w_{u,1,s,8} + a_6 w_{u,1,s,9} + \gamma_{\{111,113,118,126\}} w_{u,1,s,11}\\
&\quad + \gamma_{\{113,126\}} w_{u,1,s,10} + \gamma_{122} w_{u,1,s,12} + \gamma_{118} w_{u,1,s,13} + \gamma_{126} w_{u,1,s,14} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases} \gamma_{\{81,152\}}, & s = 3,\\ \gamma_{81}, & s \geqslant 4, \end{cases} \quad a_4 = \begin{cases} \gamma_{\{114,128,137,144,146,147,148,152\}}, & s = 3,\\ \gamma_{137}, & s \geqslant 4, \end{cases}$$

$$a_2 = \begin{cases} \gamma_{\{84,153\}}, & s = 3, \\ \gamma_{84}, & s \geqslant 4, \end{cases} \quad a_5 = \begin{cases} \gamma_{\{114,127,138,145,146,149,153\}}, & s = 3, \\ \gamma_{138}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{103,154\}}, & s = 3, \\ \gamma_{103}, & s \geqslant 4, \end{cases} \quad a_6 = \begin{cases} \gamma_{\{127,128,143,147,150,151,154\}}, & s = 3, \\ \gamma_{143}, & s \geqslant 4. \end{cases}$$

These equalities imply

$$\begin{cases} a_i = 0, i = 1, 2, \ldots, 6, \\ \gamma_j = 0,\ j = 67, \ldots, 70, 77, 78, 79, 80, \\ 82, \ldots, 86, 100, 101, 102, 104, \ldots, 113, \\ 115, \ldots, 126, 129, \ldots, 136, 139, \ldots, 142. \end{cases} \tag{8.7.5.3}$$

With the aid of (8.7.5.2) and (8.7.5.3), the homomorphism h sends (8.7.5.1) to

$$\begin{aligned} &a_7 w_{u,1,s,1} + a_8 w_{u,1,s,2} + \gamma_{144} w_{u,1,s,3} + \gamma_{145} w_{u,1,s,4} + \gamma_{146} w_{u,1,s,5} \\ &\quad + \gamma_{114} w_{u,1,s,6} + a_9 w_{u,1,s,7} + \gamma_{147} w_{u,1,s,8} + \gamma_{127} w_{u,1,s,9} + \gamma_{148} w_{u,1,s,10} \\ &\quad + \gamma_{149} w_{u,1,s,11} + \gamma_{150} w_{u,1,s,12} + \gamma_{151} w_{u,1,s,13} + \gamma_{128} w_{u,1,s,14} = 0, \end{aligned}$$

where

$$a_7 = \begin{cases} \gamma_{137}, & s = 3, \\ \gamma_{152}, & s \geqslant 4, \end{cases} \quad a_8 = \begin{cases} \gamma_{138}, & s = 3, \\ \gamma_{153}, & s \geqslant 4, \end{cases} \quad a_9 = \begin{cases} \gamma_{143}, & s = 3, \\ \gamma_{154}, & s \geqslant 4. \end{cases}$$

From the above equalities, it implies

$$a_7 = a_8 = a_9 = 0, \gamma_j = 0,\ j = 114, 127, 128, 144, \ldots, 151. \tag{8.7.5.4}$$

Combining (8.7.5.2), (8.7.5.3) and (8.7.5.4), we get $\gamma_j = 0$ for $1 \leqslant j \leqslant 154$. The proposition is proved. □

8.8. **The case $s \geqslant 2, t \geqslant 2$ and $u = 1$.**

According to Kameko [8], for $s \geqslant 2, t \geqslant 2$, the dimension of the space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+t+1}+2^{s+t}+2^s-3}$ is 14 with a basis given by the classes $w_{1,t,s,j}, 1 \leqslant j \leqslant 14$, which are determined as follows:

1. $[2^s - 1, 2^{s+t} - 1, 2^{s+t+1} - 1]$,
2. $[2^s - 1, 2^{s+t+1} - 1, 2^{s+t} - 1]$,
3. $[2^{s+t} - 1, 2^s - 1, 2^{s+t+1} - 1]$,
4. $[2^{s+t} - 1, 2^{s+t+1} - 1, 2^s - 1]$,
5. $[2^{s+t+1} - 1, 2^s - 1, 2^{s+t} - 1]$,
6. $[2^{s+t+1} - 1, 2^{s+t} - 1, 2^s - 1]$,
7. $[2^{s+1} - 1, 2^{s+t} - 2^s - 1, 2^{s+t+1} - 1]$,
8. $[2^{s+1} - 1, 2^{s+t+1} - 1, 2^{s+t} - 2^s - 1]$,
9. $[2^{s+t+1} - 1, 2^{s+1} - 1, 2^{s+t} - 2^s - 1]$,
10. $[2^{s+1} - 1, 2^{s+t} - 1, 2^{s+t+1} - 2^s - 1]$,
11. $[2^{s+1} - 1, 2^{s+t+1} - 2^s - 1, 2^{s+t} - 1]$,
12. $[2^{s+t} - 1, 2^{s+1} - 1, 2^{s+t+1} - 2^s - 1]$,

13. $[2^{s+2}-1, 2^{s+t+1}-2^{s+1}-1, 2^{s+t}-2^s-1]$,
14. $[2^{s+2}-1, 2^{s+2}-1, 2^{s+2}+2^s-1]$, for $t=2$,
14. $[2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 2^{s+t+1}-2^s-1]$, for $t>2$.

So, we easily obtain

Proposition 8.8.1. *For any positive integer $s, t \geqslant 2$, we have*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+t+1}+2^{s+t}+2^s-3} = 56.$$

Now, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+t}+2^s-3}$. We have

Theorem 8.8.2. *$(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+t}+2^s-3}$ is an $\mathbb{F}_2$-vector space of dimension 154 with a basis consisting of all the classes represented by the monomials $a_{1,t,s,j}, 1 \leqslant j \leqslant 154$, which are determined as follows:*

For $s \geqslant 2$ and $t \geqslant 2$,

1. $(1, 2^s-2, 2^{s+t}-1, 2^{s+t+1}-1)$,
2. $(1, 2^s-2, 2^{s+t+1}-1, 2^{s+t}-1)$,
3. $(1, 2^{s+t}-1, 2^s-2, 2^{s+t+1}-1)$,
4. $(1, 2^{s+t}-1, 2^{s+t+1}-1, 2^s-2)$,
5. $(1, 2^{s+t+1}-1, 2^s-2, 2^{s+t}-1)$,
6. $(1, 2^{s+t+1}-1, 2^{s+t}-1, 2^s-2)$,
7. $(2^{s+t}-1, 1, 2^s-2, 2^{s+t+1}-1)$,
8. $(2^{s+t}-1, 1, 2^{s+t+1}-1, 2^s-2)$,
9. $(2^{s+t}-1, 2^{s+t+1}-1, 1, 2^s-2)$,
10. $(2^{s+t+1}-1, 1, 2^s-2, 2^{s+t}-1)$,
11. $(2^{s+t+1}-1, 1, 2^{s+t}-1, 2^s-2)$,
12. $(2^{s+t+1}-1, 2^{s+t}-1, 1, 2^s-2)$,
13. $(1, 2^s-1, 2^{s+t}-2, 2^{s+t+1}-1)$,
14. $(1, 2^s-1, 2^{s+t+1}-1, 2^{s+t}-2)$,
15. $(1, 2^{s+t}-2, 2^s-1, 2^{s+t+1}-1)$,
16. $(1, 2^{s+t}-2, 2^{s+t+1}-1, 2^s-1)$,
17. $(1, 2^{s+t+1}-1, 2^s-1, 2^{s+t}-2)$,
18. $(1, 2^{s+t+1}-1, 2^{s+t}-2, 2^s-1)$,
19. $(2^s-1, 1, 2^{s+t}-2, 2^{s+t+1}-1)$,
20. $(2^s-1, 1, 2^{s+t+1}-1, 2^{s+t}-2)$,
21. $(2^s-1, 2^{s+t+1}-1, 1, 2^{s+t}-2)$,
22. $(2^{s+t+1}-1, 1, 2^s-1, 2^{s+t}-2)$,
23. $(2^{s+t+1}-1, 1, 2^{s+t}-2, 2^s-1)$,
24. $(2^{s+t+1}-1, 2^s-1, 1, 2^{s+t}-2)$,
25. $(1, 2^s-1, 2^{s+t}-1, 2^{s+t+1}-2)$,
26. $(1, 2^s-1, 2^{s+t+1}-2, 2^{s+t}-1)$,
27. $(1, 2^{s+t}-1, 2^s-1, 2^{s+t+1}-2)$,
28. $(1, 2^{s+t}-1, 2^{s+t+1}-2, 2^s-1)$,
29. $(1, 2^{s+t+1}-2, 2^s-1, 2^{s+t}-1)$,
30. $(1, 2^{s+t+1}-2, 2^{s+t}-1, 2^s-1)$,

31. $(2^s-1, 1, 2^{s+t}-1, 2^{s+t+1}-2)$,
32. $(2^s-1, 1, 2^{s+t+1}-2, 2^{s+t}-1)$,
33. $(2^s-1, 2^{s+t}-1, 1, 2^{s+t+1}-2)$,
34. $(2^{s+t}-1, 1, 2^s-1, 2^{s+t+1}-2)$,
35. $(2^{s+t}-1, 1, 2^{s+t+1}-2, 2^s-1)$,
36. $(2^{s+t}-1, 2^s-1, 1, 2^{s+t+1}-2)$,
37. $(1, 2^{s+1}-2, 2^{s+t}-2^s-1, 2^{s+t+1}-1)$,
38. $(1, 2^{s+1}-2, 2^{s+t+1}-1, 2^{s+t}-2^s-1)$,
39. $(1, 2^{s+t+1}-1, 2^{s+1}-2, 2^{s+t}-2^s-1)$,
40. $(2^{s+t+1}-1, 1, 2^{s+1}-2, 2^{s+t}-2^s-1)$,
41. $(1, 2^{s+1}-1, 2^{s+t}-2^s-2, 2^{s+t+1}-1)$,
42. $(1, 2^{s+1}-1, 2^{s+t+1}-1, 2^{s+t}-2^s-2)$,
43. $(1, 2^{s+t+1}-1, 2^{s+1}-1, 2^{s+t}-2^s-2)$,
44. $(2^{s+1}-1, 1, 2^{s+t}-2^s-2, 2^{s+t+1}-1)$,
45. $(2^{s+1}-1, 1, 2^{s+t+1}-1, 2^{s+t}-2^s-2)$,
46. $(2^{s+1}-1, 2^{s+t+1}-1, 1, 2^{s+t}-2^s-2)$,
47. $(2^{s+t+1}-1, 1, 2^{s+1}-1, 2^{s+t}-2^s-2)$,
48. $(2^{s+t+1}-1, 2^{s+1}-1, 1, 2^{s+t}-2^s-2)$,
49. $(1, 2^{s+1}-1, 2^{s+t}-2^s-1, 2^{s+t+1}-2)$,
50. $(1, 2^{s+1}-1, 2^{s+t+1}-2, 2^{s+t}-2^s-1)$,
51. $(1, 2^{s+t+1}-2, 2^{s+1}-1, 2^{s+t}-2^s-1)$,
52. $(2^{s+1}-1, 1, 2^{s+t}-2^s-1, 2^{s+t+1}-2)$,
53. $(2^{s+1}-1, 1, 2^{s+t+1}-2, 2^{s+t}-2^s-1)$,
54. $(2^{s+1}-1, 2^{s+t}-2^s-1, 1, 2^{s+t+1}-2)$,
55. $(1, 2^{s+1}-2, 2^{s+t}-1, 2^{s+t+1}-2^s-1)$,
56. $(1, 2^{s+1}-2, 2^{s+t+1}-2^s-1, 2^{s+t}-1)$,
57. $(1, 2^{s+t}-1, 2^{s+1}-2, 2^{s+t+1}-2^s-1)$,
58. $(2^{s+t}-1, 1, 2^{s+1}-2, 2^{s+t+1}-2^s-1)$,
59. $(1, 2^{s+1}-1, 2^{s+t}-2, 2^{s+t+1}-2^s-1)$,
60. $(1, 2^{s+1}-1, 2^{s+t+1}-2^s-1, 2^{s+t}-2)$,
61. $(1, 2^{s+t}-2, 2^{s+1}-1, 2^{s+t+1}-2^s-1)$,
62. $(2^{s+1}-1, 1, 2^{s+t}-2, 2^{s+t+1}-2^s-1)$,
63. $(2^{s+1}-1, 1, 2^{s+t+1}-2^s-1, 2^{s+t}-2)$,
64. $(2^{s+1}-1, 2^{s+t+1}-2^s-1, 1, 2^{s+t}-2)$,
65. $(1, 2^{s+1}-1, 2^{s+t}-1, 2^{s+t+1}-2^s-2)$,
66. $(1, 2^{s+1}-1, 2^{s+t+1}-2^s-2, 2^{s+t}-1)$,
67. $(1, 2^{s+t}-1, 2^{s+1}-1, 2^{s+t+1}-2^s-2)$,
68. $(2^{s+1}-1, 1, 2^{s+t}-1, 2^{s+t+1}-2^s-2)$,
69. $(2^{s+1}-1, 1, 2^{s+t+1}-2^s-2, 2^{s+t}-1)$,
70. $(2^{s+1}-1, 2^{s+t}-1, 1, 2^{s+t+1}-2^s-2)$,
71. $(2^{s+t}-1, 1, 2^{s+1}-1, 2^{s+t+1}-2^s-2)$,
72. $(2^{s+t}-1, 2^{s+1}-1, 1, 2^{s+t+1}-2^s-2)$,
73. $(3, 2^{s+t}-3, 2^s-2, 2^{s+t+1}-1)$,
74. $(3, 2^{s+t}-3, 2^{s+t+1}-1, 2^s-2)$,
75. $(3, 2^{s+t+1}-1, 2^{s+t}-3, 2^s-2)$,

76. $(2^{s+t+1}-1, 3, 2^{s+t}-3, 2^s-2)$,
77. $(3, 2^{s+t}-1, 2^{s+t+1}-3, 2^s-2)$,
78. $(3, 2^{s+t+1}-3, 2^s-2, 2^{s+t}-1)$,
79. $(3, 2^{s+t+1}-3, 2^{s+t}-1, 2^s-2)$,
80. $(2^{s+t}-1, 3, 2^{s+t+1}-3, 2^s-2)$,
81. $(3, 2^s-1, 2^{s+t}-3, 2^{s+t+1}-2)$,
82. $(3, 2^{s+t}-3, 2^s-1, 2^{s+t+1}-2)$,
83. $(3, 2^{s+t}-3, 2^{s+t+1}-2, 2^s-1)$,
84. $(3, 2^s-1, 2^{s+t+1}-3, 2^{s+t}-2)$,
85. $(3, 2^{s+t+1}-3, 2^s-1, 2^{s+t}-2)$,
86. $(3, 2^{s+t+1}-3, 2^{s+t}-2, 2^s-1)$,
87. $(1, 2^{s+2}-2, 2^{s+t+1}-2^{s+1}-1, 2^{s+t}-2^s-1)$,
88. $(1, 2^{s+2}-1, 2^{s+t+1}-2^{s+1}-1, 2^{s+t}-2^s-2)$,
89. $(2^{s+2}-1, 1, 2^{s+t+1}-2^{s+1}-1, 2^{s+t}-2^s-2)$,
90. $(2^{s+2}-1, 2^{s+t+1}-2^{s+1}-1, 1, 2^{s+t}-2^s-2)$,
91. $(1, 2^{s+2}-1, 2^{s+t+1}-2^{s+1}-2, 2^{s+t}-2^s-1)$,
92. $(2^{s+2}-1, 1, 2^{s+t+1}-2^{s+1}-2, 2^{s+t}-2^s-1)$,
93. $(3, 2^{s+1}-3, 2^{s+t}-2^s-2, 2^{s+t+1}-1)$,
94. $(3, 2^{s+1}-3, 2^{s+t+1}-1, 2^{s+t}-2^s-2)$,
95. $(3, 2^{s+t+1}-1, 2^{s+1}-3, 2^{s+t}-2^s-2)$,
96. $(2^{s+t+1}-1, 3, 2^{s+1}-3, 2^{s+t}-2^s-2)$,
97. $(3, 2^{s+1}-3, 2^{s+t}-2^s-1, 2^{s+t+1}-2)$,
98. $(3, 2^{s+1}-3, 2^{s+t+1}-2, 2^{s+t}-2^s-1)$,
99. $(3, 2^{s+1}-3, 2^{s+t}-2, 2^{s+t+1}-2^s-1)$,
100. $(3, 2^{s+1}-3, 2^{s+t+1}-2^s-1, 2^{s+t}-2)$,
101. $(3, 2^{s+t+1}-3, 2^{s+1}-2, 2^{s+t}-2^s-1)$,
102. $(3, 2^{s+1}-3, 2^{s+t}-1, 2^{s+t+1}-2^s-2)$,
103. $(3, 2^{s+1}-3, 2^{s+t+1}-2^s-2, 2^{s+t}-1)$,
104. $(3, 2^{s+t}-1, 2^{s+1}-3, 2^{s+t+1}-2^s-2)$,
105. $(2^{s+t}-1, 3, 2^{s+1}-3, 2^{s+t+1}-2^s-2)$,
106. $(3, 2^{s+1}-1, 2^{s+t}-2^s-3, 2^{s+t+1}-2)$,
107. $(2^{s+1}-1, 3, 2^{s+t}-2^s-3, 2^{s+t+1}-2)$,
108. $(3, 2^{s+1}-1, 2^{s+t+1}-3, 2^{s+t}-2^s-2)$,
109. $(3, 2^{s+t+1}-3, 2^{s+1}-1, 2^{s+t}-2^s-2)$,
110. $(2^{s+1}-1, 3, 2^{s+t+1}-3, 2^{s+t}-2^s-2)$,
111. $(3, 2^{s+t}-3, 2^{s+1}-2, 2^{s+t+1}-2^s-1)$,
112. $(3, 2^{s+1}-1, 2^{s+t}-3, 2^{s+t+1}-2^s-2)$,
113. $(3, 2^{s+t}-3, 2^{s+1}-1, 2^{s+t+1}-2^s-2)$,
114. $(2^{s+1}-1, 3, 2^{s+t}-3, 2^{s+t+1}-2^s-2)$,
115. $(3, 2^{s+1}-1, 2^{s+t+1}-2^s-3, 2^{s+t}-2)$,
116. $(2^{s+1}-1, 3, 2^{s+t+1}-2^s-3, 2^{s+t}-2)$,
117. $(7, 2^{s+t}-5, 2^{s+t+1}-3, 2^s-2)$,
118. $(7, 2^{s+t+1}-5, 2^{s+t}-3, 2^s-2)$,
119. $(3, 2^{s+2}-3, 2^{s+t+1}-2^{s+1}-1, 2^{s+t}-2^s-2)$,
120. $(3, 2^{s+2}-3, 2^{s+t+1}-2^{s+1}-2, 2^{s+t}-2^s-1)$,

121. $(3, 2^{s+2}-1, 2^{s+t+1}-2^{s+1}-3, 2^{s+t}-2^s-2)$,
122. $(2^{s+2}-1, 3, 2^{s+t+1}-2^{s+1}-3, 2^{s+t}-2^s-2)$,
123. $(7, 2^{s+t+1}-5, 2^{s+1}-3, 2^{s+t}-2^s-2)$,
124. $(7, 2^{s+t}-5, 2^{s+1}-3, 2^{s+t+1}-2^s-2)$,
125. $(7, 2^{s+2}-5, 2^{s+t+1}-2^{s+1}-3, 2^{s+t}-2^s-2)$.

For $s = 2$ and $t \geqslant 2$,

126. $(3, 3, 2^{t+2}-4, 2^{t+3}-1)$,
127. $(3, 3, 2^{t+3}-1, 2^{t+2}-4)$,
128. $(3, 2^{t+3}-1, 3, 2^{t+2}-4)$,
129. $(2^{t+3}-1, 3, 3, 2^{t+2}-4)$,
130. $(3, 3, 2^{t+2}-1, 2^{t+3}-4)$,
131. $(3, 3, 2^{t+3}-4, 2^{t+2}-1)$,
132. $(3, 2^{t+2}-1, 3, 2^{t+3}-4)$,
133. $(2^{t+2}-1, 3, 3, 2^{t+3}-4)$,
134. $(3, 7, 2^{t+2}-5, 2^{t+3}-4)$,
135. $(7, 3, 2^{t+2}-5, 2^{t+3}-4)$,
136. $(7, 2^{t+2}-5, 3, 2^{t+3}-4)$,
137. $(3, 7, 2^{t+3}-5, 2^{t+2}-4)$,
138. $(7, 3, 2^{t+3}-5, 2^{t+2}-4)$,
139. $(7, 2^{t+3}-5, 3, 2^{t+2}-4)$,
140. $(7, 7, 2^{t+2}-8, 2^{t+3}-5)$,
141. $(7, 7, 2^{t+2}-7, 2^{t+3}-6)$,
142. $(7, 7, 2^{t+3}-7, 2^{t+2}-6)$,
143. $(7, 7, 2^{t+3}-8, 2^{t+2}-5)$.

For $s \geqslant 3$ and $t \geqslant 2$,

126. $(3, 2^s-3, 2^{s+t}-2, 2^{s+t+1}-1)$,
127. $(3, 2^s-3, 2^{s+t+1}-1, 2^{s+t}-2)$,
128. $(3, 2^{s+t+1}-1, 2^s-3, 2^{s+t}-2)$,
129. $(2^{s+t+1}-1, 3, 2^s-3, 2^{s+t}-2)$,
130. $(3, 2^s-3, 2^{s+t}-1, 2^{s+t+1}-2)$,
131. $(3, 2^s-3, 2^{s+t+1}-2, 2^{s+t}-1)$,
132. $(3, 2^{s+t}-1, 2^s-3, 2^{s+t+1}-2)$,
133. $(2^{s+t}-1, 3, 2^s-3, 2^{s+t+1}-2)$,
134. $(2^s-1, 3, 2^{s+t}-3, 2^{s+t+1}-2)$,
135. $(2^s-1, 3, 2^{s+t+1}-3, 2^{s+t}-2)$,
136. $(7, 2^{s+t}-5, 2^s-3, 2^{s+t+1}-2)$,
137. $(7, 2^{s+t+1}-5, 2^s-3, 2^{s+t}-2)$,
138. $(7, 2^{s+1}-5, 2^{s+t}-2^s-3, 2^{s+t+1}-2)$,
139. $(7, 2^{s+1}-5, 2^{s+t+1}-3, 2^{s+t}-2^s-2)$,
140. $(7, 2^{s+1}-5, 2^{s+t}-3, 2^{s+t+1}-2^s-2)$,
141. $(7, 2^{s+1}-5, 2^{s+t+1}-2^s-3, 2^{s+t}-2)$.

For $s = 3$ and $t \geqslant 2$,

142. $(7, 7, 2^{t+3} - 7, 2^{t+4} - 2)$,
143. $(7, 7, 2^{t+4} - 7, 2^{t+3} - 2)$.

For $s \geqslant 4$ and $t \geqslant 2$,

142. $(7, 2^s - 5, 2^{s+t} - 3, 2^{s+t+1} - 2)$,
143. $(7, 2^s - 5, 2^{s+t+1} - 3, 2^{s+t} - 2)$.

For $s \geqslant 2$ and $t = 2$,

144. $(1, 2^{s+2} - 2, 2^{s+2} - 1, 2^{s+2} + 2^s - 1)$,
145. $(1, 2^{s+2} - 1, 2^{s+2} - 2, 2^{s+2} + 2^s - 1)$,
146. $(2^{s+2} - 1, 1, 2^{s+2} - 2, 2^{s+2} + 2^s - 1)$,
147. $(1, 2^{s+2} - 1, 2^{s+2} - 1, 2^{s+2} + 2^s - 2)$,
148. $(2^{s+2} - 1, 1, 2^{s+2} - 1, 2^{s+2} + 2^s - 2)$,
149. $(2^{s+2} - 1, 2^{s+2} - 1, 1, 2^{s+2} + 2^s - 2)$,
150. $(3, 2^{s+2} - 3, 2^{s+2} - 1, 2^{s+2} + 2^s - 2)$,
151. $(3, 2^{s+2} - 3, 2^{s+2} - 2, 2^{s+2} + 2^s - 1)$,
152. $(3, 2^{s+2} - 1, 2^{s+2} - 3, 2^{s+2} + 2^s - 2)$,
153. $(2^{s+2} - 1, 3, 2^{s+2} - 3, 2^{s+2} + 2^s - 2)$,
154. $(7, 2^{s+2} - 5, 2^{s+2} - 3, 2^{s+2} + 2^s - 2)$.

For $s \geqslant 2$ and $t \geqslant 3$,

144. $(1, 2^{s+2} - 2, 2^{s+t} - 2^{s+1} - 1, 2^{s+t+1} - 2^s - 1)$,
145. $(1, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 2, 2^{s+t+1} - 2^s - 1)$,
146. $(2^{s+2} - 1, 1, 2^{s+t} - 2^{s+1} - 2, 2^{s+t+1} - 2^s - 1)$,
147. $(1, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1, 2^{s+t+1} - 2^s - 2)$,
148. $(2^{s+2} - 1, 1, 2^{s+t} - 2^{s+1} - 1, 2^{s+t+1} - 2^s - 2)$,
149. $(2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 1, 1, 2^{s+t+1} - 2^s - 2)$,
150. $(3, 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 2, 2^{s+t+1} - 2^s - 1)$,
151. $(3, 2^{s+2} - 3, 2^{s+t} - 2^{s+1} - 1, 2^{s+t+1} - 2^s - 2)$,
152. $(3, 2^{s+2} - 1, 2^{s+t} - 2^{s+1} - 3, 2^{s+t+1} - 2^s - 2)$,
153. $(2^{s+2} - 1, 3, 2^{s+t} - 2^{s+1} - 3, 2^{s+t+1} - 2^s - 2)$,
154. $(7, 2^{s+2} - 5, 2^{s+t} - 2^{s+1} - 3, 2^{s+t+1} - 2^s - 2)$.

We prove the theorem by proving the following.

Proposition 8.8.3. *The $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+t}+2^s-3}$ is generated by the elements listed in Theorem 8.8.2.*

We need the following for the proof of this proposition.

Lemma 8.8.4. *The following matrices are strictly inadmissible*

$$\begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&0&0\\1&1&0&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&1&0&0\\1&1&0&0\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&0&0\\0&0&1&1\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&1\\1&0&1&0\\1&0&0&1\\0&1&0&0\end{pmatrix}$$

$$\begin{pmatrix}1&1&1&0\\1&1&0&1\\1&1&0&0\\0&1&0&1\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&0&1\\1&1&0&0\\1&0&0&1\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&0&1&0\\0&1&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&0&0\\0&1&1&0\\0&0&0&1\end{pmatrix}$$

$$\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&1&0&0\\0&1&0&1\\0&0&1&0\end{pmatrix}\quad\begin{pmatrix}1&1&1&0\\1&1&1&0\\1&1&0&0\\1&0&1&0\\0&0&0&1\end{pmatrix}\quad\begin{pmatrix}1&1&0&1\\1&1&0&1\\1&1&0&0\\1&0&0&1\\0&0&1&0\end{pmatrix}.$$

Proof. The monomials corresponding to the above matrices respectively are

$$(15,15,17,2),(7,7,27,8),(3,15,15,16),(15,3,15,16),(15,15,3,16),$$
$$(15,15,16,3),(7,7,11,24),(15,19,5,10),(7,15,17,10),(15,7,17,10),$$
$$(7,11,15,16),(7,15,11,16),(7,15,16,11),(15,7,11,16),(15,7,16,11).$$

We prove the lemma for the matrices associated with the monomials

$$(15,15,17,2),(7,7,27,8),(3,15,15,16),(15,15,16,3),(7,7,11,24),$$
$$(15,19,5,10),(15,7,17,10),(7,11,15,16),(15,7,11,16),(15,7,16,11).$$

By a direct computation, we have

$$\begin{aligned}(15,15,17,2) &= Sq^1(15,15,15,3)+Sq^2(15,15,15,2)\\
&\quad+Sq^4\big((15,13,15,2)+(15,12,15,3)\big)+Sq^8\big((9,15,15,2)\\
&\quad+(11,13,15,2)+(8,15,15,3)+(11,12,15,3)\big)+(9,23,15,2)\\
&\quad+(9,15,23,2)+(11,21,15,2)+(11,13,23,2)+(15,13,19,2)\\
&\quad+(8,23,15,3)+(8,15,23,3)+(15,12,19,3)+(15,15,16,3)\\
&\quad+(11,20,15,3)+(11,12,23,3)\mod \mathcal{L}_4(3;3;2;2;1),\\
(7,7,27,8) &= Sq^1\big((7,7,27,7)+(7,5,29,7)+(7,7,21,13)\\
&\quad+(7,7,15,19)+(7,5,15,21)\big)+Sq^2\big((7,6,27,7)\\
&\quad+(7,7,26,7)+(7,7,22,11)+(7,3,30,7)+(7,7,19,14)\\
&\quad+(7,7,15,18)+(7,6,15,19)+(7,3,23,14)\big)\\
&\quad+Sq^4\big((4,7,27,7)+(5,6,27,7)+(5,7,26,7)+(11,5,22,7)\end{aligned}$$

$$
\begin{aligned}
&+ (5,7,22,11) + (5,3,30,7) + (5,7,19,14) + (11,5,15,14)\\
&+ (5,7,15,18) + (4,7,15,19) + (5,6,15,19) + (5,3,23,14)\big)\\
&+ Sq^8\big((7,5,22,7) + (7,5,15,14)\big) + (4,11,27,7)\\
&+ (4,7,27,11) + (5,10,27,7) + (5,6,27,11) + (7,6,27,9)\\
&+ (7,7,26,9) + (5,11,26,7) + (7,5,26,11) + (5,11,22,11)\\
&+ (7,7,24,11) + (5,3,30,11) + (7,3,30,9) + (5,11,19,14)\\
&+ (5,11,15,18) + (4,11,15,19) + (5,10,15,19)\\
&+ (7,3,25,14) + (5,3,27,14) \mod \mathcal{L}_4(3;3;2;2;1),\\
(3,15,15,16) &= Sq^1(3,15,15,15) + Sq^2(2,15,15,15)\\
&+ Sq^4\big((3,12,15,15) + (3,15,12,15)\big) + Sq^8(3,11,12,15)\\
&+ (2,17,15,15) + (2,15,17,15) + (2,15,15,17)\\
&+ (3,12,19,15) + (3,12,15,19) + (3,15,12,19)\\
&+ (3,11,20,15) + (3,11,12,23) \mod \mathcal{L}_4(3;3;2;2;1),\\
(7,7,11,24) &= Sq^1\big((7,7,7,27) + (7,5,7,29) + (7,7,13,21)\\
&+ (7,7,19,15) + (7,5,21,15)\big) + Sq^2\big((7,7,11,22) + (7,7,7,26)\\
&+ (7,6,7,27) + (7,3,7,30) + (7,7,14,19) + (7,7,18,15)\\
&+ (7,6,19,15) + (7,3,22,15)\big) + Sq^4\big((5,7,11,22) + (11,5,7,22)\\
&+ (5,7,7,26) + (4,7,7,27) + (5,6,7,27) + (5,3,7,30)\\
&+ (5,7,14,19) + (11,5,14,15) + (5,7,18,15) + (4,7,19,15)\\
&+ (5,6,19,15) + (5,3,22,15)\big) + Sq^8\big((7,5,7,22) + (7,5,14,15)\big)\\
&+ (5,11,11,22) + (7,5,11,26) + (7,7,9,26) + (5,11,7,26)\\
&+ (7,7,8,27) + (4,11,7,27) + (4,7,11,27) + (7,6,9,27)\\
&+ (5,10,7,27) + (5,6,11,27) + (7,3,9,30) + (5,3,11,30)\\
&+ (5,11,14,19) + (5,11,18,15) + (4,11,19,15) + (5,10,19,15)\\
&+ (7,3,24,15) + (5,3,26,15) \mod \mathcal{L}_4(3;3;2;2;1),\\
(15,15,16,3) &= Sq^1\big((15,15,13,5) + (15,15,11,7) + (15,15,9,9)\\
&+ (15,15,7,11) + (15,15,5,13)\big) + Sq^2\big((15,15,14,3) + (15,15,11,6)\\
&+ (15,15,10,7) + (15,15,7,10) + (15,15,6,11) + (15,15,3,14)\big)\\
&+ Sq^4\big((15,13,14,3) + (15,13,11,6) + (15,12,11,7) + (15,13,10,7)\\
&+ (15,13,7,10) + (15,12,7,11) + (15,13,6,11) + (15,13,3,14)\big)\\
&+ Sq^8\big((9,15,14,3) + (11,13,14,3) + (9,15,11,6) + (11,13,11,6)\\
&+ (8,15,11,7) + (11,12,11,7) + (9,15,10,7) + (11,13,10,7)\\
&+ (9,15,7,10) + (11,13,7,10) + (8,15,7,11) + (11,12,7,11)
\end{aligned}
$$

$$\begin{aligned}
&+ (9,15,6,11) + (11,13,6,11) + (9,15,3,14) + (11,13,3,14)\big)\\
&+ (9,23,14,3) + (9,15,22,3) + (15,13,18,3) + (11,21,14,3)\\
&+ (11,13,22,3) + (9,23,11,6) + (9,15,19,6) + (11,21,11,6)\\
&+ (11,13,19,6) + (8,23,11,7) + (8,15,19,7) + (11,20,11,7)\\
&+ (11,12,19,7) + (9,23,10,7) + (9,15,18,7) + (11,21,10,7)\\
&+ (11,13,18,7) + (9,23,7,10) + (9,15,7,18) + (11,21,7,10)\\
&+ (11,13,7,18) + (8,23,7,11) + (8,15,7,19) + (11,20,7,11)\\
&+ (11,12,7,19) + (9,23,6,11) + (9,15,6,19) + (11,21,6,11)\\
&+ (11,13,6,19) + (15,15,3,16) + (9,23,3,14) + (9,15,3,22)\\
&+ (15,13,3,18) + (11,21,3,14) + (11,13,3,22) \mod \mathcal{L}_4(3;3;2;2;1),
\end{aligned}$$

$$\begin{aligned}
(15,19,5,10) &= Sq^1(15,15,7,11) + Sq^2(15,15,7,10) + Sq^4\big((15,15,5,10)\\
&+ (15,15,4,11)\big) + Sq^8\big((11,15,5,10) + (9,15,7,10)\\
&+ (8,15,7,11) + (11,15,4,11)\big) + (11,23,5,10) + (11,15,5,18)\\
&+ (9,23,7,10) + (9,15,7,18) + (15,17,7,10) + (8,23,7,11)\\
&+ (8,15,7,19) + (15,16,7,11) + (11,23,4,11)\\
&+ (11,15,4,19) + (15,19,4,11) \mod \mathcal{L}_4(3;3;2;2;1),
\end{aligned}$$

$$\begin{aligned}
(15,7,17,10) &= Sq^1(15,7,15,11) + Sq^2(15,7,15,10) + Sq^4\big((15,5,15,10)\\
&+ (15,4,15,11)\big) + Sq^8\big((9,7,15,10) + (8,7,15,11) + (11,5,15,10)\\
&+ (11,4,15,11)\big) + (11,5,23,10) + (11,5,15,18) + (15,5,19,10)\\
&+ (9,7,23,10) + (9,7,15,18) + (11,4,23,11)\\
&+ (11,4,15,19) + (15,4,19,11) + (8,7,23,11)\\
&+ (8,7,15,19) + (15,7,16,11) \mod \mathcal{L}_4(3;3;2;2;1),
\end{aligned}$$

$$\begin{aligned}
(7,11,15,16) &= Sq^1\big((7,11,15,15) + (7,7,19,15) + (7,7,11,23)\big)\\
&+ Sq^2(7,10,15,15) + Sq^4\big((4,11,15,15) + (5,10,15,15)\\
&+ (11,7,12,15) + (4,7,19,15) + (4,7,11,23)\big) + Sq^8(7,7,12,15)\\
&+ (4,11,19,15) + (4,11,15,19) + (7,10,17,15) + (7,10,15,17)\\
&+ (5,10,19,15) + (5,10,15,19) + (7,11,12,19) + (7,8,19,15)\\
&+ (4,11,19,15) + (7,8,11,23) + (7,7,11,24)\\
&+ (4,11,11,23) + (4,7,11,27) \mod \mathcal{L}_4(3;3;2;2;1),
\end{aligned}$$

$$\begin{aligned}
(15,7,11,16) &= Sq^1(15,7,11,15) + Sq^2(15,7,10,15) + Sq^4\big((15,4,11,15)\\
&+ (15,5,10,15)\big) + Sq^8\big((8,7,11,15) + (11,4,11,15) + (9,7,10,15)\\
&+ (11,5,10,15)\big) + (8,7,19,15) + (8,7,11,23) + (15,4,11,19)\\
&+ (11,4,19,15) + (11,4,11,23) + (9,7,18,15)
\end{aligned}$$

$$\begin{aligned}
&+ (9,7,10,23) + (15,5,10,19) + (11,5,18,15)\\
&+ (11,5,10,23) + (15,7,10,17) \mod \mathcal{L}_4(3;3;2;2;1),\\
(15,7,16,11) &= Sq^1(15,7,13,13) + Sq^2\big((15,7,14,11) + (15,7,11,14)\big)\\
&+ Sq^4\big((15,5,14,11) + (15,5,11,14)\big) + Sq^8\big((9,7,14,11)\\
&+ (11,5,14,11) + (9,7,11,14) + (11,5,11,14)\big) + (9,7,22,11)\\
&+ (9,7,14,19) + (11,5,22,11) + (11,5,14,19) + (15,5,18,11)\\
&+ (15,7,11,16) + (9,7,19,14) + (9,7,11,22) + (15,5,11,18)\\
&+ (11,5,19,14) + (11,5,11,22) \mod \mathcal{L}_4(3;3;2;2;1).
\end{aligned}$$

The lemma is proved. □

The results in Section 7, the lemmas in Sections 3, 5, 6, Lemmas 8.3.4, 8.3.5, 8.8.4 and Theorem 2.5 imply Proposition 8.8.3

Next, we prove that the classes listed in Theorem 8.8.2 are linearly independent.

Proposition 8.8.5. *The elements* $[a_{1,2,2,j}], 1 \leqslant j \leqslant 154$, *are linearly independent in* $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{49}$.

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{154} \gamma_j [a_{1,2,2,j}] = 0, \tag{8.8.5.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.8.5.1) and we obtain

$$\begin{aligned}
&\gamma_1[3,15,31] + \gamma_2[3,31,15] + \gamma_{15}[15,3,31] + \gamma_{16}[15,31,3]\\
&\quad + \gamma_{29}[31,3,15] + \gamma_{30}[31,15,3] + \gamma_{37}[7,11,31] + \gamma_{38}[7,31,11]\\
&\quad + \gamma_{51}[31,7,11] + \gamma_{55}[7,15,27] + \gamma_{56}[7,27,15]\\
&\quad + \gamma_{61}[15,7,27] + \gamma_{87}[15,23,11] + \gamma_{144}[15,15,19] = 0,\\
&\gamma_3[3,15,31] + \gamma_5[3,31,15] + \gamma_{\{13,126\}}[15,3,31] + \gamma_{18}[15,31,3]\\
&\quad + \gamma_{\{26,131\}}[31,3,15] + \gamma_{28}[31,15,3] + \gamma_{41}[7,11,31] + \gamma_{39}[7,31,11]\\
&\quad + \gamma_{\{50,143\}}[31,7,11] + \gamma_{\{57,66\}}[7,15,27] + \gamma_{66}[7,27,15]\\
&\quad + \gamma_{\{59,66,140\}}[15,7,27] + \gamma_{91}[15,23,11] + \gamma_{\{66,91,145\}}[15,15,19] = 0,\\
&\gamma_4[3,15,31] + \gamma_6[3,31,15] + \gamma_{\{14,127\}}[15,3,31] + \gamma_{\{17,128\}}[15,31,3]\\
&\quad + \gamma_{\{25,130\}}[31,3,15] + \gamma_{\{27,132\}}[31,15,3] + \gamma_{42}[7,11,31] + \gamma_{43}[7,31,11]\\
&\quad + \gamma_{\{49,134\}}[31,7,11] + \gamma_{\{65,88,147\}}[7,15,27] + \gamma_{\{65,147\}}[7,27,15]\\
&\quad + \gamma_{\{60,65,137\}}[15,7,27] + \gamma_{67}[15,23,11] + \gamma_{\{65,67\}}[15,15,19] = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_{\{19,73,93,126\}}[3,15,31] + \gamma_{\{32,78,103,131\}}[3,31,15] + \gamma_7[15,3,31]\\
&\quad + \gamma_{35}[15,31,3] + \gamma_{10}[31,3,15] + \gamma_{23}[31,15,3] + \gamma_{44}[7,11,31]\\
&\quad + \gamma_{\{53,143\}}[7,31,11] + \gamma_{40}[31,7,11] + \gamma_{\{62,140\}}[7,15,27]\\
&\quad + \gamma_{69}[7,27,15] + \gamma_{58}[15,7,27] + \gamma_{92}[15,23,11] + \gamma_{146}[15,15,19] = 0,\\
&\gamma_{\{20,74,94,127\}}[3,15,31] + \gamma_{\{31,79,102,130,150\}}[3,31,15] + \gamma_8[15,3,31]\\
&\quad + \gamma_{\{34,133\}}[15,31,3] + \gamma_{11}[31,3,15] + \gamma_{\{22,129\}}[31,15,3] + \gamma_{45}[7,11,31]\\
&\quad + \gamma_{\{52,135\}}[7,31,11] + \gamma_{\{63,138,148\}}[7,15,27] + \gamma_{\{68,148\}}[7,27,15]\\
&\quad + \gamma_{47}[31,7,11] + \gamma_{89}[15,7,27] + \gamma_{71}[15,23,11] + \gamma_{148}[15,15,19] = 0,\\
&\gamma_{\{33,77,104,121,132,152\}}[3,15,31] + \gamma_{\{21,75,95,128\}}[3,31,15]\\
&\quad + \gamma_{\{36,80,105,122,133,153\}}[15,3,31] + \gamma_9[15,31,3] + \gamma_{\{24,76,96,129\}}[31,3,15]\\
&\quad + \gamma_{12}[31,15,3] + \gamma_{\{54,117,124,125,136,154\}}[7,11,31] + \gamma_{46}[7,31,11]\\
&\quad + \gamma_{48}[31,7,11] + \gamma_{70}[7,15,27] + \gamma_{\{64,118,123,139\}}[7,27,15]\\
&\quad + \gamma_{72}[15,7,27] + \gamma_{90}[15,23,11] + \gamma_{149}[15,15,19] = 0.
\end{aligned}$$

Computing from these equalities, we obtain

$$\text{(8.8.5.2)}\quad \begin{cases}
\gamma_j = 0,\ j = 1,\dots,12,15,16,18,23,28,29,\\
\qquad\qquad 30,35,37,\dots,48,51,55,56,57,58,61,\\
\qquad\qquad 65,\dots,72,87,\dots,92,144,\dots,149,\\
\gamma_{\{13,126\}} = \gamma_{\{26,131\}} = \gamma_{\{50,143\}} = \gamma_{\{59,140\}} = \gamma_{\{14,127\}} = 0,\\
\gamma_{\{17,128\}} = \gamma_{\{25,130\}} = \gamma_{\{27,132\}} = \gamma_{\{60,137\}} = \gamma_{\{53,143\}} = 0,\\
\gamma_{\{49,134\}} = \gamma_{\{19,73,93,126\}} = \gamma_{\{32,78,103,131\}} = \gamma_{\{62,140\}} = 0,\\
\gamma_{\{20,74,94,127\}} = \gamma_{\{31,79,102,130,150\}} = \gamma_{\{34,133\}} = \gamma_{\{22,129\}} = 0,\\
\gamma_{\{52,135\}} = \gamma_{\{63,138\}} = \gamma_{\{21,75,95,128\}} = \gamma_{\{24,76,96,129\}} = 0,\\
\gamma_{\{33,77,104,121,132,152\}} = \gamma_{\{36,80,105,122,133,153\}} = 0,\\
\gamma_{\{54,117,124,125,136,154\}} = \gamma_{\{64,118,123,139\}} = 0.
\end{cases}$$

With the aid of (8.8.5.2), the homomorphisms g_1, g_2 send (8.8.5.1) to

$$\begin{aligned}
&\gamma_{19}[3,15,31] + \gamma_{32}[3,31,15] + \gamma_{73}[15,3,31] + \gamma_{83}[15,31,3]\\
&\quad + \gamma_{78}[31,3,15] + \gamma_{86}[31,15,3] + \gamma_{93}[7,11,31] + \gamma_{\{50,98\}}[7,31,11]\\
&\quad + \gamma_{101}[31,7,11] + \gamma_{\{59,99\}}[7,15,27] + \gamma_{103}[7,27,15]\\
&\quad + \gamma_{111}[15,7,27] + \gamma_{120}[15,23,11] + \gamma_{151}[15,15,19] = 0,\\
&\gamma_{20}[3,15,31] + \gamma_{31}[3,31,15] + \gamma_{74}[15,3,31] + \gamma_{\{82,136\}}[15,31,3]\\
&\quad + \gamma_{79}[31,3,15] + \gamma_{\{85,139\}}[31,15,3] + \gamma_{94}[7,11,31] + \gamma_{109}[31,7,11]\\
&\quad + \gamma_{97}[7,31,11] + \gamma_{\{100,150\}}[7,15,27] + \gamma_{119}[15,7,27]\\
&\quad + \gamma_{\{102,150\}}[7,27,15] + \gamma_{113}[15,23,11] + \gamma_{150}[15,15,19] = 0.
\end{aligned}$$

These equalities imply

$$\text{(8.8.5.3)}\qquad \begin{cases} \gamma_j = 0,\ j = 19, 20, 31, 32, 73, 74, 78, 79, 83, 86, 93, 94, 97, \\ \qquad 100, 101, 102, 103, 109, 111, 113, 119, 120, 150, 151, \\ \gamma_{\{50,98\}} = \gamma_{\{59,99\}} = \gamma_{\{82,136\}} = \gamma_{\{85,139\}} = 0. \end{cases}$$

With the aid of (8.8.5.2) and (8.8.5.3), the homomorphisms g_3, g_4 send (8.8.5.1) to

$$\begin{aligned} &\gamma_{21}[3,15,31] + \gamma_{33}[3,31,15] + \gamma_{75}[15,3,31] + a_1[15,31,3] \\ &\quad + \gamma_{77}[31,3,15] + a_2[31,15,3] + \gamma_{95}[7,11,31] + a_3[7,31,11] \\ &\quad + \gamma_{\{108,117,142\}}[31,7,11] + \gamma_{\{64,121,123,139,152\}}[7,15,27] \\ &\quad + \gamma_{\{104,121,152\}}[7,27,15] + \gamma_{\{118,121\}}[15,7,27] \\ &\quad + a_4[15,23,11] + \gamma_{\{50,115,121,125,142,152\}}[15,15,19] = 0, \\ &\gamma_{24}[3,15,31] + \gamma_{36}[3,31,15] + \gamma_{76}[15,3,31] + a_5[15,31,3] \\ &\quad + \gamma_{80}[31,3,15] + a_6[31,15,3] + \gamma_{96}[7,11,31] + a_7[7,31,11] \\ &\quad + a_8[31,7,11] + \gamma_{\{122,153\}}[7,15,27] + \gamma_{\{105,122,153\}}[7,27,15] \\ &\quad + \gamma_{122}[15,7,27] + a_9[15,23,11] + a_{10}[15,15,19] = 0, \end{aligned}$$

where

$$\begin{aligned} &a_1 = \gamma_{\{22,24,34,36,52,76,81,96,105,107,114,153\}},\ a_2 = \gamma_{\{22,24,63,76,80,84,96,110,116,122\}}, \\ &a_3 = \gamma_{\{54,59,106,124,136,141\}},\ a_4 = \gamma_{\{50,112,115,125,141,142,154\}}, \\ &a_5 = \gamma_{\{13,17,21,25,27,33,49,75,81,82,85,95,99,104,106,112,152\}}, \\ &a_6 = \gamma_{\{14,17,21,26,60,75,77,84,85,95,98,108,115,121\}}, \\ &a_7 = \gamma_{\{54,59,64,107,136,139,141\}},\ a_8 = \gamma_{\{110,117,118,123,125,142\}}, \\ &a_9 = \gamma_{\{50,64,114,116,118,123,124,139,141,142\}},\ a_{10} = \gamma_{\{50,64,116,122,139,142,153\}}. \end{aligned}$$

From the above equalities, we obtain

$$\text{(8.8.5.4)}\qquad \begin{cases} a_i = 0,\ i = 1, 2, \ldots, 10, \quad \gamma_j = 0,\ j = 21, 24, 33, 36, \\ \qquad 75, 76, 77, 80, 95, 96, 105, 122, 153, \\ \gamma_{\{108,117,142\}} = \gamma_{\{64,121,123,139,152\}} = \gamma_{\{104,121,152\}} = 0, \\ \gamma_{\{118,121\}} = \gamma_{\{50,115,121,125,142,152\}} = 0. \end{cases}$$

With the aid of (8.8.5.2), (8.8.5.3) and (8.8.5.4), the homomorphism h sends (8.8.5.1) to

$$\begin{aligned} &\gamma_{\{59,81,107,114\}}[3,15,31] + \gamma_{\{50,84,110,116\}}[3,31,15] + \gamma_{54}[15,3,31] \\ &\quad + \gamma_{117}[15,31,3] + \gamma_{64}[31,3,15] + \gamma_{118}[31,15,3] + a_{11}[7,11,31] \\ &\quad + \gamma_{\{108,110,117,142\}}[7,31,11] + \gamma_{123}[31,7,11] + a_{12}[7,15,27] \\ &\quad + a_{13}[7,27,15] + \gamma_{\{64,85,104,123,124\}}[15,7,27] \\ &\quad + \gamma_{125}[15,23,11] + \gamma_{\{64,85,118,123,152,154\}}[15,15,19] = 0, \end{aligned}$$

where

$$a_{11} = \gamma_{\{54,59,82,106,107,124,141,155\}},$$
$$a_{12} = \gamma_{\{64,85,112,114,123,141,154,155\}}, \quad a_{13} = \gamma_{\{50,64,85,115,116,123,125,142\}}.$$

From the above equalities, it implies

$$\text{(8.8.5.5)} \quad \begin{cases} a_{11} = a_{12} = a_{13} = 0,\ \gamma_j = 0,\ j = 54, 64, 117, 118, 123, 125, \\ \gamma_{\{59,81,107,114\}} = \gamma_{\{50,84,110,116\}} = 0, \\ \gamma_{\{108,110,142\}} = \gamma_{\{85,104,124\}} = \gamma_{\{85,152,154\}} = 0, \end{cases}$$

Combining (8.8.5.2), (8.8.5.3), (8.8.5.4) and (8.8.5.5), we get $\gamma_j = 0$ for any j. The proposition is proved. □

Proposition 8.8.6. *For $t \geqslant 3$, the elements $[a_{1,t,2,j}], 1 \leqslant j \leqslant 154$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+3}+2^{t+2}+1}$.*

Proof. Suppose that there is a linear relation

$$\text{(8.8.6.1)} \quad \sum_{j=1}^{154} \gamma_j [a_{1,t,2,j}] = 0,$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.8.6.1), we obtain

$$\begin{aligned}
&\gamma_1 w_{1,t,2,1} + \gamma_2 w_{1,t,2,2} + \gamma_{15} w_{1,t,2,3} + \gamma_{16} w_{1,t,2,4} + \gamma_{29} w_{1,t,2,5} \\
&\quad + \gamma_{30} w_{1,t,2,6} + \gamma_{37} w_{1,t,2,7} + \gamma_{38} w_{1,t,2,8} + \gamma_{51} w_{1,t,2,9} + \gamma_{55} w_{1,t,2,10} \\
&\quad + \gamma_{56} w_{1,t,2,11} + \gamma_{61} w_{1,t,2,12} + \gamma_{87} w_{1,t,2,13} + \gamma_{144} w_{1,t,2,14} = 0, \\
&\gamma_3 w_{1,t,2,1} + \gamma_5 w_{1,t,2,2} + \gamma_{\{13,126\}} w_{1,t,2,3} + \gamma_{18} w_{1,t,2,4} + \gamma_{\{26,131\}} w_{1,t,2,5} \\
&\quad + \gamma_{28} w_{1,t,2,6} + \gamma_{41} w_{1,t,2,7} + \gamma_{39} w_{1,t,2,8} + \gamma_{\{50,143\}} w_{1,t,2,9} + \gamma_{57} w_{1,t,2,10} \\
&\quad + \gamma_{66} w_{1,t,2,11} + \gamma_{\{59,140\}} w_{1,t,2,12} + \gamma_{91} w_{1,t,2,13} + \gamma_{\{66,91,145\}} w_{1,t,2,14} = 0, \\
&\gamma_4 w_{1,t,2,1} + \gamma_6 w_{1,t,2,2} + \gamma_{\{14,127\}} w_{1,t,2,3} + \gamma_{\{17,128\}} w_{1,t,2,4} \\
&\quad + \gamma_{\{25,130\}} w_{1,t,2,5} + \gamma_{\{27,132\}} w_{1,t,2,6} + \gamma_{42} w_{1,t,2,7} + \gamma_{43} w_{1,t,2,8} \\
&\quad + \gamma_{\{49,134\}} w_{1,t,2,9} + \gamma_{67} w_{1,t,2,10} + \gamma_{65} w_{1,t,2,11} + \gamma_{\{60,137\}} w_{1,t,2,12} \\
&\quad + \gamma_{\{67,147\}} w_{1,t,2,13} + \gamma_{\{65,88,147\}} w_{1,t,2,14} = 0, \\
&\gamma_{\{19,73,93,126\}} w_{1,t,2,1} + \gamma_{\{32,78,103,131\}} w_{1,t,2,2} + \gamma_7 w_{1,t,2,3} \\
&\quad + \gamma_{35} w_{1,t,2,4} + \gamma_{10} w_{1,t,2,5} + \gamma_{23} w_{1,t,2,6} + \gamma_{44} w_{1,t,2,7} \\
&\quad + \gamma_{\{53,143\}} w_{1,t,2,8} + \gamma_{40} w_{1,t,2,9} + \gamma_{\{62,140\}} w_{1,t,2,10} + \gamma_{69} w_{1,t,2,11} \\
&\quad + \gamma_{58} w_{1,t,2,12} + \gamma_{92} w_{1,t,2,13} + \gamma_{146} w_{1,t,2,14} = 0, \\
&\gamma_{\{20,74,94,127\}} w_{1,t,2,1} + \gamma_{\{31,79,102,130\}} w_{1,t,2,2} + \gamma_8 w_{1,t,2,3} \\
&\quad + \gamma_{\{34,133\}} w_{1,t,2,4} + \gamma_{11} w_{1,t,2,5} + \gamma_{\{22,129\}} w_{1,t,2,6} + \gamma_{45} w_{1,t,2,7} \\
&\quad + \gamma_{\{52,135\}} w_{1,t,2,8} + \gamma_{47} w_{1,t,2,9} + \gamma_{\{63,138\}} w_{1,t,2,10} + \gamma_{68} w_{1,t,2,11} \\
&\quad + \gamma_{71} w_{1,t,2,12} + \gamma_{\{71,148\}} w_{1,t,2,13} + \gamma_{\{71,89\}} w_{1,t,2,14} = 0,
\end{aligned}$$

$$\gamma_{\{33,77,104,132\}}w_{1,t,2,1} + \gamma_{\{21,75,95,128\}}w_{1,t,2,2} + \gamma_{\{36,80,105,133\}}w_{1,t,2,3} + \gamma_9 w_{1,t,2,4} + \gamma_{\{24,76,96,129\}}w_{1,t,2,5} + \gamma_{12}w_{1,t,2,6} + \gamma_{\{54,117,124,136\}}w_{1,t,2,7} + \gamma_{46}w_{1,t,2,8} + \gamma_{48}w_{1,t,2,9} + \gamma_{70}w_{1,t,2,10} + \gamma_{\{64,118,123,139\}}w_{1,t,2,11} + \gamma_{72}w_{1,t,2,12} + \gamma_{90}w_{1,t,2,13} + \gamma_{149}w_{1,t,2,14} = 0.$$

Computing from these equalities gives

$$(8.8.6.2)\quad \begin{cases} \gamma_j = 0,\ j = 1,\dots,12,15,16,18,23,28,29, \\ \qquad\qquad 30,35,37,\dots,48,51,55,56,57,58,61, \\ \qquad\qquad 65,\dots,72,87,\dots,92,144,\dots,149, \\ \gamma_{\{13,126\}} = \gamma_{\{26,131\}} = \gamma_{\{50,143\}} = \gamma_{\{59,140\}} = \gamma_{\{14,127\}} = 0, \\ \gamma_{\{17,128\}} = \gamma_{\{25,130\}} = \gamma_{\{27,132\}} = \gamma_{\{49,134\}} = \gamma_{\{60,137\}} = 0, \\ \gamma_{\{19,73,93,126\}} = \gamma_{\{32,78,103,131\}} = \gamma_{\{53,143\}} = \gamma_{\{62,140\}} = 0, \\ \gamma_{\{20,74,94,127\}} = \gamma_{\{31,79,102,130\}} = \gamma_{\{34,133\}} = \gamma_{\{22,129\}} = 0, \\ \gamma_{\{52,135\}} = \gamma_{\{63,138\}} = \gamma_{\{33,77,104,132\}} = \gamma_{\{21,75,95,128\}} = 0, \\ \gamma_{\{36,80,105,133\}} = \gamma_{\{24,76,96,129\}} = 0, \\ \gamma_{\{54,117,124,136\}} = \gamma_{\{64,118,123,139\}} = 0. \end{cases}$$

With the aid of (8.8.6.2), the homomorphisms g_1, g_2 send (8.8.6.1) to

$$\gamma_{19}w_{1,t,2,1} + \gamma_{32}w_{1,t,2,2} + \gamma_{73}w_{1,t,2,3} + \gamma_{83}w_{1,t,2,4} + \gamma_{78}w_{1,t,2,5} + \gamma_{86}w_{1,t,2,6} + \gamma_{93}w_{1,t,2,7} + \gamma_{\{50,98\}}w_{1,t,2,8} + \gamma_{101}w_{1,t,2,9} + \gamma_{\{59,99\}}w_{1,t,2,10} + \gamma_{103}w_{1,t,2,11} + \gamma_{111}w_{1,t,2,12} + \gamma_{120}w_{1,t,2,13} + \gamma_{150}w_{1,t,2,14} = 0,$$

$$\gamma_{20}w_{1,t,2,1} + \gamma_{31}w_{1,t,2,2} + \gamma_{74}w_{1,t,2,3} + \gamma_{\{82,136\}}w_{1,t,2,4} + \gamma_{79}w_{1,t,2,5} + \gamma_{\{85,139\}}w_{1,t,2,6} + \gamma_{94}w_{1,t,2,7} + \gamma_{97}w_{1,t,2,8} + \gamma_{109}w_{1,t,2,9} + \gamma_{100}w_{1,t,2,10} + \gamma_{102}w_{1,t,2,11} + \gamma_{113}w_{1,t,2,12} + \gamma_{\{113,151\}}w_{1,t,2,13} + \gamma_{\{113,119\}}w_{1,t,2,14} = 0.$$

These equalities imply

$$(8.8.6.3)\quad \begin{cases} \gamma_j = 0,\ j = 19,20,31,32,73,74,78,79,83,86,93,94, \\ \qquad 97,100,101,102,103,109,111,113,119,120,150,151, \\ \gamma_{\{50,98\}} = \gamma_{\{59,99\}} = \gamma_{\{82,136\}} = \gamma_{\{85,139\}} = 0. \end{cases}$$

With the aid of (8.8.6.2) and (8.8.6.3), the homomorphisms g_3, g_4 send (8.8.6.1) to

$$\gamma_{21}w_{1,t,2,1} + \gamma_{33}w_{1,t,2,2} + \gamma_{75}w_{1,t,2,3} + a_1 w_{1,t,2,4} + \gamma_{77}w_{1,t,2,5} + a_2 w_{1,t,2,6} + \gamma_{95}w_{1,t,2,7} + a_3 w_{1,t,2,8} + \gamma_{\{108,117,142\}}w_{1,t,2,9} + a_4 w_{1,t,2,10} + \gamma_{104}w_{1,t,2,11} + \gamma_{\{112,118,141\}}w_{1,t,2,12} + a_5 w_{1,t,2,13} + \gamma_{\{112,121,141\}}w_{1,t,2,14} = 0,$$

$$\gamma_{24}w_{1,t,2,1} + \gamma_{36}w_{1,t,2,2} + \gamma_{76}w_{1,t,2,3} + a_6 w_{1,t,2,4} + \gamma_{80}w_{1,t,2,5} + a_7 w_{1,t,2,6} + \gamma_{96}w_{1,t,2,7} + a_8 w_{1,t,2,8} + a_9 w_{1,t,2,9} + \gamma_{\{50,64,116,139,142\}}w_{1,t,2,10} + \gamma_{105}w_{1,t,2,11} + a_{10}w_{1,t,2,12} + a_{11}w_{1,t,2,13} + a_{12}w_{1,t,2,14} = 0,$$

where

$$a_1 = \gamma_{\{22,24,34,36,52,76,81,96,105,107,114,153\}},\ a_2 = \gamma_{\{22,24,63,76,80,84,96,110,116,122\}},$$
$$a_3 = \gamma_{\{54,59,106,124,136,141,154\}},\ a_4 = \gamma_{\{50,64,115,123,125,139,142\}},$$
$$a_5 = \gamma_{\{50,112,115,121,125,141,142,152\}},\ a_8 = \gamma_{\{54,59,64,107,136,139,141\}},$$
$$a_6 = \gamma_{\{13,17,21,25,27,33,49,75,81,82,85,95,99,104,106,112,152\}},$$
$$a_7 = \gamma_{\{14,17,21,26,60,75,77,84,85,95,98,108,115,121\}},\ a_9 = \gamma_{\{110,117,118,123,125,142\}}$$
$$a_{11} = \gamma_{\{50,64,114,116,118,122,123,124,139,141,142,153,154\}},$$
$$a_{10} = \gamma_{\{114,118,123,124,141,154\}},\ a_{12} = \gamma_{\{114,118,122,123,124,141,154\}}.$$

From the above equalities, we obtain

$$\text{(8.8.6.4)}\qquad \begin{cases} a_i = 0,\ i = 1,2,\ldots,12, \quad \gamma_j = 0,\ j = 21,24,33,36,\\ \qquad\qquad 75,76,77,80,95,96,104,105,\\ \gamma_{\{108,117,142\}} = \gamma_{\{112,118,141\}} = 0,\\ \gamma_{\{112,121,141\}} = \gamma_{\{50,64,116,139,142\}} = 0. \end{cases}$$

With the aid of (8.8.6.2), (8.8.6.3) and (8.8.6.4), the homomorphism h sends (8.8.6.1) to

$$\begin{aligned} &\gamma_{\{59,81,107,114,153\}}w_{1,t,2,1} + \gamma_{\{50,84,110,116,122\}}w_{1,t,2,2} + \gamma_{54}w_{1,t,2,3} + \gamma_{117}w_{1,t,2,4}\\ &\quad + \gamma_{64}w_{1,t,2,5} + \gamma_{118}w_{1,t,2,6} + a_{13}w_{1,t,2,7} + \gamma_{\{108,110,117,122,142\}}w_{1,t,2,8}\\ &\quad + \gamma_{123}w_{1,t,2,9} + \gamma_{\{112,114,118,141,153\}}w_{1,t,2,10} + a_{14}w_{1,t,2,11} + \gamma_{124}w_{1,t,2,12}\\ &\quad + \gamma_{\{118,121,122,125\}}w_{1,t,2,13} + \gamma_{\{64,85,123,152,153,154\}}w_{1,t,2,14} = 0, \end{aligned}$$

where

$$a_{13} = \gamma_{\{54,59,82,106,107,124,141,153,154\}},\ a_{14} = \gamma_{\{50,64,85,115,116,122,123,125,142\}}.$$

From the above equalities, it implies

$$\text{(8.8.6.5)}\qquad \begin{cases} a_{13} = a_{14} = 0,\ \gamma_j = 0,\ j = 54,64,117,118,123,124,\\ \gamma_{\{59,81,107,114,153\}} = \gamma_{\{50,84,110,116,122\}} = \gamma_{\{108,110,122,142\}} = 0,\\ \gamma_{\{112,114,141,153\}} = \gamma_{\{121,122,125\}} = \gamma_{\{85,152,153,154\}} = 0. \end{cases}$$

Combining (8.8.6.2), (8.8.6.3), (8.8.6.4) and (8.8.6.5), we get $\gamma_j = 0$ for all $1 \leqslant j \leqslant 154$. The proposition is proved. □

Proposition 8.8.7. *For $s \geqslant 3$, the elements $[a_{1,2,s,j}], 1 \leqslant j \leqslant 154$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+3}+2^{s+2}+2^s-3}$.*

Proof. Suppose that there is a linear relation

$$\text{(8.8.7.1)}\qquad \sum_{j=1}^{154} \gamma_j[a_{1,2,s,j}] = 0,$$

with $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.8.7.1) and we have

$$\begin{aligned}
&\gamma_1 w_{1,2,s,1} + \gamma_2 w_{1,2,s,2} + \gamma_{15} w_{1,2,s,3} + \gamma_{16} w_{1,2,s,4} + \gamma_{29} w_{1,2,s,5}\\
&\quad + \gamma_{30} w_{1,2,s,6} + \gamma_{37} w_{1,2,s,7} + \gamma_{38} w_{1,2,s,8} + \gamma_{51} w_{1,2,s,9} + \gamma_{55} w_{1,2,s,10}\\
&\quad + \gamma_{56} w_{1,2,s,11} + \gamma_{61} w_{1,2,s,12} + \gamma_{87} w_{1,2,s,13} + \gamma_{144} w_{1,2,s,14} = 0,\\
&\gamma_3 w_{1,2,s,1} + \gamma_5 w_{1,2,s,2} + \gamma_{13} w_{1,2,s,3} + \gamma_{18} w_{1,2,s,4} + \gamma_{26} w_{1,2,s,5}\\
&\quad + \gamma_{28} w_{1,2,s,6} + \gamma_{41} w_{1,2,s,7} + \gamma_{39} w_{1,2,s,8} + \gamma_{50} w_{1,2,s,9} + \gamma_{\{57,66\}} w_{1,2,s,10}\\
&\quad + \gamma_{66} w_{1,2,s,11} + \gamma_{\{59,66\}} w_{1,2,s,12} + \gamma_{91} w_{1,2,s,13} + \gamma_{\{66,91,145\}} w_{1,2,s,14} = 0,\\
&\gamma_4 w_{1,2,s,1} + \gamma_6 w_{1,2,s,2} + \gamma_{14} w_{1,2,s,3} + \gamma_{17} w_{1,2,s,4} + \gamma_{25} w_{1,2,s,5}\\
&\quad + \gamma_{27} w_{1,2,s,6} + \gamma_{42} w_{1,2,s,7} + \gamma_{43} w_{1,2,s,8} + \gamma_{49} w_{1,2,s,9} + \gamma_{\{65,88,147\}} w_{1,2,s,10}\\
&\quad + \gamma_{\{65,147\}} w_{1,2,s,11} + \gamma_{\{60,65\}} w_{1,2,s,12} + \gamma_{67} w_{1,2,s,13} + \gamma_{\{65,67\}} w_{1,2,s,14} = 0,\\
&\gamma_{19} w_{1,2,s,1} + \gamma_{32} w_{1,2,s,2} + \gamma_7 w_{1,2,s,3} + \gamma_{35} w_{1,2,s,4} + \gamma_{10} w_{1,2,s,5}\\
&\quad + \gamma_{23} w_{1,2,s,6} + \gamma_{44} w_{1,2,s,7} + \gamma_{53} w_{1,2,s,8} + \gamma_{40} w_{1,2,s,9} + \gamma_{62} w_{1,2,s,10}\\
&\quad + \gamma_{69} w_{1,2,s,11} + \gamma_{58} w_{1,2,s,12} + \gamma_{92} w_{1,2,s,13} + \gamma_{146} w_{1,2,s,14} = 0,\\
&\gamma_{20} w_{1,2,s,1} + \gamma_{31} w_{1,2,s,2} + \gamma_8 w_{1,2,s,3} + \gamma_{34} w_{1,2,s,4} + \gamma_{11} w_{1,2,s,5}\\
&\quad + \gamma_{22} w_{1,2,s,6} + \gamma_{45} w_{1,2,s,7} + \gamma_{52} w_{1,2,s,8} + \gamma_{47} w_{1,2,s,9} + \gamma_{\{63,148\}} w_{1,2,s,10}\\
&\quad + \gamma_{\{68,148\}} w_{1,2,s,11} + \gamma_{89} w_{1,2,s,12} + \gamma_{71} w_{1,2,s,13} + \gamma_{148} w_{1,2,s,14} = 0,\\
&\gamma_{33} w_{1,2,s,1} + \gamma_{21} w_{1,2,s,2} + \gamma_{36} w_{1,2,s,3} + \gamma_9 w_{1,2,s,4} + \gamma_{24} w_{1,2,s,5}\\
&\quad + \gamma_{12} w_{1,2,s,6} + \gamma_{54} w_{1,2,s,7} + \gamma_{46} w_{1,2,s,8} + \gamma_{48} w_{1,2,s,9} + \gamma_{70} w_{1,2,s,10}\\
&\quad + \gamma_{64} w_{1,2,s,11} + \gamma_{72} w_{1,2,s,12} + \gamma_{90} w_{1,2,s,13} + \gamma_{149} w_{1,2,s,14} = 0.
\end{aligned}$$

Computing from these equalities gives

$$\gamma_j = 0,\ j = 1, \ldots, 72, 87, \ldots, 92, 144, \ldots, 149. \tag{8.8.7.2}$$

With the aid of (8.8.7.2), the homomorphisms g_1, g_2, g_3, g_4 send (8.8.7.1) to

$$\begin{aligned}
&\gamma_{126} w_{1,2,s,1} + \gamma_{131} w_{1,2,s,2} + \gamma_{73} w_{1,2,s,3} + \gamma_{83} w_{1,2,s,4} + \gamma_{78} w_{1,2,s,5}\\
&\quad + \gamma_{86} w_{1,2,s,6} + \gamma_{93} w_{1,2,s,7} + \gamma_{98} w_{1,2,s,8} + \gamma_{101} w_{1,2,s,9} + \gamma_{99} w_{1,2,s,10}\\
&\quad + \gamma_{103} w_{1,2,s,11} + \gamma_{111} w_{1,2,s,12} + \gamma_{120} w_{1,2,s,13} + \gamma_{151} w_{1,2,s,14} = 0,\\
&\gamma_{127} w_{1,2,s,1} + \gamma_{130} w_{1,2,s,2} + \gamma_{74} w_{1,2,s,3} + \gamma_{82} w_{1,2,s,4} + \gamma_{79} w_{1,2,s,5}\\
&\quad + \gamma_{85} w_{1,2,s,6} + \gamma_{94} w_{1,2,s,7} + \gamma_{97} w_{1,2,s,8} + \gamma_{109} w_{1,2,s,9} + \gamma_{\{100,150\}} w_{1,2,s,10}\\
&\quad + \gamma_{\{102,150\}} w_{1,2,s,11} + \gamma_{119} w_{1,2,s,12} + \gamma_{113} w_{1,2,s,13} + \gamma_{150} w_{1,2,s,14} = 0,\\
&\gamma_{128} w_{1,2,s,1} + \gamma_{132} w_{1,2,s,2} + \gamma_{75} w_{1,2,s,3} + a_1 w_{1,2,s,4} + \gamma_{77} w_{1,2,s,5} + a_2 w_{1,2,s,6}\\
&\quad + \gamma_{95} w_{1,2,s,7} + \gamma_{106} w_{1,2,s,8} + \gamma_{108} w_{1,2,s,9} + \gamma_{\{121,152\}} w_{1,2,s,10} + \gamma_{121} w_{1,2,s,12}\\
&\quad + \gamma_{\{104,121,152\}} w_{1,2,s,11} + \gamma_{\{112,115\}} w_{1,2,s,13} + \gamma_{\{115,121,152\}} w_{1,2,s,14} = 0,
\end{aligned}$$

$$\gamma_{129}w_{1,2,s,1} + \gamma_{133}w_{1,2,s,2} + \gamma_{76}w_{1,2,s,3} + a_3w_{1,2,s,4} + \gamma_{80}w_{1,2,s,5} + a_6w_{1,2,s,6} + \gamma_{96}w_{1,2,s,7} + \gamma_{107}w_{1,2,s,8} + \gamma_{110}w_{1,2,s,9} + \gamma_{\{122,153\}}w_{1,2,s,10} + \gamma_{122}w_{1,2,s,12} + \gamma_{\{105,122,153\}}w_{1,2,s,11} + \gamma_{\{114,116\}}w_{1,2,s,13} + \gamma_{\{116,122,153\}}w_{1,2,s,14} = 0,$$

where

$$a_1 = \begin{cases} \gamma_{\{81,142\}}, & s = 3, \\ \gamma_{81}, & s \geqslant 4, \end{cases} \quad a_2 = \begin{cases} \gamma_{\{84,143\}}, & s = 3, \\ \gamma_{84}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{118,123,124,134,136,137,138,140,142,154\}}, & s = 3, \\ \gamma_{134}, & s \geqslant 4, \end{cases}$$

$$a_4 = \begin{cases} \gamma_{\{117,118,123,125,135,137,139,141,143\}}, & s = 3, \\ \gamma_{135}, & s \geqslant 4. \end{cases}$$

These equalities imply

$$\begin{cases} \gamma_j = 0,\ j = 73, \ldots, 80, 82, 83, 85, 86, 93, \ldots, 116, \\ \qquad 119, \ldots, 122, 126, \ldots, 133, 150, \ldots, 153, \\ a_1 = a_2 = a_3 = a_4 = 0. \end{cases} \tag{8.8.7.3}$$

With the aid of (8.8.7.2) and (8.8.7.3), the homomorphism h sends (8.8.7.1) to

$$a_5w_{1,2,s,1} + a_6w_{1,2,s,2} + \gamma_{136}w_{1,2,s,3} + \gamma_{117}w_{1,2,s,4} + \gamma_{137}w_{1,2,s,5} + \gamma_{118}w_{1,2,s,6} + \gamma_{138}w_{1,2,s,7} + \gamma_{139}w_{1,2,s,8} + \gamma_{123}w_{1,2,s,9} + \gamma_{140}w_{1,2,s,10} + \gamma_{141}w_{1,2,s,11} + \gamma_{124}w_{1,2,s,12} + \gamma_{125}w_{1,2,s,13} + \gamma_{154}w_{1,2,s,14} = 0,$$

where

$$a_5 = \begin{cases} \gamma_{134}, & s = 3, \\ \gamma_{142}, & s \geqslant 4, \end{cases} \quad a_6 = \begin{cases} \gamma_{135}, & s = 3, \\ \gamma_{143}, & s \geqslant 4. \end{cases}$$

From the above equalities, it implies

$$\begin{cases} a_5 = a_6 = 0, \quad \gamma_j = 0,\ j = 117, 118, 123, \\ 124, 125, 136, 137, 138, 139, 140, 141, 154. \end{cases} \tag{8.8.7.4}$$

Combining (8.8.7.2), (8.8.7.3) and (8.8.7.4), we get $\gamma_j = 0$ for all $1 \leqslant j \leqslant 154$. The proposition is proved. □

Proposition 8.8.8. *For $s \geqslant 3$ and $t \geqslant 3$, the elements $[a_{1,t,s,j}], 1 \leqslant j \leqslant 154$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+1}+2^{s+t}+2^s-3}$.*

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{154} \gamma_j[a_{1,t,s,j}] = 0, \tag{8.8.8.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.8.8.1), we obtain

$$\begin{aligned}
&\gamma_1 w_{1,t,s,1} + \gamma_2 w_{1,t,s,2} + \gamma_{15} w_{1,t,s,3} + \gamma_{16} w_{1,t,s,4} + \gamma_{29} w_{1,t,s,5}\\
&\quad + \gamma_{30} w_{1,t,s,6} + \gamma_{37} w_{1,t,s,7} + \gamma_{38} w_{1,t,s,8} + \gamma_{51} w_{1,t,s,9} + \gamma_{55} w_{1,t,s,10}\\
&\quad + \gamma_{56} w_{1,t,s,11} + \gamma_{61} w_{1,t,s,12} + \gamma_{87} w_{1,t,s,13} + \gamma_{144} w_{1,t,s,14} = 0,\\
&\gamma_3 w_{1,t,s,1} + \gamma_5 w_{1,t,s,2} + \gamma_{13} w_{1,t,s,3} + \gamma_{18} w_{1,t,s,4} + \gamma_{26} w_{1,t,s,5}\\
&\quad + \gamma_{28} w_{1,t,s,6} + \gamma_{41} w_{1,t,s,7} + \gamma_{39} w_{1,t,s,8} + \gamma_{50} w_{1,t,s,9} + \gamma_{57} w_{1,t,s,10}\\
&\quad + \gamma_{66} w_{1,t,s,11} + \gamma_{59} w_{1,t,s,12} + \gamma_{91} w_{1,t,s,13} + \gamma_{\{66,91,145\}} w_{1,t,s,14} = 0,\\
&\gamma_4 w_{1,t,s,1} + \gamma_6 w_{1,t,s,2} + \gamma_{14} w_{1,t,s,3} + \gamma_{17} w_{1,t,s,4} + \gamma_{25} w_{1,t,s,5}\\
&\quad + \gamma_{27} w_{1,t,s,6} + \gamma_{42} w_{1,t,s,7} + \gamma_{43} w_{1,t,s,8} + \gamma_{49} w_{1,t,s,9} + \gamma_{67} w_{1,t,s,10}\\
&\quad + \gamma_{65} w_{1,t,s,11} + \gamma_{60} w_{1,t,s,12} + \gamma_{\{67,147\}} w_{1,t,s,13} + \gamma_{\{65,88,147\}} w_{1,t,s,14} = 0,\\
&\gamma_{19} w_{1,t,s,1} + \gamma_{32} w_{1,t,s,2} + \gamma_7 w_{1,t,s,3} + \gamma_{35} w_{1,t,s,4} + \gamma_{10} w_{1,t,s,5}\\
&\quad + \gamma_{23} w_{1,t,s,6} + \gamma_{44} w_{1,t,s,7} + \gamma_{53} w_{1,t,s,8} + \gamma_{40} w_{1,t,s,9} + \gamma_{62} w_{1,t,s,10}\\
&\quad + \gamma_{69} w_{1,t,s,11} + \gamma_{58} w_{1,t,s,12} + \gamma_{92} w_{1,t,s,13} + \gamma_{146} w_{1,t,s,14} = 0,\\
&\gamma_{20} w_{1,t,s,1} + \gamma_{31} w_{1,t,s,2} + \gamma_8 w_{1,t,s,3} + \gamma_{34} w_{1,t,s,4} + \gamma_{11} w_{1,t,s,5}\\
&\quad + \gamma_{22} w_{1,t,s,6} + \gamma_{45} w_{1,t,s,7} + \gamma_{52} w_{1,t,s,8} + \gamma_{47} w_{1,t,s,9} + \gamma_{63} w_{1,t,s,10}\\
&\quad + \gamma_{68} w_{1,t,s,11} + \gamma_{71} w_{1,t,s,12} + \gamma_{\{71,148\}} w_{1,t,s,13} + \gamma_{\{71,89\}} w_{1,t,s,14} = 0,\\
&\gamma_{33} w_{1,t,s,1} + \gamma_{21} w_{1,t,s,2} + \gamma_{36} w_{1,t,s,3} + \gamma_9 w_{1,t,s,4} + \gamma_{24} w_{1,t,s,5}\\
&\quad + \gamma_{12} w_{1,t,s,6} + \gamma_{54} w_{1,t,s,7} + \gamma_{46} w_{1,t,s,8} + \gamma_{48} w_{1,t,s,9} + \gamma_{70} w_{1,t,s,10}\\
&\quad + \gamma_{64} w_{1,t,s,11} + \gamma_{72} w_{1,t,s,12} + \gamma_{90} w_{1,t,s,13} + \gamma_{149} w_{1,t,s,14} = 0.
\end{aligned}$$

Computing from these equalities gives

$$\gamma_j = 0,\ j = 1, \ldots, 72, 87, \ldots, 92, 144, \ldots, 149. \tag{8.8.8.2}$$

With the aid of (8.8.8.2), the homomorphisms g_1, g_2, g_3, g_4 send (8.8.8.1) to

$$\begin{aligned}
&\gamma_{126} w_{1,t,s,1} + \gamma_{131} w_{1,t,s,2} + \gamma_{73} w_{1,t,s,3} + \gamma_{83} w_{1,t,s,4} + \gamma_{78} w_{1,t,s,5}\\
&\quad + \gamma_{86} w_{1,t,s,6} + \gamma_{93} w_{1,t,s,7} + \gamma_{98} w_{1,t,s,8} + \gamma_{101} w_{1,t,s,9} + \gamma_{99} w_{1,t,s,10}\\
&\quad + \gamma_{103} w_{1,t,s,11} + \gamma_{111} w_{1,t,s,12} + \gamma_{120} w_{1,t,s,13} + \gamma_{150} w_{1,t,s,14} = 0,\\
&\gamma_{127} w_{1,t,s,1} + \gamma_{130} w_{1,t,s,2} + \gamma_{74} w_{1,t,s,3} + \gamma_{82} w_{1,t,s,4} + \gamma_{79} w_{1,t,s,5}\\
&\quad + \gamma_{85} w_{1,t,s,6} + \gamma_{94} w_{1,t,s,7} + \gamma_{97} w_{1,t,s,8} + \gamma_{109} w_{1,t,s,9} + \gamma_{100} w_{1,t,s,10}\\
&\quad + \gamma_{102} w_{1,t,s,11} + \gamma_{113} w_{1,t,s,12} + \gamma_{\{113,151\}} w_{1,t,s,13} + \gamma_{\{113,119\}} w_{1,t,s,14} = 0,\\
&\gamma_{128} w_{1,t,s,1} + \gamma_{132} w_{1,t,s,2} + \gamma_{75} w_{1,t,s,3} + a_1 w_{1,t,s,4} + \gamma_{77} w_{1,t,s,5} + a_2 w_{1,t,s,6}\\
&\quad + \gamma_{95} w_{1,t,s,7} + \gamma_{106} w_{1,t,s,8} + \gamma_{108} w_{1,t,s,9} + \gamma_{115} w_{1,t,s,10} + \gamma_{104} w_{1,t,s,11}\\
&\quad + \gamma_{112} w_{1,t,s,12} + \gamma_{\{112,115,121,152\}} w_{1,t,s,13} + \gamma_{\{112,121\}} w_{1,t,s,14} = 0,
\end{aligned}$$

$$\gamma_{129}w_{1,t,s,1} + \gamma_{133}w_{1,t,s,2} + \gamma_{76}w_{1,t,s,3} + a_3w_{1,t,s,4} + \gamma_{80}w_{1,t,s,5} + a_4w_{1,t,s,6} + \gamma_{96}w_{1,t,s,7} + \gamma_{107}w_{1,t,s,8} + \gamma_{110}w_{1,t,s,9} + \gamma_{116}w_{1,t,s,10} + \gamma_{105}w_{1,t,s,11} + \gamma_{114}w_{1,t,s,12} + \gamma_{\{114,116,122,153\}}w_{1,t,s,13} + \gamma_{\{114,122\}}w_{1,t,s,14} = 0,$$

where

$$a_1 = \begin{cases} \gamma_{\{81,142\}}, & s = 3, \\ \gamma_{81}, & s \geqslant 4, \end{cases} \quad a_2 = \begin{cases} \gamma_{\{84,143\}}, & s = 3, \\ \gamma_{84}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{118,123,124,134,136,137,138,140,142,154\}}, & s = 3, \\ \gamma_{134}, & s \geqslant 4, \end{cases}$$

$$a_4 = \begin{cases} \gamma_{\{117,118,123,125,135,137,139,141,143\}}, & s = 3, \\ \gamma_{135}, & s \geqslant 4. \end{cases}$$

These equalities imply

$$\begin{cases} \gamma_j = 0,\ j = 73, \ldots, 83, 85, 86, 93, \ldots, 116, \\ 119, \ldots, 122, 126, \ldots, 133, 150, 151, 152, 153, \\ a_1 = a_2 = a_3 = a_4 = 0. \end{cases} \tag{8.8.8.3}$$

With the aid of (8.8.8.2) and (8.8.8.3), the homomorphisms h sends (8.8.8.1) to

$$a_5w_{1,t,s,1} + a_6w_{1,t,s,2} + \gamma_{136}w_{1,t,s,3} + \gamma_{117}w_{1,t,s,4} + \gamma_{137}w_{1,t,s,5} + \gamma_{118}w_{1,t,s,6} + \gamma_{138}w_{1,t,s,7} + \gamma_{139}w_{1,t,s,8} + \gamma_{123}w_{1,t,s,9} + \gamma_{140}w_{1,t,s,10} + \gamma_{141}w_{1,t,s,11} + \gamma_{124}w_{1,t,s,12} + \gamma_{125}w_{1,t,s,13} + \gamma_{154}w_{1,t,s,14} = 0,$$

where

$$a_5 = \begin{cases} \gamma_{134}, & s = 3, \\ \gamma_{142}, & s \geqslant 4, \end{cases} \quad a_6 = \begin{cases} \gamma_{135}, & s = 3, \\ \gamma_{143}, & s \geqslant 4. \end{cases}$$

From the above equalities, we obtain

$$\begin{cases} a_5 = a_6 = 0, \quad \gamma_j = 0,\ j = 117, 118, 123, \\ 124, 125, 136, 137, 138, 139, 140, 141, 154. \end{cases} \tag{8.8.8.4}$$

Combining (8.8.8.2), (8.8.8.3) and (8.8.8.4), we get $\gamma_j = 0$ for all $1 \leqslant j \leqslant 154$. The proposition is proved. □

8.9. **The case $s \geqslant 2, t \geqslant 2$ and $u \geqslant 2$.**

According to Kameko [8], for $s \geqslant 2, t \geqslant 2, u \geqslant 2$, the dimension of the space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} P_3)_{2^{s+t+u}+2^{s+t}+2^s-3}$ is 21 with a basis given by the following classes:

$$\begin{aligned} w_{u,t,s,1} &= [2^s - 1, 2^{s+t} - 1, 2^{s+t+u} - 1], \\ w_{u,t,s,2} &= [2^s - 1, 2^{s+t+u} - 1, 2^{s+t} - 1], \\ w_{u,t,s,3} &= [2^{s+t} - 1, 2^s - 1, 2^{s+t+u} - 1], \\ w_{u,t,s,4} &= [2^{s+t} - 1, 2^{s+t+u} - 1, 2^s - 1], \end{aligned}$$

$w_{u,t,s,5} = [2^{s+t+u}-1, 2^s-1, 2^{s+t}-1]$,
$w_{u,t,s,6} = [2^{s+t+u}-1, 2^{s+t}-1, 2^s-1]$,
$w_{u,t,s,7} = [2^s-1, 2^{s+t+1}-1, 2^{s+t+u}-2^{s+t}-1]$,
$w_{u,t,s,8} = [2^{s+t+1}-1, 2^s-1, 2^{s+t+u}-2^{s+t}-1]$,
$w_{u,t,s,9} = [2^{s+t+1}-1, 2^{s+t+u}-2^{s+t}-1, 2^s-1]$,
$w_{u,t,s,10} = [2^{s+1}-1, 2^{s+t}-2^s-1, 2^{s+t+u}-1]$,
$w_{u,t,s,11} = [2^{s+1}-1, 2^{s+t+u}-1, 2^{s+t}-2^s-1]$,
$w_{u,t,s,12} = [2^{s+t+u}-1, 2^{s+1}-1, 2^{s+t}-2^s-1]$,
$w_{u,t,s,13} = [2^{s+1}-1, 2^{s+t}-1, 2^{s+t+u}-2^s-1]$,
$w_{u,t,s,14} = [2^{s+1}-1, 2^{s+t+u}-2^s-1, 2^{s+t}-1]$,
$w_{u,t,s,15} = [2^{s+t}-1, 2^{s+1}-1, 2^{s+t+u}-2^s-1]$,
$w_{u,t,s,16} = [2^{s+1}-1, 2^{s+t+1}-2^s-1, 2^{s+t+u}-2^{s+t}-1]$,
$w_{u,t,s,17} = [2^{s+1}-1, 2^{s+t+1}-1, 2^{s+t+u}-2^{s+t}-2^s-1]$,
$w_{u,t,s,18} = [2^{s+t+1}-1, 2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-1]$,
$w_{u,t,s,19} = [2^{s+2}-1, 2^{s+t+1}-2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-1]$,
$w_{u,t,s,20} = [2^{s+2}-1, 2^{s+t+u}-2^{s+1}-1, 2^{s+t}-2^s-1]$,
$w_{u,2,s,21} = [2^{s+2}-1, 2^{s+2}-1, 2^{s+u+2}-2^{s+1}-2^s-1]$, for $t=2$,
$w_{u,t,s,21} = [2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 2^{s+t+u}-2^s-1,]$, for $t \geqslant 3$.

So, we easily obtain

Proposition 8.9.1. *For any positive integer $s,t,u \geqslant 2$, we have*

$$\dim(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} Q_4)_{2^{s+t+u}+2^{s+t}+2^s-3} = 84.$$

Next, we determine $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+u}+2^{s+t}+2^s-3}$. We have

Theorem 8.9.2. *For $s,t,u \geqslant 2$, $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+u}+2^{s+t}+2^s-3}$ is an $\mathbb{F}_2$-vector space of dimension 231 with a basis consisting of all the classes represented by the monomials $a_{u,t,s,j}, 1 \leqslant j \leqslant 231$, which are determined as follows:*

For $s,t,u \geqslant 2$

1. $(1, 2^s-2, 2^{s+t}-1, 2^{s+t+u}-1)$,
2. $(1, 2^s-2, 2^{s+t+u}-1, 2^{s+t}-1)$,
3. $(1, 2^{s+t}-1, 2^s-2, 2^{s+t+u}-1)$,
4. $(1, 2^{s+t}-1, 2^{s+t+u}-1, 2^s-2)$,
5. $(1, 2^{s+t+u}-1, 2^s-2, 2^{s+t}-1)$,
6. $(1, 2^{s+t+u}-1, 2^{s+t}-1, 2^s-2)$,
7. $(2^{s+t}-1, 1, 2^s-2, 2^{s+t+u}-1)$,
8. $(2^{s+t}-1, 1, 2^{s+t+u}-1, 2^s-2)$,
9. $(2^{s+t}-1, 2^{s+t+u}-1, 1, 2^s-2)$,
10. $(2^{s+t+u}-1, 1, 2^s-2, 2^{s+t}-1)$,
11. $(2^{s+t+u}-1, 1, 2^{s+t}-1, 2^s-2)$,
12. $(2^{s+t+u}-1, 2^{s+t}-1, 1, 2^s-2)$,
13. $(1, 2^s-1, 2^{s+t}-2, 2^{s+t+u}-1)$,
14. $(1, 2^s-1, 2^{s+t+u}-1, 2^{s+t}-2)$,

15. $(1, 2^{s+t} - 2, 2^s - 1, 2^{s+t+u} - 1)$,
16. $(1, 2^{s+t} - 2, 2^{s+t+u} - 1, 2^s - 1)$,
17. $(1, 2^{s+t+u} - 1, 2^s - 1, 2^{s+t} - 2)$,
18. $(1, 2^{s+t+u} - 1, 2^{s+t} - 2, 2^s - 1)$,
19. $(2^s - 1, 1, 2^{s+t} - 2, 2^{s+t+u} - 1)$,
20. $(2^s - 1, 1, 2^{s+t+u} - 1, 2^{s+t} - 2)$,
21. $(2^s - 1, 2^{s+t+u} - 1, 1, 2^{s+t} - 2)$,
22. $(2^{s+t+u} - 1, 1, 2^s - 1, 2^{s+t} - 2)$,
23. $(2^{s+t+u} - 1, 1, 2^{s+t} - 2, 2^s - 1)$,
24. $(2^{s+t+u} - 1, 2^s - 1, 1, 2^{s+t} - 2)$,
25. $(1, 2^s - 1, 2^{s+t} - 1, 2^{s+t+u} - 2)$,
26. $(1, 2^s - 1, 2^{s+t+u} - 2, 2^{s+t} - 1)$,
27. $(1, 2^{s+t} - 1, 2^s - 1, 2^{s+t+u} - 2)$,
28. $(1, 2^{s+t} - 1, 2^{s+t+u} - 2, 2^s - 1)$,
29. $(1, 2^{s+t+u} - 2, 2^s - 1, 2^{s+t} - 1)$,
30. $(1, 2^{s+t+u} - 2, 2^{s+t} - 1, 2^s - 1)$,
31. $(2^s - 1, 1, 2^{s+t} - 1, 2^{s+t+u} - 2)$,
32. $(2^s - 1, 1, 2^{s+t+u} - 2, 2^{s+t} - 1)$,
33. $(2^s - 1, 2^{s+t} - 1, 1, 2^{s+t+u} - 2)$,
34. $(2^{s+t} - 1, 1, 2^s - 1, 2^{s+t+u} - 2)$,
35. $(2^{s+t} - 1, 1, 2^{s+t+u} - 2, 2^s - 1)$,
36. $(2^{s+t} - 1, 2^s - 1, 1, 2^{s+t+u} - 2)$,
37. $(1, 2^s - 2, 2^{s+t+1} - 1, 2^{s+t+u} - 2^{s+t} - 1)$,
38. $(1, 2^{s+t+1} - 1, 2^s - 2, 2^{s+t+u} - 2^{s+t} - 1)$,
39. $(1, 2^{s+t+1} - 1, 2^{s+t+u} - 2^{s+t} - 1, 2^s - 2)$,
40. $(2^{s+t+1} - 1, 1, 2^s - 2, 2^{s+t+u} - 2^{s+t} - 1)$,
41. $(2^{s+t+1} - 1, 1, 2^{s+t+u} - 2^{s+t} - 1, 2^s - 2)$,
42. $(2^{s+t+1} - 1, 2^{s+t+u} - 2^{s+t} - 1, 1, 2^s - 2)$,
43. $(1, 2^s - 1, 2^{s+t+1} - 2, 2^{s+t+u} - 2^{s+t} - 1)$,
44. $(1, 2^{s+t+1} - 2, 2^s - 1, 2^{s+t+u} - 2^{s+t} - 1)$,
45. $(1, 2^{s+t+1} - 2, 2^{s+t+u} - 2^{s+t} - 1, 2^s - 1)$,
46. $(2^s - 1, 1, 2^{s+t+1} - 2, 2^{s+t+u} - 2^{s+t} - 1)$,
47. $(1, 2^s - 1, 2^{s+t+1} - 1, 2^{s+t+u} - 2^{s+t} - 2)$,
48. $(1, 2^{s+t+1} - 1, 2^s - 1, 2^{s+t+u} - 2^{s+t} - 2)$,
49. $(1, 2^{s+t+1} - 1, 2^{s+t+u} - 2^{s+t} - 2, 2^s - 1)$,
50. $(2^s - 1, 1, 2^{s+t+1} - 1, 2^{s+t+u} - 2^{s+t} - 2)$,
51. $(2^s - 1, 2^{s+t+1} - 1, 1, 2^{s+t+u} - 2^{s+t} - 2)$,
52. $(2^{s+t+1} - 1, 1, 2^s - 1, 2^{s+t+u} - 2^{s+t} - 2)$,
53. $(2^{s+t+1} - 1, 1, 2^{s+t+u} - 2^{s+t} - 2, 2^s - 1)$,
54. $(2^{s+t+1} - 1, 2^s - 1, 1, 2^{s+t+u} - 2^{s+t} - 2)$,
55. $(1, 2^{s+1} - 2, 2^{s+t} - 2^s - 1, 2^{s+t+u} - 1)$,
56. $(1, 2^{s+1} - 2, 2^{s+t+u} - 1, 2^{s+t} - 2^s - 1)$,
57. $(1, 2^{s+t+u} - 1, 2^{s+1} - 2, 2^{s+t} - 2^s - 1)$,
58. $(2^{s+t+u} - 1, 1, 2^{s+1} - 2, 2^{s+t} - 2^s - 1)$,
59. $(1, 2^{s+1} - 1, 2^{s+t} - 2^s - 2, 2^{s+t+u} - 1)$,

60. $(1, 2^{s+1}-1, 2^{s+t+u}-1, 2^{s+t}-2^s-2)$,
61. $(1, 2^{s+t+u}-1, 2^{s+1}-1, 2^{s+t}-2^s-2)$,
62. $(2^{s+1}-1, 1, 2^{s+t}-2^s-2, 2^{s+t+u}-1)$,
63. $(2^{s+1}-1, 1, 2^{s+t+u}-1, 2^{s+t}-2^s-2)$,
64. $(2^{s+1}-1, 2^{s+t+u}-1, 1, 2^{s+t}-2^s-2)$,
65. $(2^{s+t+u}-1, 1, 2^{s+1}-1, 2^{s+t}-2^s-2)$,
66. $(2^{s+t+u}-1, 2^{s+1}-1, 1, 2^{s+t}-2^s-2)$,
67. $(1, 2^{s+1}-1, 2^{s+t}-2^s-1, 2^{s+t+u}-2)$,
68. $(1, 2^{s+1}-1, 2^{s+t+u}-2, 2^{s+t}-2^s-1)$,
69. $(1, 2^{s+t+u}-2, 2^{s+1}-1, 2^{s+t}-2^s-1)$,
70. $(2^{s+1}-1, 1, 2^{s+t}-2^s-1, 2^{s+t+u}-2)$,
71. $(2^{s+1}-1, 1, 2^{s+t+u}-2, 2^{s+t}-2^s-1)$,
72. $(2^{s+1}-1, 2^{s+t}-2^s-1, 1, 2^{s+t+u}-2)$,
73. $(1, 2^{s+1}-2, 2^{s+t}-1, 2^{s+t+u}-2^s-1)$,
74. $(1, 2^{s+1}-2, 2^{s+t+u}-2^s-1, 2^{s+t}-1)$,
75. $(1, 2^{s+t}-1, 2^{s+1}-2, 2^{s+t+u}-2^s-1)$,
76. $(2^{s+t}-1, 1, 2^{s+1}-2, 2^{s+t+u}-2^s-1)$,
77. $(1, 2^{s+1}-1, 2^{s+t}-2, 2^{s+t+u}-2^s-1)$,
78. $(1, 2^{s+1}-1, 2^{s+t+u}-2^s-1, 2^{s+t}-2)$,
79. $(1, 2^{s+t}-2, 2^{s+1}-1, 2^{s+t+u}-2^s-1)$,
80. $(2^{s+1}-1, 1, 2^{s+t}-2, 2^{s+t+u}-2^s-1)$,
81. $(2^{s+1}-1, 1, 2^{s+t+u}-2^s-1, 2^{s+t}-2)$,
82. $(2^{s+1}-1, 2^{s+t+u}-2^s-1, 1, 2^{s+t}-2)$,
83. $(1, 2^{s+1}-1, 2^{s+t}-1, 2^{s+t+u}-2^s-2)$,
84. $(1, 2^{s+1}-1, 2^{s+t+u}-2^s-2, 2^{s+t}-1)$,
85. $(1, 2^{s+t}-1, 2^{s+1}-1, 2^{s+t+u}-2^s-2)$,
86. $(2^{s+1}-1, 1, 2^{s+t}-1, 2^{s+t+u}-2^s-2)$,
87. $(2^{s+1}-1, 1, 2^{s+t+u}-2^s-2, 2^{s+t}-1)$,
88. $(2^{s+1}-1, 2^{s+t}-1, 1, 2^{s+t+u}-2^s-2)$,
89. $(2^{s+t}-1, 1, 2^{s+1}-1, 2^{s+t+u}-2^s-2)$,
90. $(2^{s+t}-1, 2^{s+1}-1, 1, 2^{s+t+u}-2^s-2)$,
91. $(3, 2^{s+t}-3, 2^s-2, 2^{s+t+u}-1)$,
92. $(3, 2^{s+t}-3, 2^{s+t+u}-1, 2^s-2)$,
93. $(3, 2^{s+t+u}-1, 2^{s+t}-3, 2^s-2)$,
94. $(2^{s+t+u}-1, 3, 2^{s+t}-3, 2^s-2)$,
95. $(3, 2^{s+t}-1, 2^{s+t+u}-3, 2^s-2)$,
96. $(3, 2^{s+t+u}-3, 2^s-2, 2^{s+t}-1)$,
97. $(3, 2^{s+t+u}-3, 2^{s+t}-1, 2^s-2)$,
98. $(2^{s+t}-1, 3, 2^{s+t+u}-3, 2^s-2)$,
99. $(1, 2^{s+1}-2, 2^{s+t+1}-2^s-1, 2^{s+t+u}-2^{s+t}-1)$,
100. $(1, 2^{s+1}-2, 2^{s+t+1}-1, 2^{s+t+u}-2^{s+t}-2^s-1)$,
101. $(1, 2^{s+t+1}-1, 2^{s+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
102. $(2^{s+t+1}-1, 1, 2^{s+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
103. $(1, 2^{s+1}-1, 2^{s+t+1}-2^s-2, 2^{s+t+u}-2^{s+t}-1)$,
104. $(2^{s+1}-1, 1, 2^{s+t+1}-2^s-2, 2^{s+t+u}-2^{s+t}-1)$,

105. $(1, 2^{s+1}-1, 2^{s+t+1}-2^s-1, 2^{s+t+u}-2^{s+t}-2)$,
106. $(2^{s+1}-1, 1, 2^{s+t+1}-2^s-1, 2^{s+t+u}-2^{s+t}-2)$,
107. $(2^{s+1}-1, 2^{s+t+1}-2^s-1, 1, 2^{s+t+u}-2^{s+t}-2)$,
108. $(3, 2^s-1, 2^{s+t}-3, 2^{s+t+u}-2)$,
109. $(3, 2^{s+t}-3, 2^s-1, 2^{s+t+u}-2)$,
110. $(3, 2^{s+t}-3, 2^{s+t+u}-2, 2^s-1)$,
111. $(3, 2^s-1, 2^{s+t+u}-3, 2^{s+t}-2)$,
112. $(3, 2^{s+t+u}-3, 2^s-1, 2^{s+t}-2)$,
113. $(3, 2^{s+t+u}-3, 2^{s+t}-2, 2^s-1)$,
114. $(1, 2^{s+1}-1, 2^{s+t+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
115. $(1, 2^{s+t+1}-2, 2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-1)$,
116. $(2^{s+1}-1, 1, 2^{s+t+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
117. $(1, 2^{s+1}-1, 2^{s+t+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
118. $(1, 2^{s+t+1}-1, 2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
119. $(2^{s+1}-1, 1, 2^{s+t+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
120. $(2^{s+1}-1, 2^{s+t+1}-1, 1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
121. $(2^{s+t+1}-1, 1, 2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
122. $(2^{s+t+1}-1, 2^{s+1}-1, 1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
123. $(1, 2^{s+2}-2, 2^{s+t+u}-2^{s+1}-1, 2^{s+t}-2^s-1)$,
124. $(1, 2^{s+2}-1, 2^{s+t+u}-2^{s+1}-1, 2^{s+t}-2^s-2)$,
125. $(2^{s+2}-1, 1, 2^{s+t+u}-2^{s+1}-1, 2^{s+t}-2^s-2)$,
126. $(2^{s+2}-1, 2^{s+t+u}-2^{s+1}-1, 1, 2^{s+t}-2^s-2)$,
127. $(1, 2^{s+2}-1, 2^{s+t+u}-2^{s+1}-2, 2^{s+t}-2^s-1)$,
128. $(2^{s+2}-1, 1, 2^{s+t+u}-2^{s+1}-2, 2^{s+t}-2^s-1)$,
129. $(3, 2^{s+t+1}-3, 2^s-2, 2^{s+t+u}-2^{s+t}-1)$,
130. $(3, 2^{s+t+1}-3, 2^{s+t+u}-2^{s+t}-1, 2^s-2)$,
131. $(3, 2^{s+t+1}-1, 2^{s+t+u}-2^{s+t}-3, 2^s-2)$,
132. $(2^{s+t+1}-1, 3, 2^{s+t+u}-2^{s+t}-3, 2^s-2)$,
133. $(3, 2^s-1, 2^{s+t+1}-3, 2^{s+t+u}-2^{s+t}-2)$,
134. $(3, 2^{s+t+1}-3, 2^s-1, 2^{s+t+u}-2^{s+t}-2)$,
135. $(3, 2^{s+t+1}-3, 2^{s+t+u}-2^{s+t}-2, 2^s-1)$,
136. $(3, 2^{s+1}-3, 2^{s+t}-2^s-2, 2^{s+t+u}-1)$,
137. $(3, 2^{s+1}-3, 2^{s+t+u}-1, 2^{s+t}-2^s-2)$,
138. $(3, 2^{s+t+u}-1, 2^{s+1}-3, 2^{s+t}-2^s-2)$,
139. $(2^{s+t+u}-1, 3, 2^{s+1}-3, 2^{s+t}-2^s-2)$,
140. $(3, 2^{s+1}-3, 2^{s+t}-2^s-1, 2^{s+t+u}-2)$,
141. $(3, 2^{s+1}-3, 2^{s+t+u}-2, 2^{s+t}-2^s-1)$,
142. $(3, 2^{s+t+u}-3, 2^{s+1}-2, 2^{s+t}-2^s-1)$,
143. $(1, 2^{s+2}-2, 2^{s+t+1}-2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-1)$,
144. $(1, 2^{s+2}-1, 2^{s+t+1}-2^{s+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
145. $(2^{s+2}-1, 1, 2^{s+t+1}-2^{s+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
146. $(1, 2^{s+2}-1, 2^{s+t+1}-2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
147. $(2^{s+2}-1, 1, 2^{s+t+1}-2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
148. $(2^{s+2}-1, 2^{s+t+1}-2^{s+1}-1, 1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
149. $(3, 2^{s+1}-1, 2^{s+t}-2^s-3, 2^{s+t+u}-2)$,

150. $(2^{s+1}-1, 3, 2^{s+t}-2^s-3, 2^{s+t+u}-2)$,
151. $(3, 2^{s+1}-1, 2^{s+t+u}-3, 2^{s+t}-2^s-2)$,
152. $(3, 2^{s+t+u}-3, 2^{s+1}-1, 2^{s+t}-2^s-2)$,
153. $(2^{s+1}-1, 3, 2^{s+t+u}-3, 2^{s+t}-2^s-2)$,
154. $(3, 2^{s+1}-3, 2^{s+t}-2, 2^{s+t+u}-2^s-1)$,
155. $(3, 2^{s+1}-3, 2^{s+t+u}-2^s-1, 2^{s+t}-2)$,
156. $(3, 2^{s+1}-3, 2^{s+t}-1, 2^{s+t+u}-2^s-2)$,
157. $(3, 2^{s+1}-3, 2^{s+t+u}-2^s-2, 2^{s+t}-1)$,
158. $(3, 2^{s+t}-1, 2^{s+1}-3, 2^{s+t+u}-2^s-2)$,
159. $(2^{s+t}-1, 3, 2^{s+1}-3, 2^{s+t+u}-2^s-2)$,
160. $(3, 2^{s+t}-3, 2^{s+1}-2, 2^{s+t+u}-2^s-1)$,
161. $(3, 2^{s+1}-1, 2^{s+t}-3, 2^{s+t+u}-2^s-2)$,
162. $(3, 2^{s+t}-3, 2^{s+1}-1, 2^{s+t+u}-2^s-2)$,
163. $(2^{s+1}-1, 3, 2^{s+t}-3, 2^{s+t+u}-2^s-2)$,
164. $(3, 2^{s+1}-1, 2^{s+t+u}-2^s-3, 2^{s+t}-2)$,
165. $(2^{s+1}-1, 3, 2^{s+t+u}-2^s-3, 2^{s+t}-2)$,
166. $(7, 2^{s+t}-5, 2^{s+t+u}-3, 2^s-2)$,
167. $(7, 2^{s+t+u}-5, 2^{s+t}-3, 2^s-2)$,
168. $(3, 2^{s+1}-3, 2^{s+t+1}-2^s-2, 2^{s+t+u}-2^{s+t}-1)$,
169. $(3, 2^{s+1}-3, 2^{s+t+1}-2^s-1, 2^{s+t+u}-2^{s+t}-2)$,
170. $(3, 2^{s+1}-3, 2^{s+t+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
171. $(3, 2^{s+1}-3, 2^{s+t+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
172. $(3, 2^{s+t+1}-1, 2^{s+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$,
173. $(2^{s+t+1}-1, 3, 2^{s+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$,
174. $(3, 2^{s+t+1}-3, 2^{s+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
175. $(3, 2^{s+1}-1, 2^{s+t+1}-2^s-3, 2^{s+t+u}-2^{s+t}-2)$,
176. $(2^{s+1}-1, 3, 2^{s+t+1}-2^s-3, 2^{s+t+u}-2^{s+t}-2)$,
177. $(3, 2^{s+1}-1, 2^{s+t+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$,
178. $(3, 2^{s+t+1}-3, 2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
179. $(2^{s+1}-1, 3, 2^{s+t+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$,
180. $(7, 2^{s+t+1}-5, 2^{s+t+u}-2^{s+t}-3, 2^s-2)$,
181. $(3, 2^{s+2}-3, 2^{s+t+u}-2^{s+1}-1, 2^{s+t}-2^s-2)$,
182. $(3, 2^{s+2}-3, 2^{s+t+u}-2^{s+1}-2, 2^{s+t}-2^s-1)$,
183. $(3, 2^{s+2}-1, 2^{s+t+u}-2^{s+1}-3, 2^{s+t}-2^s-2)$,
184. $(2^{s+2}-1, 3, 2^{s+t+u}-2^{s+1}-3, 2^{s+t}-2^s-2)$,
185. $(7, 2^{s+t+u}-5, 2^{s+1}-3, 2^{s+t}-2^s-2)$,
186. $(7, 2^{s+t}-5, 2^{s+1}-3, 2^{s+t+u}-2^s-2)$,
187. $(3, 2^{s+2}-3, 2^{s+t+1}-2^{s+1}-2, 2^{s+t+u}-2^{s+t}-2^s-1)$,
188. $(3, 2^{s+2}-3, 2^{s+t+1}-2^{s+1}-1, 2^{s+t+u}-2^{s+t}-2^s-2)$,
189. $(3, 2^{s+2}-1, 2^{s+t+1}-2^{s+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$,
190. $(2^{s+2}-1, 3, 2^{s+t+1}-2^{s+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$,
191. $(7, 2^{s+t+1}-5, 2^{s+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$,
192. $(7, 2^{s+2}-5, 2^{s+t+u}-2^{s+1}-3, 2^{s+t}-2^s-2)$,
193. $(7, 2^{s+2}-5, 2^{s+t+1}-2^{s+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$.

For $s = 2, t, u \geqslant 2$,

194. $(3, 3, 2^{t+2} - 4, 2^{t+u+2} - 1)$,
195. $(3, 3, 2^{t+u+2} - 1, 2^{t+2} - 4)$,
196. $(3, 2^{t+u+2} - 1, 3, 2^{t+2} - 4)$,
197. $(2^{t+u+2} - 1, 3, 3, 2^{t+2} - 4)$,
198. $(3, 3, 2^{t+2} - 1, 2^{t+u+2} - 4)$,
199. $(3, 3, 2^{t+u+2} - 4, 2^{t+2} - 1)$,
200. $(3, 2^{t+2} - 1, 3, 2^{t+u+2} - 4)$,
201. $(2^{t+2} - 1, 3, 3, 2^{t+u+2} - 4)$,
202. $(3, 7, 2^{t+2} - 5, 2^{t+u+2} - 4)$,
203. $(7, 3, 2^{t+2} - 5, 2^{t+u+2} - 4)$,
204. $(3, 3, 2^{t+3} - 4, 2^{t+u+2} - 2^{t+2} - 1)$,
205. $(3, 3, 2^{t+3} - 1, 2^{t+u+2} - 2^{t+2} - 4)$,
206. $(3, 2^{t+3} - 1, 3, 2^{t+u+2} - 2^{t+2} - 4)$,
207. $(2^{t+3} - 1, 3, 3, 2^{t+u+2} - 2^{t+2} - 4)$,
208. $(3, 7, 2^{t+u+2} - 5, 2^{t+2} - 4)$,
209. $(7, 2^{t+2} - 5, 3, 2^{t+u+2} - 4)$,
210. $(7, 2^{t+u+2} - 5, 3, 2^{t+2} - 4)$,
211. $(7, 2^{t+3} - 5, 3, 2^{t+u+2} - 2^{t+2} - 4)$,
212. $(7, 3, 2^{t+3} - 5, 2^{t+u+2} - 2^{t+2} - 4)$,
213. $(7, 7, 2^{t+2} - 8, 2^{t+u+2} - 5)$,
214. $(7, 7, 2^{t+2} - 7, 2^{t+u+2} - 6)$,
215. $(7, 7, 2^{t+u+2} - 7, 2^{t+2} - 6)$,
216. $(7, 7, 2^{t+u+2} - 8, 2^{t+2} - 5)$,
217. $(7, 7, 2^{t+3} - 8, 2^{t+u+2} - 2^{t+2} - 5)$,
218. $(7, 7, 2^{t+3} - 7, 2^{t+u+2} - 2^{t+2} - 6)$,
219. $(3, 7, 2^{t+3} - 5, 2^{t+u+2} - 2^{t+2} - 4)$,
220. $(7, 3, 2^{t+u+2} - 5, 2^{t+2} - 4)$.

For $s \geqslant 3, t, u \geqslant 2$,

194. $(3, 2^s - 3, 2^{s+t} - 2, 2^{s+t+u} - 1)$,
195. $(3, 2^s - 3, 2^{s+t+u} - 1, 2^{s+t} - 2)$,
196. $(3, 2^{s+t+u} - 1, 2^s - 3, 2^{s+t} - 2)$,
197. $(2^{s+t+u} - 1, 3, 2^s - 3, 2^{s+t} - 2)$,
198. $(3, 2^s - 3, 2^{s+t} - 1, 2^{s+t+u} - 2)$,
199. $(3, 2^s - 3, 2^{s+t+u} - 2, 2^{s+t} - 1)$,
200. $(3, 2^{s+t} - 1, 2^s - 3, 2^{s+t+u} - 2)$,
201. $(2^{s+t} - 1, 3, 2^s - 3, 2^{s+t+u} - 2)$,
202. $(2^s - 1, 3, 2^{s+t} - 3, 2^{s+t+u} - 2)$,
203. $(2^s - 1, 3, 2^{s+t+u} - 3, 2^{s+t} - 2)$,
204. $(3, 2^s - 3, 2^{s+t+1} - 2, 2^{s+t+u} - 2^{s+t} - 1)$,
205. $(3, 2^s - 3, 2^{s+t+1} - 1, 2^{s+t+u} - 2^{s+t} - 2)$,
206. $(3, 2^{s+t+1} - 1, 2^s - 3, 2^{s+t+u} - 2^{s+t} - 2)$,
207. $(2^{s+t+1} - 1, 3, 2^s - 3, 2^{s+t+u} - 2^{s+t} - 2)$,

208. $(2^s-1, 3, 2^{s+t+1}-3, 2^{s+t+u}-2^{s+t}-2)$,
209. $(7, 2^{s+t}-5, 2^s-3, 2^{s+t+u}-2)$,
210. $(7, 2^{s+t+u}-5, 2^s-3, 2^{s+t}-2)$,
211. $(7, 2^{s+t+1}-5, 2^s-3, 2^{s+t+u}-2^{s+t}-2)$,
212. $(7, 2^{s+1}-5, 2^{s+t}-2^s-3, 2^{s+t+u}-2)$,
213. $(7, 2^{s+1}-5, 2^{s+t+u}-3, 2^{s+t}-2^s-2)$,
214. $(7, 2^{s+1}-5, 2^{s+t}-3, 2^{s+t+u}-2^s-2)$,
215. $(7, 2^{s+1}-5, 2^{s+t+u}-2^s-3, 2^{s+t}-2)$,
216. $(7, 2^{s+1}-5, 2^{s+t+1}-2^s-3, 2^{s+t+u}-2^{s+t}-2)$,
217. $(7, 2^{s+1}-5, 2^{s+t+1}-3, 2^{s+t+u}-2^{s+t}-2^s-2)$.

For $s=3, t, u \geqslant 2$,

218. $(7, 7, 2^{t+3}-7, 2^{t+u+3}-2)$,
219. $(7, 7, 2^{t+u+3}-7, 2^{t+3}-2)$,
220. $(7, 7, 2^{t+4}-7, 2^{t+u+3}-2^{t+3}-2)$.

For $s \geqslant 4, t, u \geqslant 2$,

218. $(7, 2^s-5, 2^{s+t}-3, 2^{s+t+u}-2)$,
219. $(7, 2^s-5, 2^{s+t+u}-3, 2^{s+t}-2)$,
220. $(7, 2^s-5, 2^{s+t+1}-3, 2^{s+t+u}-2^{s+t}-2)$.

For $s \geqslant 2, t=2, u \geqslant 2$,

221. $(1, 2^{s+2}-2, 2^{s+2}-1, 2^{s+u+2}-2^{s+1}-2^s-1)$,
222. $(1, 2^{s+2}-1, 2^{s+2}-2, 2^{s+u+2}-2^{s+1}-2^s-1)$,
223. $(2^{s+2}-1, 1, 2^{s+2}-2, 2^{s+u+2}-2^{s+1}-2^s-1)$,
224. $(1, 2^{s+2}-1, 2^{s+2}-1, 2^{s+u+2}-2^{s+1}-2^s-2)$,
225. $(2^{s+2}-1, 1, 2^{s+2}-1, 2^{s+u+2}-2^{s+1}-2^s-2)$,
226. $(2^{s+2}-1, 2^{s+2}-1, 1, 2^{s+u+2}-2^{s+1}-2^s-2)$,
227. $(3, 2^{s+2}-3, 2^{s+2}-2, 2^{s+u+2}-2^{s+1}-2^s-1)$,
228. $(3, 2^{s+2}-3, 2^{s+2}-1, 2^{s+u+2}-2^{s+1}-2^s-2)$,
229. $(3, 2^{s+2}-1, 2^{s+2}-3, 2^{s+u+2}-2^{s+1}-2^s-2)$,
230. $(2^{s+2}-1, 3, 2^{s+2}-3, 2^{s+u+2}-2^{s+1}-2^s-2)$,
231. $(7, 2^{s+2}-5, 2^{s+2}-3, 2^{s+u+2}-2^{s+1}-2^s-2)$.

For $s \geqslant 2, t \geqslant 3, u \geqslant 2$,

221. $(1, 2^{s+2}-2, 2^{s+t}-2^{s+1}-1, 2^{s+t+u}-2^s-1)$,
222. $(1, 2^{s+2}-1, 2^{s+t}-2^{s+1}-2, 2^{s+t+u}-2^s-1)$,
223. $(2^{s+2}-1, 1, 2^{s+t}-2^{s+1}-2, 2^{s+t+u}-2^s-1)$,
224. $(1, 2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 2^{s+t+u}-2^s-2)$,
225. $(2^{s+2}-1, 1, 2^{s+t}-2^{s+1}-1, 2^{s+t+u}-2^s-2)$,
226. $(2^{s+2}-1, 2^{s+t}-2^{s+1}-1, 1, 2^{s+t+u}-2^s-2)$,
227. $(3, 2^{s+2}-3, 2^{s+t}-2^{s+1}-2, 2^{s+t+u}-2^s-1)$,
228. $(3, 2^{s+2}-3, 2^{s+t}-2^{s+1}-1, 2^{s+t+u}-2^s-2)$,
229. $(3, 2^{s+2}-1, 2^{s+t}-2^{s+1}-3, 2^{s+t+u}-2^s-2)$,
230. $(2^{s+2}-1, 3, 2^{s+t}-2^{s+1}-3, 2^{s+t+u}-2^s-2)$,
231. $(7, 2^{s+2}-5, 2^{s+t}-2^{s+1}-3, 2^{s+t+u}-2^s-2)$.

We prove this theorem by combining some propositions.

Proposition 8.9.3. *For $n = 2^{s+t+u} + 2^{s+t} + 2^s - 3$ with $s, t, u \geqslant 2$, the $\mathbb{F}_2$-vector space $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_n$ is generated by the elements listed in Theorem 8.9.2.*

We need the following lemma.

Lemma 8.9.4. *The following matrix is strictly inadmissible*

$$\begin{pmatrix} 1 & 1 & 1 & 0\\ 1 & 1 & 1 & 0\\ 1 & 1 & 0 & 0\\ 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Proof. The monomials corresponding to the above matrix is (15,15,19,32). By a direct calculation, we get

$$\begin{aligned}(15,15,19,32) &= Sq^1(15,15,19,31) + Sq^2(15,15,18,31)\\ &\quad + Sq^4\big((4,23,19,31) + (15,12,19,31) + (7,20,19,31) + (5,23,18,31)\\ &\quad + (15,13,18,31) + (7,21,18,31)\big) + Sq^8\big((8,15,19,31) + (4,27,11,31)\\ &\quad + (11,12,19,31) + (7,24,11,31) + (9,15,18,31) + (5,27,10,31)\\ &\quad + (11,13,18,31) + (7,25,10,31)\big) + (8,15,19,39) + (4,35,11,31)\\ &\quad + (4,27,11,39) + (15,12,19,35) + (11,12,19,39) + (7,32,11,31)\\ &\quad + (7,24,11,39) + (15,15,18,33) + (9,15,18,39)\\ &\quad + (5,35,10,31) + (5,27,10,39) + (15,13,18,35) + (11,13,18,39)\\ &\quad + (7,33,10,31) + (7,25,10,39) \mod \mathcal{L}_4(3;3;2;2;1;1).\end{aligned}$$

The lemma is proved. □

Using the results in Section 7, the lemmas in Sections 3, 5, 6, Lemmas 8.3.4, 8.3.5, 8.4.4, 8.8.4, 8.9.4 and Theorem 2.5, we get Proposition 8.9.3.

Now, we prove that the classes listed in Theorem 8.9.2 are linearly independent.

Proposition 8.9.5. *For $u \geqslant 2$, the elements $[a_{u,2,2,j}], 1 \leqslant j \leqslant 231$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{u+4}+17}$.*

Proof. Suppose that there is a linear relation

$$\sum_{j=1}^{231} \gamma_j [a_{u,2,2,j}] = 0, \tag{8.9.5.1}$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.9.5.1), we obtain

$$\begin{aligned}
&\gamma_1 w_{u,2,2,1} + \gamma_2 w_{u,2,2,2} + \gamma_{15} w_{u,2,2,3} + \gamma_{16} w_{u,2,2,4} + \gamma_{29} w_{u,2,2,5} \\
&\quad + \gamma_{30} w_{u,2,2,6} + \gamma_{37} w_{u,2,2,7} + \gamma_{44} w_{u,2,2,8} + \gamma_{45} w_{u,2,2,9} \\
&\quad + \gamma_{55} w_{u,2,2,10} + \gamma_{56} w_{u,2,2,11} + \gamma_{69} w_{u,2,2,12} + \gamma_{73} w_{u,2,2,13} \\
&\quad + \gamma_{74} w_{u,2,2,14} + \gamma_{79} w_{u,2,2,15} + \gamma_{99} w_{u,2,2,16} + \gamma_{100} w_{u,2,2,17} \\
&\quad + \gamma_{115} w_{u,2,2,18} + \gamma_{143} w_{u,2,2,19} + \gamma_{123} w_{u,2,2,20} + \gamma_{221} w_{u,2,2,21} = 0, \\
&\gamma_3 w_{u,2,2,1} + \gamma_5 w_{u,2,2,2} + \gamma_{\{13,194\}} w_{u,2,2,3} + \gamma_{18} w_{u,2,2,4} + \gamma_{\{26,199\}} w_{u,2,2,5} \\
&\quad + \gamma_{28} w_{u,2,2,6} + \gamma_{38} w_{u,2,2,7} + \gamma_{\{43,204\}} w_{u,2,2,8} + \gamma_{49} w_{u,2,2,9} + \gamma_{59} w_{u,2,2,10} \\
&\quad + \gamma_{57} w_{u,2,2,11} + \gamma_{\{68,216\}} w_{u,2,2,12} + \gamma_{\{75,84,103\}} w_{u,2,2,13} + \gamma_{84} w_{u,2,2,14} \\
&\quad + \gamma_{\{77,84,103,213\}} w_{u,2,2,15} + \gamma_{103} w_{u,2,2,16} + \gamma_{101} w_{u,2,2,17} + \gamma_{\{114,217\}} w_{u,2,2,18} \\
&\quad + \gamma_{144} w_{u,2,2,19} + \gamma_{127} w_{u,2,2,20} + \gamma_{\{84,103,127,144,222\}} w_{u,2,2,21} = 0, \\
&\gamma_4 w_{u,2,2,1} + \gamma_6 w_{u,2,2,2} + \gamma_{\{14,195\}} w_{u,2,2,3} + \gamma_{\{17,196\}} w_{u,2,2,4} + \gamma_{\{25,198\}} w_{u,2,2,5} \\
&\quad + \gamma_{\{27,200\}} w_{u,2,2,6} + \gamma_{39} w_{u,2,2,7} + \gamma_{\{47,205\}} w_{u,2,2,8} + \gamma_{\{48,206\}} w_{u,2,2,9} \\
&\quad + \gamma_{60} w_{u,2,2,10} + \gamma_{61} w_{u,2,2,11} + \gamma_{\{67,202\}} w_{u,2,2,12} \\
&\quad + \gamma_{\{83,117,118,124,224\}} w_{u,2,2,13} + \gamma_{\{83,224\}} w_{u,2,2,14} + \gamma_{\{78,83,117,208\}} w_{u,2,2,15} \\
&\quad + \gamma_{117} w_{u,2,2,16} + \gamma_{118} w_{u,2,2,17} + \gamma_{\{105,219\}} w_{u,2,2,18} + \gamma_{146} w_{u,2,2,19} \\
&\quad + \gamma_{\{85,118\}} w_{u,2,2,20} + \gamma_{\{83,85,117,118,146\}} w_{u,2,2,21} = 0, \\
&\gamma_{\{19,91,136,194\}} w_{u,2,2,1} + \gamma_{\{32,96,157,199\}} w_{u,2,2,2} + \gamma_7 w_{u,2,2,3} + \gamma_{35} w_{u,2,2,4} \\
&\quad + \gamma_{10} w_{u,2,2,5} + \gamma_{23} w_{u,2,2,6} + \gamma_{\{46,129,168,204\}} w_{u,2,2,7} + \gamma_{40} w_{u,2,2,8} \\
&\quad + \gamma_{53} w_{u,2,2,9} + \gamma_{62} w_{u,2,2,10} + \gamma_{\{71,216\}} w_{u,2,2,11} + \gamma_{58} w_{u,2,2,12} + \gamma_{87} w_{u,2,2,14} \\
&\quad + \gamma_{\{80,213\}} w_{u,2,2,13} + \gamma_{76} w_{u,2,2,15} + \gamma_{104} w_{u,2,2,16} + \gamma_{\{116,217\}} w_{u,2,2,17} \\
&\quad + \gamma_{110} w_{u,2,2,18} + \gamma_{145} w_{u,2,2,19} + \gamma_{128} w_{u,2,2,20} + \gamma_{223} w_{u,2,2,21} = 0, \\
&\gamma_{\{20,92,137,195\}} w_{u,2,2,1} + \gamma_{\{31,97,156,198,228\}} w_{u,2,2,2} + \gamma_8 w_{u,2,2,3} \\
&\quad + \gamma_{\{34,201\}} w_{u,2,2,4} + \gamma_{11} w_{u,2,2,5} + \gamma_{\{22,197\}} w_{u,2,2,6} + \gamma_{\{50,130,171,205\}} w_{u,2,2,7} \\
&\quad + \gamma_{41} w_{u,2,2,8} + \gamma_{\{52,207\}} w_{u,2,2,9} + \gamma_{63} w_{u,2,2,10} + \gamma_{\{70,203\}} w_{u,2,2,11} \\
&\quad + \gamma_{65} w_{u,2,2,12} + \gamma_{\{81,220,225\}} w_{u,2,2,13} + \gamma_{\{86,225\}} w_{u,2,2,14} \\
&\quad + \gamma_{\{121,125\}} w_{u,2,2,15} + \gamma_{119} w_{u,2,2,16} + \gamma_{\{106,212\}} w_{u,2,2,17} + \gamma_{121} w_{u,2,2,18} \\
&\quad + \gamma_{147} w_{u,2,2,19} + \gamma_{\{89,121\}} w_{u,2,2,20} + \gamma_{225} w_{u,2,2,21} = 0, \\
&\gamma_{\{33,95,158,183,189,200,229\}} w_{u,2,2,1} + \gamma_{\{21,93,138,196\}} w_{u,2,2,2} \\
&\quad + \gamma_{\{36,98,159,184,190,201,230\}} w_{u,2,2,3} + \gamma_9 w_{u,2,2,4} + \gamma_{\{24,94,139,197\}} w_{u,2,2,5} \\
&\quad + \gamma_{12} w_{u,2,2,6} + \gamma_{\{51,131,172,206\}} w_{u,2,2,7} + \gamma_{\{54,132,173,207\}} w_{u,2,2,8} \\
&\quad + \gamma_{42} w_{u,2,2,9} + \gamma_{\{72,166,186,192,193,209,231\}} w_{u,2,2,10} + \gamma_{64} w_{u,2,2,11} \\
&\quad + \gamma_{66} w_{u,2,2,12} + \gamma_{88} w_{u,2,2,13} + \gamma_{\{82,167,185,210\}} w_{u,2,2,14} + \gamma_{90} w_{u,2,2,15} \\
&\quad + \gamma_{\{107,180,191,211\}} w_{u,2,2,16} + \gamma_{120} w_{u,2,2,17} + \gamma_{122} w_{u,2,2,18} \\
&\quad + \gamma_{148} w_{u,2,2,19} + \gamma_{126} w_{u,2,2,20} + \gamma_{226} w_{u,2,2,21} = 0.
\end{aligned}$$

Computing directly from the above equalities, we obtain

$$(8.9.5.2)\quad \begin{cases} \gamma_j = 0,\ j = 1, \ldots, 12, 15, 16, 18, 23, 28, 29, 30, 35,\\ 37, \ldots, 42, 44, 45, 49, 53, 55, \ldots, 66, 69, 73, 74, 75,\\ 76, 79, 83, \ldots, 90, 99, 100, 101, 103, 104, 110, 115,\\ 117, \ldots, 128, 143, 144, \ldots, 148, 221, \ldots, 226,\\ \gamma_{\{13,194\}} = \gamma_{\{26,199\}} = \gamma_{\{43,204\}} = \gamma_{\{68,216\}} = \gamma_{\{77,213\}} = 0,\\ \gamma_{\{114,217\}} = \gamma_{\{14,195\}} = \gamma_{\{17,196\}} = \gamma_{\{25,198\}} = \gamma_{\{27,200\}} = 0,\\ \gamma_{\{47,205\}} = \gamma_{\{48,206\}} = \gamma_{\{78,208\}} = \gamma_{\{105,219\}} = \gamma_{\{71,216\}} = 0,\\ \gamma_{\{67,202\}} = \gamma_{\{19,91,136,194\}} = \gamma_{\{80,213\}} = \gamma_{\{32,96,157,199\}} = 0,\\ \gamma_{\{46,129,168,204\}} = \gamma_{\{116,217\}} = \gamma_{\{34,201\}} = \gamma_{\{20,92,137,195\}} = 0,\\ \gamma_{\{22,197\}} = \gamma_{\{31,97,156,198,228\}} = \gamma_{\{52,207\}} = \gamma_{\{70,203\}} = 0,\\ \gamma_{\{50,130,171,205\}} = \gamma_{\{81,220\}} = \gamma_{\{33,95,158,183,189,200,229\}} = 0,\\ \gamma_{\{106,212\}} = \gamma_{\{21,93,138,196\}} = \gamma_{\{36,98,159,184,190,201,230\}} = 0,\\ \gamma_{\{24,94,139,197\}} = \gamma_{\{51,131,172,206\}} = \gamma_{\{54,132,173,207\}} = 0,\\ \gamma_{\{72,166,186,192,193,209,231\}} = 0,\\ \gamma_{\{82,167,185,210\}} = \gamma_{\{107,180,191,211\}} = 0. \end{cases}$$

With the aid of (8.9.5.2), the homomorphisms g_1, g_2 send (8.9.5.1) to

$$\begin{aligned} &\gamma_{19}w_{u,2,2,1} + \gamma_{32}w_{u,2,2,2} + \gamma_{91}w_{u,2,2,3} + \gamma_{109}w_{u,2,2,4} + \gamma_{96}w_{u,2,2,5}\\ &\quad + \gamma_{113}w_{u,2,2,6} + \gamma_{46}w_{u,2,2,7} + \gamma_{129}w_{u,2,2,8} + \gamma_{135}w_{u,2,2,9} + \gamma_{136}w_{u,2,2,10}\\ &\quad + \gamma_{\{68,141\}}w_{u,2,2,11} + \gamma_{142}w_{u,2,2,12} + \gamma_{\{77,154\}}w_{u,2,2,13} + \gamma_{157}w_{u,2,2,14}\\ &\quad + \gamma_{160}w_{u,2,2,15} + \gamma_{168}w_{u,2,2,16} + \gamma_{\{114,170\}}w_{u,2,2,17} + \gamma_{174}w_{u,2,2,18}\\ &\quad + \gamma_{187}w_{u,2,2,19} + \gamma_{182}w_{u,2,2,20} + \gamma_{227}w_{u,2,2,21} = 0,\\ &\gamma_{20}w_{u,2,2,1} + \gamma_{31}w_{u,2,2,2} + \gamma_{92}w_{u,2,2,3} + \gamma_{\{108,209\}}w_{u,2,2,4} + \gamma_{97}w_{u,2,2,5}\\ &\quad + \gamma_{\{112,210\}}w_{u,2,2,6} + \gamma_{50}w_{u,2,2,7} + \gamma_{130}w_{u,2,2,8} + \gamma_{\{134,211\}}w_{u,2,2,9}\\ &\quad + \gamma_{137}w_{u,2,2,10} + \gamma_{140}w_{u,2,2,11} + \gamma_{152}w_{u,2,2,12} + \gamma_{\{155,228\}}w_{u,2,2,13}\\ &\quad + \gamma_{\{156,228\}}w_{u,2,2,14} + \gamma_{\{178,181\}}w_{u,2,2,15} + \gamma_{171}w_{u,2,2,16} + \gamma_{169}w_{u,2,2,17}\\ &\quad + \gamma_{178}w_{u,2,2,18} + \gamma_{188}w_{u,2,2,19} + \gamma_{\{162,178\}}w_{u,2,2,20} + \gamma_{228}w_{u,2,2,21} = 0. \end{aligned}$$

These equalities imply

$$(8.9.5.3)\quad \begin{cases} \gamma_j = 0,\ j = 19, 20, 31, 32, 46, 50, 91, 92, 96, 97, 109, 113,\\ 129, 130, 135, 136, 137, 140, 142, 152, 155, 156, 157, 160,\\ 162, 168, 169, 171, 174, 178, 181, 182, 187, 188, 227, 228,\\ \gamma_{\{68,141\}} = \gamma_{\{77,154\}} = \gamma_{\{114,170\}} = \gamma_{\{108,209\}} = 0,\\ \gamma_{\{112,210\}} = \gamma_{\{134,211\}} = 0. \end{cases}$$

With the aid of (8.9.5.2) and (8.9.5.3), the homomorphisms g_3, g_4 send (8.9.5.1) to

$$\begin{aligned}
&\gamma_{21}w_{u,2,2,1} + \gamma_{33}w_{u,2,2,2} + \gamma_{93}w_{u,2,2,3} + a_1w_{u,2,2,4} + \gamma_{95}w_{u,2,2,5}\\
&\quad + a_2w_{u,2,2,6} + \gamma_{51}w_{u,2,2,7} + \gamma_{131}w_{u,2,2,8} + a_3w_{u,2,2,9} + \gamma_{138}w_{u,2,2,10}\\
&\quad + a_4w_{u,2,2,11} + \gamma_{\{151,166,215\}}w_{u,2,2,12} + a_5w_{u,2,2,13} + \gamma_{172}w_{u,2,2,16}\\
&\quad + \gamma_{\{158,183,189,229\}}w_{u,2,2,14} + \gamma_{\{167,177,183,218\}}w_{u,2,2,15} + a_6w_{u,2,2,17}\\
&\quad + \gamma_{\{177,180,218\}}w_{u,2,2,18} + \gamma_{189}w_{u,2,2,19} + a_7w_{u,2,2,20} + a_8w_{u,2,2,21} = 0,\\
&\gamma_{24}w_{u,2,2,1} + \gamma_{36}w_{u,2,2,2} + \gamma_{94}w_{u,2,2,3} + a_9w_{u,2,2,4} + \gamma_{98}w_{u,2,2,5}\\
&\quad + a_{10}w_{u,2,2,6} + \gamma_{54}w_{u,2,2,7} + \gamma_{132}w_{u,2,2,8} + a_{11}w_{u,2,2,9} + \gamma_{139}w_{u,2,2,10}\\
&\quad + a_{12}w_{u,2,2,11} + a_{13}w_{u,2,2,12} + a_{14}w_{u,2,2,13} + \gamma_{\{159,184,190,230\}}w_{u,2,2,14}\\
&\quad + a_{15}w_{u,2,2,15} + \gamma_{173}w_{u,2,2,16} + \gamma_{\{107,114,176,211,218\}}w_{u,2,2,17} + \gamma_{190}w_{u,2,2,19}\\
&\quad + \gamma_{\{179,180,191,193,218\}}w_{u,2,2,18} + a_{16}w_{u,2,2,20} + a_{17}w_{u,2,2,21} = 0,
\end{aligned}$$

where

$$\begin{aligned}
a_1 &= \gamma_{\{22,24,34,36,52,54,70,94,102,139,150,159,163,173,230,\}},\\
a_2 &= \gamma_{\{22,24,81,94,98,111,132,139,153,165,184\}},\\
a_3 &= \gamma_{\{52,54,106,132,133,173,176,179,190\}}, \quad a_4 = \gamma_{\{72,77,149,186,209,214\}},\\
a_5 &= \gamma_{\{82,114,175,183,185,189,193,210,218,229\}},\\
a_6 &= \gamma_{\{107,114,175,191,193,211,218\}},\\
a_7 &= \gamma_{\{68,114,161,164,175,177,192,193,214,215,231\}},\\
a_8 &= \gamma_{\{68,114,164,175,183,189,192,193,215,218,229\}},\\
a_9 &= \gamma_{\{13,17,21,25,27,33,48,51,67,93,102,108,112,134,138,149,154,158,161,172,229\}},\\
a_{10} &= \gamma_{\{14,17,21,26,78,93,95,111,112,131,138,141,151,164,183\}},\\
a_{11} &= \gamma_{\{43,47,48,51,105,131,133,134,170,172,175,177,189\}},\\
a_{12} &= \gamma_{\{72,77,82,107,150,209,210,211,214\}}, \quad a_{13} = \gamma_{\{153,166,167,180,185,192,215\}},\\
a_{14} &= \gamma_{\{107,114,176,184,190,211,218,230\}}, \quad a_{15} = \gamma_{\{179,180,184,191,193,218\}},\\
a_{16} &= \gamma_{\{68,82,107,114,163,165,167,176,179,180,185,186,193,210,211,214,215,231\}},\\
a_{17} &= \gamma_{\{68,82,107,114,165,176,184,190,210,211,215,218,230\}}.
\end{aligned}$$

From the above equalities, we obtain

$$\text{(8.9.5.4)} \qquad \begin{cases}
a_i = 0, \ i = 1, 2, \ldots, 17,\\
\gamma_j = 0, \ j = 21, 24, 33, 36, 51, 54, 93, 94,\\
95, 98, 131, 132, 138, 139, 172, 173, 189, 190,\\
\gamma_{\{151,166,215\}} = \gamma_{\{158,183,229\}} = 0,\\
\gamma_{\{167,177,183,218\}} = \gamma_{\{177,180,218\}} = 0.
\end{cases}$$

With the aid of (8.9.5.2), (8.9.5.3) and (8.9.5.4), the homomorphism h sends (8.9.5.1) to

$$\begin{aligned}
&a_{18}w_{u,2,2,1} + \gamma_{\{68,111,153,165,184\}}w_{u,2,2,2} + \gamma_{72}w_{u,2,2,3} + \gamma_{166}w_{u,2,2,4}\\
&+ \gamma_{82}w_{u,2,2,5} + \gamma_{167}w_{u,2,2,6} + \gamma_{\{114,133,176,179\}}w_{u,2,2,7} + \gamma_{107}w_{u,2,2,8}\\
&+ \gamma_{180}w_{u,2,2,9} + a_{19}w_{u,2,2,10} + \gamma_{\{151,153,166,184,215\}}w_{u,2,2,11} + \gamma_{185}w_{u,2,2,12}\\
&+ a_{20}w_{u,2,2,13} + a_{21}w_{u,2,2,14} + a_{22}w_{u,2,2,15} + a_{23}w_{u,2,2,16}\\
&+ \gamma_{\{177,179,180,218\}}w_{u,2,2,17} + \gamma_{191}w_{u,2,2,18} + \gamma_{193}w_{u,2,2,19}\\
&+ \gamma_{\{167,180,183,184,192\}}w_{u,2,2,20} + a_{24}w_{u,2,2,21} = 0,
\end{aligned}$$

where

$$\begin{aligned}
a_{18} &= \gamma_{\{77,102,150,159,163,230\}},\\
a_{19} &= \gamma_{\{72,77,108,149,150,159,186,214\}},\\
a_{20} &= \gamma_{\{82,107,112,134,159,161,163,180,185,191,214,231\}},\\
a_{21} &= \gamma_{\{68,82,112,164,165,184,185,192,215\}},\\
a_{22} &= \gamma_{\{82,107,112,134,158,159,185,186,191\}},\\
a_{23} &= \gamma_{\{107,114,134,175,176,191,193,218\}},\\
a_{24} &= \gamma_{\{82,107,112,134,167,180,185,191,229,230\}}.
\end{aligned}$$

From the above equalities, it implies

$$\text{(8.9.5.5)}\qquad \begin{cases}
a_i = 0,\ i = 18, 19, \ldots, 24,\\
\gamma_j = 0,\ j = 72, 82, 107, 166, 167, 180, 185, 191, 193,\\
\gamma_{\{68,111,153,165,184\}} = 0,\\
\gamma_{\{114,133,176,179\}} = 0,\\
\gamma_{\{151,153,184,215\}} = 0,\\
\gamma_{\{177,179,218\}} = \gamma_{\{183,184,192\}} = 0.
\end{cases}$$

Combining (8.9.5.2), (8.9.5.3), (8.9.5.4) and (8.9.5.5), we get $\gamma_j = 0$ for any j. The proposition is proved. □

Proposition 8.9.6. *For $u \geqslant 2$ and $t \geqslant 3$, the elements $[a_{u,t,2,j}], 1 \leqslant j \leqslant 231$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{t+u+2}+2^{t+2}+1}$.*

Proof. Suppose that there is a linear relation

$$\text{(8.9.6.1)}\qquad \sum_{j=1}^{231} \gamma_j [a_{u,t,2,j}] = 0,$$

with $\gamma_j \in \mathbb{F}_2$.

Applying the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.9.6.1), we get

$$\begin{aligned}
&\gamma_1 w_{u,t,2,1} + \gamma_2 w_{u,t,2,2} + \gamma_{15} w_{u,t,2,3} + \gamma_{16} w_{u,t,2,4} + \gamma_{29} w_{u,t,2,5} \\
&\quad + \gamma_{30} w_{u,t,2,6} + \gamma_{37} w_{u,t,2,7} + \gamma_{44} w_{u,t,2,8} + \gamma_{45} w_{u,t,2,9} + \gamma_{55} w_{u,t,2,10} \\
&\quad + \gamma_{56} w_{u,t,2,11} + \gamma_{69} w_{u,t,2,12} + \gamma_{73} w_{u,t,2,13} + \gamma_{74} w_{u,t,2,14} \\
&\quad + \gamma_{79} w_{u,t,2,15} + \gamma_{99} w_{u,t,2,16} + \gamma_{100} w_{u,t,2,17} + \gamma_{115} w_{u,t,2,18} \\
&\quad + \gamma_{143} w_{u,t,2,19} + \gamma_{123} w_{u,t,2,20} + \gamma_{221} w_{u,t,2,21} = 0, \\
&\gamma_3 w_{u,t,2,1} + \gamma_5 w_{u,t,2,2} + \gamma_{\{13,194\}} w_{u,t,2,3} + \gamma_{18} w_{u,t,2,4} + \gamma_{\{26,199\}} w_{u,t,2,5} \\
&\quad + \gamma_{28} w_{u,t,2,6} + \gamma_{38} w_{u,t,2,7} + \gamma_{\{43,204\}} w_{u,t,2,8} + \gamma_{49} w_{u,t,2,9} + \gamma_{59} w_{u,t,2,10} \\
&\quad + \gamma_{57} w_{u,t,2,11} + \gamma_{\{68,216\}} w_{u,t,2,12} + \gamma_{75} w_{u,t,2,13} + \gamma_{84} w_{u,t,2,14} \\
&\quad + \gamma_{\{77,213\}} w_{u,t,2,15} + \gamma_{103} w_{u,t,2,16} + \gamma_{101} w_{u,t,2,17} + \gamma_{\{114,217\}} w_{u,t,2,18} \\
&\quad + \gamma_{144} w_{u,t,2,19} + \gamma_{127} w_{u,t,2,20} + \gamma_{\{84,103,127,144,222\}} w_{u,t,2,21} = 0, \\
&\gamma_4 w_{u,t,2,1} + \gamma_6 w_{u,t,2,2} + \gamma_{\{14,195\}} w_{u,t,2,3} + \gamma_{\{17,196\}} w_{u,t,2,4} \\
&\quad + \gamma_{\{25,198\}} w_{u,t,2,5} + \gamma_{\{27,200\}} w_{u,t,2,6} + \gamma_{39} w_{u,t,2,7} + \gamma_{\{47,205\}} w_{u,t,2,8} \\
&\quad + \gamma_{\{48,206\}} w_{u,t,2,9} + \gamma_{60} w_{u,t,2,10} + \gamma_{61} w_{u,t,2,11} + \gamma_{\{67,202\}} w_{u,t,2,12} \\
&\quad + \gamma_{85} w_{u,t,2,13} + \gamma_{83} w_{u,t,2,14} + \gamma_{\{78,208\}} w_{u,t,2,15} + \gamma_{117} w_{u,t,2,16} \\
&\quad + \gamma_{118} w_{u,t,2,17} + \gamma_{\{105,219\}} w_{u,t,2,18} + \gamma_{146} w_{u,t,2,19} \\
&\quad + \gamma_{\{85,118,224\}} w_{u,t,2,20} + \gamma_{\{83,117,124,146,224\}} w_{u,t,2,21} = 0, \\
&\gamma_{\{19,91,136,194\}} w_{u,t,2,1} + \gamma_{\{32,96,157,199\}} w_{u,t,2,2} + \gamma_7 w_{u,t,2,3} + \gamma_{35} w_{u,t,2,4} \\
&\quad + \gamma_{10} w_{u,t,2,5} + \gamma_{23} w_{u,t,2,6} + \gamma_{\{46,129,168,204\}} w_{u,t,2,7} + \gamma_{40} w_{u,t,2,8} \\
&\quad + \gamma_{53} w_{u,t,2,9} + \gamma_{62} w_{u,t,2,10} + \gamma_{\{71,216\}} w_{u,t,2,11} + \gamma_{58} w_{u,t,2,12} + \gamma_{87} w_{u,t,2,14} \\
&\quad + \gamma_{\{80,213\}} w_{u,t,2,13} + \gamma_{76} w_{u,t,2,15} + \gamma_{104} w_{u,t,2,16} + \gamma_{\{116,217\}} w_{u,t,2,17} \\
&\quad + \gamma_{102} w_{u,t,2,18} + \gamma_{145} w_{u,t,2,19} + \gamma_{128} w_{u,t,2,20} + \gamma_{223} w_{u,t,2,21} = 0, \\
&\gamma_{\{20,92,137,195\}} w_{u,t,2,1} + \gamma_{\{31,97,156,198\}} w_{u,t,2,2} + \gamma_8 w_{u,t,2,3} + \gamma_{\{34,201\}} w_{u,t,2,4} \\
&\quad + \gamma_{11} w_{u,t,2,5} + \gamma_{\{22,197\}} w_{u,t,2,6} + \gamma_{\{50,130,171,205\}} w_{u,t,2,7} + \gamma_{41} w_{u,t,2,8} \\
&\quad + \gamma_{\{52,207\}} w_{u,t,2,9} + \gamma_{63} w_{u,t,2,10} + \gamma_{\{70,203\}} w_{u,t,2,11} + \gamma_{65} w_{u,t,2,12} \\
&\quad + \gamma_{\{81,220\}} w_{u,t,2,13} + \gamma_{86} w_{u,t,2,14} + \gamma_{89} w_{u,t,2,15} + \gamma_{119} w_{u,t,2,16} \\
&\quad + \gamma_{\{106,212\}} w_{u,t,2,17} + \gamma_{121} w_{u,t,2,18} + \gamma_{147} w_{u,t,2,19} \\
&\quad + \gamma_{\{89,121,225\}} w_{u,t,2,20} + \gamma_{\{89,121,125\}} w_{u,t,2,21} = 0, \\
&\gamma_{\{33,95,158,200\}} w_{u,t,2,1} + \gamma_{\{21,93,138,196\}} w_{u,t,2,2} + \gamma_{\{36,98,159,201\}} w_{u,t,2,3} \\
&\quad + \gamma_9 w_{u,t,2,4} + \gamma_{\{24,94,139,197\}} w_{u,t,2,5} + \gamma_{12} w_{u,t,2,6} + \gamma_{\{51,131,172,206\}} w_{u,t,2,7} \\
&\quad + \gamma_{\{54,132,173,207\}} w_{u,t,2,8} + \gamma_{42} w_{u,t,2,9} + \gamma_{\{72,166,186,209\}} w_{u,t,2,10} + \gamma_{64} w_{u,t,2,11} \\
&\quad + \gamma_{66} w_{u,t,2,12} + \gamma_{88} w_{u,t,2,13} + \gamma_{\{82,167,185,210\}} w_{u,t,2,14} + \gamma_{90} w_{u,t,2,15} \\
&\quad + \gamma_{\{107,180,191,211\}} w_{u,t,2,16} + \gamma_{120} w_{u,t,2,17} + \gamma_{122} w_{u,t,2,18} \\
&\quad + \gamma_{148} w_{u,t,2,19} + \gamma_{126} w_{u,t,2,20} + \gamma_{226} w_{u,t,2,21} = 0.
\end{aligned}$$

Computing directly from the above equalities, we obtain

$$(8.9.6.4)\quad \begin{cases} \gamma_j = 0,\ 1, \ldots, 12, 15, 16, 18, 23, 28, 29, 30, 35,\\ 37, \ldots, 42, 44, 45, 49, 53, 55, \ldots, 66, 69, 73,\\ 74, 75, 76, 79, 83, \ldots, 90, 99, \ldots, 104, 115,\\ 117, \ldots, 128, 143, 144, \ldots, 148, 221, \ldots, 226,\\ \gamma_{\{13,194\}} = \gamma_{\{26,199\}} = \gamma_{\{43,204\}} = \gamma_{\{68,216\}} = \gamma_{\{77,213\}} = 0,\\ \gamma_{\{114,217\}} = \gamma_{\{14,195\}} = \gamma_{\{17,196\}} = \gamma_{\{25,198\}} = \gamma_{\{27,200\}} = 0,\\ \gamma_{\{47,205\}} = \gamma_{\{48,206\}} = \gamma_{\{67,202\}} = \gamma_{\{78,208\}} = \gamma_{\{71,216\}} = 0,\\ \gamma_{\{105,219\}} = \gamma_{\{19,91,136,194\}} = \gamma_{\{32,96,157,199\}} = \gamma_{\{80,213\}} = 0,\\ \gamma_{\{46,129,168,204\}} = \gamma_{\{116,217\}} = \gamma_{\{20,92,137,195\}} = \gamma_{\{34,201\}} = 0,\\ \gamma_{\{31,97,156,198\}} = \gamma_{\{22,197\}} = \gamma_{\{50,130,171,205\}} = \gamma_{\{52,207\}} = 0,\\ \gamma_{\{70,203\}} = \gamma_{\{81,220\}} = \gamma_{\{106,212\}} = \gamma_{\{33,95,158,200\}} = 0,\\ \gamma_{\{21,93,138,196\}} = \gamma_{\{36,98,159,201\}} = \gamma_{\{24,94,139,197\}} = 0,\\ \gamma_{\{51,131,172,206\}} = \gamma_{\{54,132,173,207\}} = \gamma_{\{72,166,186,209\}} = 0,\\ \gamma_{\{82,167,185,210\}} = \gamma_{\{107,180,191,211\}} = 0. \end{cases}$$

With the aid of (8.9.6.2), the homomorphisms g_1, g_2 send (8.9.6.1) to

$$\begin{aligned} &\gamma_{19}w_{u,t,2,1} + \gamma_{32}w_{u,t,2,2} + \gamma_{91}w_{u,t,2,3} + \gamma_{110}w_{u,t,2,4} + \gamma_{96}w_{u,t,2,5}\\ &\quad + \gamma_{113}w_{u,t,2,6} + \gamma_{46}w_{u,t,2,7} + \gamma_{129}w_{u,t,2,8} + \gamma_{135}w_{u,t,2,9} + \gamma_{136}w_{u,t,2,10}\\ &\quad + \gamma_{68,141}w_{u,t,2,11} + \gamma_{142}w_{u,t,2,12} + \gamma_{\{77,154\}}w_{u,t,2,13} + \gamma_{157}w_{u,t,2,14}\\ &\quad + \gamma_{160}w_{u,t,2,15} + \gamma_{168}w_{u,t,2,16} + \gamma_{\{114,170\}}w_{u,t,2,17} + \gamma_{174}w_{u,t,2,18}\\ &\quad + \gamma_{187}w_{u,t,2,19} + \gamma_{182}w_{u,t,2,20} + \gamma_{227}w_{u,t,2,21} = 0,\\ &\gamma_{20}w_{u,t,2,1} + \gamma_{31}w_{u,t,2,2} + \gamma_{92}w_{u,t,2,3} + \gamma_{\{109,209\}}w_{u,t,2,4} + \gamma_{97}w_{u,t,2,5}\\ &\quad + \gamma_{\{112,210\}}w_{u,t,2,6} + \gamma_{50}w_{u,t,2,7} + \gamma_{130}w_{u,t,2,8} + \gamma_{\{134,211\}}w_{u,t,2,9}\\ &\quad + \gamma_{137}w_{u,t,2,10} + \gamma_{140}w_{u,t,2,11} + \gamma_{152}w_{u,t,2,12} + \gamma_{155}w_{u,t,2,13} + \gamma_{156}w_{u,t,2,14}\\ &\quad + \gamma_{162}w_{u,t,2,15} + \gamma_{171}w_{u,t,2,16} + \gamma_{169}w_{u,t,2,17} + \gamma_{178}w_{u,t,2,18}\\ &\quad + \gamma_{188}w_{u,t,2,19} + \gamma_{\{162,178,228\}}w_{u,t,2,20} + \gamma_{\{162,178,181\}}w_{u,t,2,21} = 0. \end{aligned}$$

These equalities imply

$$(8.9.6.3)\quad \begin{cases} \gamma_j = 0,\ j = 19, 20, 31, 32, 46, 50, 91, 92, 96, 97, 110, 113,\\ 129, 130, 135, 136, 137, 140, 142, 152, 155, 156, 157, 160,\\ 162, 168, 169, 171, 174, 178, 181, 182, 187, 188, 227, 228,\\ \gamma_{\{68,141\}} = \gamma_{\{77,154\}} = \gamma_{\{114,170\}} = 0,\\ \gamma_{\{109,209\}} = \gamma_{\{112,210\}} = \gamma_{\{134,211\}} = 0. \end{cases}$$

With the aid of (8.9.6.2) and (8.9.6.3), the homomorphisms g_3, g_4 send (8.9.6.1) to

$$\begin{aligned}
&\gamma_{21}w_{u,t,2,1} + \gamma_{33}w_{u,t,2,2} + \gamma_{93}w_{u,t,2,3} + a_1w_{u,t,2,4} + \gamma_{95}w_{u,t,2,5}\\
&\quad + a_2w_{u,t,2,6} + \gamma_{51}w_{u,t,2,7} + \gamma_{131}w_{u,t,2,8} + a_3w_{u,t,2,9} + \gamma_{138}w_{u,t,2,10}\\
&\quad + a_4w_{u,t,2,11} + \gamma_{\{151,166,215\}}w_{u,t,2,12} + a_5w_{u,t,2,13} + \gamma_{158}w_{u,t,2,14}\\
&\quad + \gamma_{\{161,167,214\}}w_{u,t,2,15} + \gamma_{172}w_{u,t,2,16} + a_6w_{u,t,2,17} + \gamma_{\{177,180,218\}}w_{u,t,2,18}\\
&\quad + \gamma_{189}w_{u,t,2,19} + a_7w_{u,t,2,20} + \gamma_{\{161,177,183,214,218\}}w_{u,t,2,21} = 0,\\
&\gamma_{24}w_{u,t,2,1} + \gamma_{36}w_{u,t,2,2} + \gamma_{94}w_{u,t,2,3} + a_8w_{u,t,2,4} + \gamma_{98}w_{u,t,2,5}\\
&\quad + a_9w_{u,t,2,6} + \gamma_{54}w_{u,t,2,7} + \gamma_{132}w_{u,t,2,8} + a_{10}w_{u,t,2,9} + \gamma_{139}w_{u,t,2,10}\\
&\quad + a_{11}w_{u,t,2,11} + a_{12}w_{u,t,2,12} + \gamma_{\{68,82,165,210,215\}}w_{u,t,2,13} + \gamma_{159}w_{u,t,2,14}\\
&\quad + a_{13}w_{u,t,2,15} + \gamma_{173}w_{u,t,2,16} + \gamma_{\{107,114,176,211,218\}}w_{u,t,2,17}\\
&\quad + \gamma_{\{179,180,191,193,218\}}w_{u,t,2,18} + \gamma_{190}w_{u,t,2,19} + a_{14}w_{u,t,2,20} + a_{15}w_{u,t,2,21} = 0,
\end{aligned}$$

where

$$\begin{aligned}
a_1 &= \gamma_{\{22,24,34,36,52,54,70,94,108,139,150,159,163,173,230\}},\\
a_2 &= \gamma_{\{22,24,81,94,98,111,132,139,153,165,184\}}, \quad a_3 = \gamma_{\{52,54,106,132,133,173,176,179,190\}},\\
a_4 &= \gamma_{\{72,77,149,186,209,214,231\}}, \quad a_5 = \gamma_{\{68,82,164,185,192,210,215\}},\\
a_6 &= \gamma_{\{107,114,175,191,193,211,218\}},\\
a_7 &= \gamma_{\{68,114,161,164,175,177,183,189,192,193,214,215,229,\}},\\
a_8 &= \gamma_{\{13,17,21,25,27,33,48,51,67,93,108,109,112,134,138,149,154,158,161,172,229\}},\\
a_9 &= \gamma_{\{14,17,21,26,78,93,95,111,112,131,138,141,151,164,183\}},\\
a_{10} &= \gamma_{\{43,47,48,51,105,131,133,134,170,172,175,177,189\}},\\
a_{11} &= \gamma_{\{72,77,82,107,150,209,210,211,214\}},\\
a_{12} &= \gamma_{\{153,166,167,180,185,192,215\}}, \quad a_{13} = \gamma_{\{163,167,185,186,191,214,231\}},\\
a_{14} &= \gamma_{\{68,82,107,114,163,165,167,176,179,180,184,185,186,190,193,210,211,214,215,230,231\}},\\
a_{15} &= \gamma_{\{163,167,179,180,184,185,186,193,214,218,231\}}.
\end{aligned}$$

From the above equalities, we obtain

$$\text{(8.9.6.4)} \qquad \begin{cases}
a_i = 0, \ i = 1, 2, \ldots, 15,\\
\gamma_j = 0, \ j = 21, 24, 33, 36, 51, 54, 93, 94, 95, 98,\\
131, 132, 138, 139, 158, 159, 172, 173, 189, 190,\\
\gamma_{\{151,166,215\}} = \gamma_{\{161,167,214\}} = \gamma_{\{161,177,183,214,218\}} = 0,\\
\gamma_{\{161,167,214\}} = \gamma_{\{177,180,218\}} = \gamma_{\{68,82,165,210,215\}} = 0,\\
\gamma_{\{107,114,176,211,218\}} = \gamma_{\{179,180,191,193,218\}} = 0.
\end{cases}$$

With the aid of (8.9.6.2), (8.9.6.3) and (8.9.6.4), the homomorphism h sends (8.9.6.1) to

$$\begin{aligned}
&\gamma_{\{77,108,150,163,230\}}w_{u,t,2,1} + \gamma_{\{68,111,153,165,184\}}w_{u,t,2,2} + \gamma_{72}w_{u,t,2,3}\\
&+ \gamma_{166}w_{u,t,2,4} + \gamma_{82}w_{u,t,2,5} + \gamma_{167}w_{u,t,2,6} + \gamma_{\{114,133,176,179\}}w_{u,t,2,7}\\
&+ \gamma_{107}w_{u,t,2,8} + \gamma_{180}w_{u,t,2,9} + a_{16}w_{u,t,2,10} + \gamma_{\{151,153,166,184,215\}}w_{u,t,2,11}\\
&+ \gamma_{185}w_{u,t,2,12} + \gamma_{\{161,163,167,214,230\}}w_{u,t,2,13} + a_{17}w_{u,t,2,14} + \gamma_{186}w_{u,t,2,15}\\
&+ a_{18}w_{u,t,2,16} + \gamma_{\{177,179,180,218\}}w_{u,t,2,17} + \gamma_{191}w_{u,t,2,18}\\
&+ \gamma_{193}w_{u,t,2,19} + \gamma_{\{167,180,183,184,192\}}w_{u,t,2,20} + a_{19}w_{u,t,2,21} = 0,
\end{aligned}$$

where

$$\begin{aligned}
&a_{16} = \gamma_{\{72,77,109,149,150,186,214,230,231\}}, \quad a_{17} = \gamma_{\{68,82,112,164,165,184,185,192,215\}},\\
&a_{18} = \gamma_{\{107,114,134,175,176,191,193,218\}}, \quad a_{19} = \gamma_{82,107,112,134,185,191,229,230,231}.
\end{aligned}$$

From the above equalities, it implies

$$\text{(8.9.6.5)} \qquad \begin{cases}
a_i = 0, \ i = 16, 17, 18, 19,\\
\gamma_j = 0, \ j = 72, 82, 107, 166, 167, 180, 185, 191, 193,\\
\gamma_{\{77,108,150,163,230\}} = \gamma_{\{68,111,153,165,184\}} = 0,\\
\gamma_{\{114,133,176,179\}} = \gamma_{\{151,153,184,215\}} = 0,\\
\gamma_{\{161,163,214,230\}} = \gamma_{\{177,179,218\}} = \gamma_{\{183,184,192\}} = 0.
\end{cases}$$

Combining (8.9.6.2), (8.9.6.3), (8.9.6.4) and (8.9.6.5), we get $\gamma_j = 0$ for all j. The proposition is proved. □

Proposition 8.9.7. *For $s \geqslant 3$ and $u \geqslant 2$, the elements $[a_{u,2,s,j}], 1 \leqslant j \leqslant 231$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+u+2}+2^{s+2}+2^s-3}$.*

Proof. Suppose that there is a linear relation

$$\text{(8.9.7.1)} \qquad \sum_{j=1}^{231} \gamma_j [a_{u,2,s,j}] = 0,$$

with $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.9.7.1) and we obtain

$$\begin{aligned}
&\gamma_1 w_{u,2,s,1} + \gamma_2 w_{u,2,s,2} + \gamma_{15} w_{u,2,s,3} + \gamma_{16} w_{u,2,s,4} + \gamma_{29} w_{u,2,s,5}\\
&+ \gamma_{30} w_{u,2,s,6} + \gamma_{37} w_{u,2,s,7} + \gamma_{44} w_{u,2,s,8} + \gamma_{45} w_{u,2,s,9}\\
&+ \gamma_{55} w_{u,2,s,10} + \gamma_{56} w_{u,2,s,11} + \gamma_{69} w_{u,2,s,12} + \gamma_{73} w_{u,2,s,13}\\
&+ \gamma_{74} w_{u,2,s,14} + \gamma_{79} w_{u,2,s,15} + \gamma_{99} w_{u,2,s,16} + \gamma_{100} w_{u,2,s,17}\\
&+ \gamma_{115} w_{u,2,s,18} + \gamma_{143} w_{u,2,s,19} + \gamma_{123} w_{u,2,s,20} + \gamma_{221} w_{u,2,s,21} = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_3 w_{u,2,s,1} + \gamma_5 w_{u,2,s,2} + \gamma_{13} w_{u,2,s,3} + \gamma_{18} w_{u,2,s,4} + \gamma_{26} w_{u,2,s,5}\\
&\quad + \gamma_{28} w_{u,2,s,6} + \gamma_{38} w_{u,2,s,7} + \gamma_{43} w_{u,2,s,8} + \gamma_{49} w_{u,2,s,9} + \gamma_{59} w_{u,2,s,10}\\
&\quad + \gamma_{57} w_{u,2,s,11} + \gamma_{68} w_{u,2,s,12} + \gamma_{\{75,84,103\}} w_{u,2,s,13} + \gamma_{84} w_{u,2,s,14}\\
&\quad + \gamma_{\{77,84,103\}} w_{u,2,s,15} + \gamma_{103} w_{u,2,s,16} + \gamma_{101} w_{u,2,s,17} + \gamma_{114} w_{u,2,s,18}\\
&\quad + \gamma_{144} w_{u,2,s,19} + \gamma_{127} w_{u,2,s,20} + \gamma_{\{84,103,127,144,222\}} w_{u,2,s,21} = 0,\\
&\gamma_4 w_{u,2,s,1} + \gamma_6 w_{u,2,s,2} + \gamma_{14} w_{u,2,s,3} + \gamma_{17} w_{u,2,s,4} + \gamma_{25} w_{u,2,s,5}\\
&\quad + \gamma_{27} w_{u,2,s,6} + \gamma_{39} w_{u,2,s,7} + \gamma_{47} w_{u,2,s,8} + \gamma_{48} w_{u,2,s,9} + \gamma_{60} w_{u,2,s,10}\\
&\quad + \gamma_{61} w_{u,2,s,11} + \gamma_{67} w_{u,2,s,12} + \gamma_{\{83,117,118,124,224\}} w_{u,2,s,13} + \gamma_{\{83,224\}} w_{u,2,s,14}\\
&\quad + \gamma_{\{78,83,117\}} w_{u,2,s,15} + \gamma_{117} w_{u,2,s,16} + \gamma_{118} w_{u,2,s,17} + \gamma_{105} w_{u,2,s,18}\\
&\quad + \gamma_{146} w_{u,2,s,19} + \gamma_{\{85,118\}} w_{u,2,s,20} + \gamma_{\{83,85,117,118,146\}} w_{u,2,s,21} = 0,\\
&\gamma_{19} w_{u,2,s,1} + \gamma_{32} w_{u,2,s,2} + \gamma_7 w_{u,2,s,3} + \gamma_{35} w_{u,2,s,4} + \gamma_{10} w_{u,2,s,5}\\
&\quad + \gamma_{23} w_{u,2,s,6} + \gamma_{46} w_{u,2,s,7} + \gamma_{40} w_{u,2,s,8} + \gamma_{53} w_{u,2,s,9}\\
&\quad + \gamma_{62} w_{u,2,s,10} + \gamma_{71} w_{u,2,s,11} + \gamma_{58} w_{u,2,s,12} + \gamma_{80} w_{u,2,s,13}\\
&\quad + \gamma_{87} w_{u,2,s,14} + \gamma_{76} w_{u,2,s,15} + \gamma_{104} w_{u,2,s,16} + \gamma_{116} w_{u,2,s,17}\\
&\quad + \gamma_{102} w_{u,2,s,18} + \gamma_{145} w_{u,2,s,19} + \gamma_{128} w_{u,2,s,20} + \gamma_{223} w_{u,2,s,21} = 0,\\
&\gamma_{20} w_{u,2,s,1} + \gamma_{31} w_{u,2,s,2} + \gamma_8 w_{u,2,s,3} + \gamma_{34} w_{u,2,s,4} + \gamma_{11} w_{u,2,s,5}\\
&\quad + \gamma_{22} w_{u,2,s,6} + \gamma_{50} w_{u,2,s,7} + \gamma_{41} w_{u,2,s,8} + \gamma_{52} w_{u,2,s,9}\\
&\quad + \gamma_{63} w_{u,2,s,10} + \gamma_{70} w_{u,2,s,11} + \gamma_{65} w_{u,2,s,12} + \gamma_{\{81,225\}} w_{u,2,s,13}\\
&\quad + \gamma_{\{86,225\}} w_{u,2,s,14} + \gamma_{\{121,125\}} w_{u,2,s,15} + \gamma_{119} w_{u,2,s,16} + \gamma_{106} w_{u,2,s,17}\\
&\quad + \gamma_{121} w_{u,2,s,18} + \gamma_{147} w_{u,2,s,19} + \gamma_{\{89,121\}} w_{u,2,s,20} + \gamma_{225} w_{u,2,s,21} = 0,\\
&\gamma_{33} w_{u,2,s,1} + \gamma_{21} w_{u,2,s,2} + \gamma_{36} w_{u,2,s,3} + \gamma_9 w_{u,2,s,4} + \gamma_{24} w_{u,2,s,5}\\
&\quad + \gamma_{12} w_{u,2,s,6} + \gamma_{51} w_{u,2,s,7} + \gamma_{54} w_{u,2,s,8} + \gamma_{42} w_{u,2,s,9}\\
&\quad + \gamma_{72} w_{u,2,s,10} + \gamma_{64} w_{u,2,s,11} + \gamma_{66} w_{u,2,s,12} + \gamma_{88} w_{u,2,s,13}\\
&\quad + \gamma_{82} w_{u,2,s,14} + \gamma_{90} w_{u,2,s,15} + \gamma_{107} w_{u,2,s,16} + \gamma_{120} w_{u,2,s,17}\\
&\quad + \gamma_{122} w_{u,2,s,18} + \gamma_{148} w_{u,2,s,19} + \gamma_{126} w_{u,2,s,20} + \gamma_{226} w_{u,2,s,21} = 0.
\end{aligned}$$

Computing directly from the above equalities, we obtain

$$\begin{aligned}
(8.9.7.4)\qquad &\gamma_j = 0,\ j = 1, \dots, 90, 99, \dots, 107,\\
&\qquad\qquad 114, \dots, 128, 143, 144, \dots, 148, 221 \dots, 226.
\end{aligned}$$

With the aid of (8.9.7.2), the homomorphisms g_1, g_2, g_3, g_4 send (8.9.7.1) to

$$\begin{aligned}
&\gamma_{194} w_{u,2,s,1} + \gamma_{199} w_{u,2,s,2} + \gamma_{91} w_{u,2,s,3} + \gamma_{110} w_{u,2,s,4} + \gamma_{96} w_{u,2,s,5}\\
&\quad + \gamma_{113} w_{u,2,s,6} + \gamma_{204} w_{u,2,s,7} + \gamma_{129} w_{u,2,s,8} + \gamma_{135} w_{u,2,s,9}\\
&\quad + \gamma_{136} w_{u,2,s,10} + \gamma_{141} w_{u,2,s,11} + \gamma_{142} w_{u,2,s,12} + \gamma_{154} w_{u,2,s,13}\\
&\quad + \gamma_{157} w_{u,2,s,14} + \gamma_{160} w_{u,2,s,15} + \gamma_{168} w_{u,2,s,16} + \gamma_{170} w_{u,2,s,17}\\
&\quad + \gamma_{174} w_{u,2,s,18} + \gamma_{187} w_{u,2,s,19} + \gamma_{182} w_{u,2,s,20} + \gamma_{227} w_{u,2,s,21} = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_{195}w_{u,2,s,1} + \gamma_{198}w_{u,2,s,2} + \gamma_{92}w_{u,2,s,3} + \gamma_{109}w_{u,2,s,4} + \gamma_{97}w_{u,2,s,5}\\
&\quad + \gamma_{112}w_{u,2,s,6} + \gamma_{205}w_{u,2,s,7} + \gamma_{130}w_{u,2,s,8} + \gamma_{134}w_{u,2,s,9} + \gamma_{137}w_{u,2,s,10}\\
&\quad + \gamma_{140}w_{u,2,s,11} + \gamma_{152}w_{u,2,s,12} + \gamma_{\{155,228\}}w_{u,2,s,13} + \gamma_{\{156,228\}}w_{u,2,s,14}\\
&\quad + \gamma_{\{178,181\}}w_{u,2,s,15} + \gamma_{171}w_{u,2,s,16} + \gamma_{169}w_{u,2,s,17} + \gamma_{178}w_{u,2,s,18}\\
&\quad + \gamma_{188}w_{u,2,s,19} + \gamma_{\{162,178\}}w_{u,2,s,20} + \gamma_{228}w_{u,2,s,21} = 0,\\
&\gamma_{196}w_{u,2,s,1} + \gamma_{200}w_{u,2,s,2} + \gamma_{93}w_{u,2,s,3} + a_1w_{u,2,s,4} + \gamma_{95}w_{u,2,s,5} + a_2w_{u,2,s,6}\\
&\quad + \gamma_{206}w_{u,2,s,7} + \gamma_{131}w_{u,2,s,8} + a_3w_{u,2,s,9} + \gamma_{138}w_{u,2,s,10} + \gamma_{149}w_{u,2,s,11}\\
&\quad + \gamma_{151}w_{u,2,s,12} + \gamma_{\{175,183,189,229\}}w_{u,2,s,13} + \gamma_{\{158,183,189,229\}}w_{u,2,s,14}\\
&\quad + \gamma_{\{177,183\}}w_{u,2,s,15} + \gamma_{172}w_{u,2,s,16} + \gamma_{175}w_{u,2,s,17} + \gamma_{177}w_{u,2,s,18}\\
&\quad + \gamma_{189}w_{u,2,s,19} + \gamma_{\{161,164,175,177\}}w_{u,2,s,20} + \gamma_{\{164,175,183,189,229\}}w_{u,2,s,21} = 0,\\
&\gamma_{197}w_{u,2,s,1} + \gamma_{201}w_{u,2,s,2} + \gamma_{94}w_{u,2,s,3} + a_4w_{u,2,s,4} + \gamma_{98}w_{u,2,s,5} + a_5w_{u,2,s,6}\\
&\quad + \gamma_{207}w_{u,2,s,7} + \gamma_{132}w_{u,2,s,8} + a_6w_{u,2,s,9} + \gamma_{139}w_{u,2,s,10} + \gamma_{150}w_{u,2,s,11}\\
&\quad + \gamma_{153}w_{u,2,s,12} + \gamma_{\{176,184,190,230\}}w_{u,2,s,13} + \gamma_{\{159,184,190,230\}}w_{u,2,s,14}\\
&\quad + \gamma_{\{179,184\}}w_{u,2,s,15} + \gamma_{173}w_{u,2,s,16} + \gamma_{176}w_{u,2,s,17} + \gamma_{179}w_{u,2,s,18}\\
&\quad + \gamma_{190}w_{u,2,s,19} + \gamma_{\{163,165,176,179\}}w_{u,2,s,20} + \gamma_{\{165,176,184,190,230\}}w_{u,2,s,21} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases} \gamma_{\{108,218\}}, & s = 3,\\ \gamma_{108}, & s \geqslant 4, \end{cases} \quad a_2 = \begin{cases} \gamma_{\{111,219\}}, & s = 3,\\ \gamma_{111}, & s \geqslant 4, \end{cases}$$

$$a_3 = \begin{cases} \gamma_{\{133,220\}}, & s = 3,\\ \gamma_{133}, & s \geqslant 4, \end{cases}$$

$$a_4 = \begin{cases} \gamma_{\{167,185,186,191,202,209,210,211,212,214,218,231\}}, & s = 3,\\ \gamma_{202}, & s \geqslant 4, \end{cases}$$

$$a_5 = \begin{cases} \gamma_{\{166,167,180,185,192,203,210,213,215,219\}}, & s = 3,\\ \gamma_{203}, & s \geqslant 4, \end{cases}$$

$$a_6 = \begin{cases} \gamma_{\{180,191,193,208,211,216,217,220\}}, & s = 3,\\ \gamma_{208}, & s \geqslant 4. \end{cases}$$

These equalities imply

$$\text{(8.9.7.3)} \quad \begin{cases} a_i = 0,\ i = 1,2,3,4,5,6,\\ \gamma_j = 0,\ j = 91,\dots,98,109,110,112,113,129,130,131,\\ 132,134,\dots,141,142,149,150,\dots,165,168,\dots,179,\\ 181,182,183,184,187,188,189,190,194,\dots,201,\\ 204,205,206,207,227,228,229,230. \end{cases}$$

With the aid of (8.9.7.2) and (8.9.7.3), the homomorphism h sends (8.9.7.1) to

$$\begin{aligned}
&a_7w_{u,2,s,1} + a_8w_{u,2,s,2} + \gamma_{209}w_{u,2,s,3} + \gamma_{166}w_{u,2,s,4} + \gamma_{210}w_{u,2,s,5}\\
&\quad + \gamma_{167}w_{u,2,s,6} + a_9w_{u,2,s,7} + \gamma_{211}w_{u,2,s,8} + \gamma_{180}w_{u,2,s,9}\\
&\quad + \gamma_{212}w_{u,2,s,10} + \gamma_{213}w_{u,2,s,11} + \gamma_{185}w_{u,2,s,12} + \gamma_{214}w_{u,2,s,13}\\
&\quad + \gamma_{215}w_{u,2,s,14} + \gamma_{186}w_{u,2,s,15} + \gamma_{216}w_{u,2,s,16} + \gamma_{217}w_{u,2,s,17}\\
&\quad + \gamma_{191}w_{u,2,s,18} + \gamma_{193}w_{u,2,s,19} + \gamma_{192}w_{u,2,s,20} + \gamma_{231}w_{u,2,s,21} = 0,
\end{aligned}$$

where

$$a_7 = \begin{cases}\gamma_{202}, & s = 3,\\ \gamma_{218}, & s \geqslant 4,\end{cases}\quad a_8 = \begin{cases}\gamma_{203}, & s = 3,\\ \gamma_{219}, & s \geqslant 4,\end{cases}\quad a_9 = \begin{cases}\gamma_{208}, & s = 3,\\ \gamma_{220}, & s \geqslant 4.\end{cases}$$

From the above equalities, it implies

$$\text{(8.9.7.4)}\quad \begin{cases} a_7 = a_8 = a_9 = 0,\ \gamma_j = 0,\ j = 166, 167, 180, 185, 186, 191,\\ 192, 193, 209, 210, 211, 212, 213, 214, 215, 216, 217, 231.\end{cases}$$

Combining (8.9.7.2), (8.9.7.3) and (8.9.7.4), we get $\gamma_j = 0$ for all j. The proposition is proved. □

Proposition 8.9.8. *For $s \geqslant 3$, $t \geqslant 3$ and $u \geqslant 2$, the elements $[a_{u,t,s,j}]$, $1 \leqslant j \leqslant 231$, are linearly independent in $(\mathbb{F}_2 \underset{\mathcal{A}}{\otimes} R_4)_{2^{s+t+u}+2^{s+t}+2^s-3}$.*

Proof. Suppose that there is a linear relation

$$\text{(8.9.8.1)}\qquad \sum_{j=1}^{231} \gamma_j[a_{u,t,s,j}] = 0,$$

with $\gamma_j \in \mathbb{F}_2$.

Apply the homomorphisms $f_1, f_2, \ldots, f_6$ to the relation (8.9.8.1) and we obtain

$$\begin{aligned}
&\gamma_1w_{u,t,s,1} + \gamma_2w_{u,t,s,2} + \gamma_{15}w_{u,t,s,3} + \gamma_{16}w_{u,t,s,4} + \gamma_{29}w_{u,t,s,5}\\
&\quad + \gamma_{30}w_{u,t,s,6} + \gamma_{37}w_{u,t,s,7} + \gamma_{44}w_{u,t,s,8} + \gamma_{45}w_{u,t,s,9}\\
&\quad + \gamma_{55}w_{u,t,s,10} + \gamma_{56}w_{u,t,s,11} + \gamma_{69}w_{u,t,s,12} + \gamma_{73}w_{u,t,s,13}\\
&\quad + \gamma_{74}w_{u,t,s,14} + \gamma_{79}w_{u,t,s,15} + \gamma_{99}w_{u,t,s,16} + \gamma_{100}w_{u,t,s,17}\\
&\quad + \gamma_{115}w_{u,t,s,18} + \gamma_{143}w_{u,t,s,19} + \gamma_{123}w_{u,t,s,20} + \gamma_{221}w_{u,t,s,21} = 0,\\
&\gamma_3w_{u,t,s,1} + \gamma_5w_{u,t,s,2} + \gamma_{13}w_{u,t,s,3} + \gamma_{18}w_{u,t,s,4} + \gamma_{26}w_{u,t,s,5}\\
&\quad + \gamma_{28}w_{u,t,s,6} + \gamma_{38}w_{u,t,s,7} + \gamma_{43}w_{u,t,s,8} + \gamma_{49}w_{u,t,s,9} + \gamma_{59}w_{u,t,s,10}\\
&\quad + \gamma_{57}w_{u,t,s,11} + \gamma_{68}w_{u,t,s,12} + \gamma_{75}w_{u,t,s,13} + \gamma_{84}w_{u,t,s,14}\\
&\quad + \gamma_{77}w_{u,t,s,15} + \gamma_{103}w_{u,t,s,16} + \gamma_{101}w_{u,t,s,17} + \gamma_{114}w_{u,t,s,18}\\
&\quad + \gamma_{144}w_{u,t,s,19} + \gamma_{127}w_{u,t,s,20} + \gamma_{\{84,103,127,144,222\}}w_{u,t,s,21} = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_4 w_{u,t,s,1} + \gamma_6 w_{u,t,s,2} + \gamma_{14} w_{u,t,s,3} + \gamma_{17} w_{u,t,s,4} + \gamma_{25} w_{u,t,s,5}\\
&\quad + \gamma_{27} w_{u,t,s,6} + \gamma_{39} w_{u,t,s,7} + \gamma_{47} w_{u,t,s,8} + \gamma_{48} w_{u,t,s,9} + \gamma_{60} w_{u,t,s,10}\\
&\quad + \gamma_{61} w_{u,t,s,11} + \gamma_{67} w_{u,t,s,12} + \gamma_{85} w_{u,t,s,13} + \gamma_{83} w_{u,t,s,14}\\
&\quad + \gamma_{78} w_{u,t,s,15} + \gamma_{117} w_{u,t,s,16} + \gamma_{118} w_{u,t,s,17} + \gamma_{105} w_{u,t,s,18}\\
&\quad + \gamma_{146} w_{u,t,s,19} + \gamma_{\{85,118,224\}} w_{u,t,s,20} + \gamma_{\{83,117,124,146,224\}} w_{u,t,s,21} = 0,\\
&\gamma_{19} w_{u,t,s,1} + \gamma_{32} w_{u,t,s,2} + \gamma_7 w_{u,t,s,3} + \gamma_{35} w_{u,t,s,4} + \gamma_{10} w_{u,t,s,5}\\
&\quad + \gamma_{23} w_{u,t,s,6} + \gamma_{46} w_{u,t,s,7} + \gamma_{40} w_{u,t,s,8} + \gamma_{53} w_{u,t,s,9} + \gamma_{62} w_{u,t,s,10}\\
&\quad + \gamma_{71} w_{u,t,s,11} + \gamma_{58} w_{u,t,s,12} + \gamma_{80} w_{u,t,s,13} + \gamma_{87} w_{u,t,s,14}\\
&\quad + \gamma_{76} w_{u,t,s,15} + \gamma_{104} w_{u,t,s,16} + \gamma_{116} w_{u,t,s,17} + \gamma_{102} w_{u,t,s,18}\\
&\quad + \gamma_{145} w_{u,t,s,19} + \gamma_{128} w_{u,t,s,20} + \gamma_{223} w_{u,t,s,21} = 0,\\
&\gamma_{20} w_{u,t,s,1} + \gamma_{31} w_{u,t,s,2} + \gamma_8 w_{u,t,s,3} + \gamma_{34} w_{u,t,s,4} + \gamma_{11} w_{u,t,s,5}\\
&\quad + \gamma_{22} w_{u,t,s,6} + \gamma_{50} w_{u,t,s,7} + \gamma_{41} w_{u,t,s,8} + \gamma_{52} w_{u,t,s,9} + \gamma_{63} w_{u,t,s,10}\\
&\quad + \gamma_{70} w_{u,t,s,11} + \gamma_{65} w_{u,t,s,12} + \gamma_{81} w_{u,t,s,13} + \gamma_{86} w_{u,t,s,14}\\
&\quad + \gamma_{89} w_{u,t,s,15} + \gamma_{119} w_{u,t,s,16} + \gamma_{106} w_{u,t,s,17} + \gamma_{121} w_{u,t,s,18}\\
&\quad + \gamma_{147} w_{u,t,s,19} + \gamma_{\{89,121,225\}} w_{u,t,s,20} + \gamma_{\{89,121,125\}} w_{u,t,s,21} = 0,\\
&\gamma_{33} w_{u,t,s,1} + \gamma_{21} w_{u,t,s,2} + \gamma_{36} w_{u,t,s,3} + \gamma_9 w_{u,t,s,4} + \gamma_{24} w_{u,t,s,5}\\
&\quad + \gamma_{12} w_{u,t,s,6} + \gamma_{51} w_{u,t,s,7} + \gamma_{54} w_{u,t,s,8} + \gamma_{42} w_{u,t,s,9}\\
&\quad + \gamma_{72} w_{u,t,s,10} + \gamma_{64} w_{u,t,s,11} + \gamma_{66} w_{u,t,s,12} + \gamma_{88} w_{u,t,s,13}\\
&\quad + \gamma_{82} w_{u,t,s,14} + \gamma_{90} w_{u,t,s,15} + \gamma_{107} w_{u,t,s,16} + \gamma_{120} w_{u,t,s,17}\\
&\quad + \gamma_{122} w_{u,t,s,18} + \gamma_{148} w_{u,t,s,19} + \gamma_{126} w_{u,t,s,20} + \gamma_{226} w_{u,t,s,21} = 0.
\end{aligned}$$

Computing directly from the above equalities, we obtain

$$\begin{aligned}
(8.9.8.2)\qquad &\gamma_j = 0,\ j = 1, \dots, 90, 99, \dots, 107,\\
&\qquad 114, \dots, 128, 143, 144, \dots, 148, 221 \dots, 226.
\end{aligned}$$

With the aid of (8.9.8.2), the homomorphisms g_1, g_2, g_3, g_4 send (8.9.8.1) to

$$\begin{aligned}
&\gamma_{194} w_{u,t,s,1} + \gamma_{199} w_{u,t,s,2} + \gamma_{91} w_{u,t,s,3} + \gamma_{110} w_{u,t,s,4} + \gamma_{96} w_{u,t,s,5}\\
&\quad + \gamma_{113} w_{u,t,s,6} + \gamma_{204} w_{u,t,s,7} + \gamma_{129} w_{u,t,s,8} + \gamma_{135} w_{u,t,s,9}\\
&\quad + \gamma_{136} w_{u,t,s,10} + \gamma_{141} w_{u,t,s,11} + \gamma_{142} w_{u,t,s,12} + \gamma_{154} w_{u,t,s,13}\\
&\quad + \gamma_{157} w_{u,t,s,14} + \gamma_{160} w_{u,t,s,15} + \gamma_{168} w_{u,t,s,16} + \gamma_{170} w_{u,t,s,17}\\
&\quad + \gamma_{174} w_{u,t,s,18} + \gamma_{187} w_{u,t,s,19} + \gamma_{182} w_{u,t,s,20} + \gamma_{227} w_{u,t,s,21} = 0,\\
&\gamma_{195} w_{u,t,s,1} + \gamma_{198} w_{u,t,s,2} + \gamma_{92} w_{u,t,s,3} + \gamma_{109} w_{u,t,s,4} + \gamma_{97} w_{u,t,s,5}\\
&\quad + \gamma_{112} w_{u,t,s,6} + \gamma_{205} w_{u,t,s,7} + \gamma_{130} w_{u,t,s,8} + \gamma_{134} w_{u,t,s,9} + \gamma_{137} w_{u,t,s,10}\\
&\quad + \gamma_{140} w_{u,t,s,11} + \gamma_{152} w_{u,t,s,12} + \gamma_{155} w_{u,t,s,13} + \gamma_{156} w_{u,t,s,14}\\
&\quad + \gamma_{162} w_{u,t,s,15} + \gamma_{171} w_{u,t,s,16} + \gamma_{169} w_{u,t,s,17} + \gamma_{178} w_{u,t,s,18}\\
&\quad + \gamma_{188} w_{u,t,s,19} + \gamma_{\{162,178,228\}} w_{u,t,s,20} + \gamma_{\{162,178,181\}} w_{u,t,s,21} = 0,
\end{aligned}$$

$$\begin{aligned}
&\gamma_{196}w_{u,t,s,1} + \gamma_{200}w_{u,t,s,2} + \gamma_{93}w_{u,t,s,3} + a_1w_{u,t,s,4} + \gamma_{95}w_{u,t,s,5} + a_2w_{u,t,s,6}\\
&\quad + \gamma_{206}w_{u,t,s,7} + \gamma_{131}w_{u,t,s,8} + a_3w_{u,t,s,9} + \gamma_{138}w_{u,t,s,10} + \gamma_{149}w_{u,t,s,11}\\
&\quad + \gamma_{151}w_{u,t,s,12} + \gamma_{164}w_{u,t,s,13} + \gamma_{158}w_{u,t,s,14} + \gamma_{161}w_{u,t,s,15}\\
&\quad + \gamma_{172}w_{u,t,s,16} + \gamma_{175}w_{u,t,s,17} + \gamma_{177}w_{u,t,s,18} + \gamma_{189}w_{u,t,s,19}\\
&\quad + \gamma_{\{161,164,175,177,183,189,229\}}w_{u,t,s,20} + \gamma_{\{161,177,183\}}w_{u,t,s,21} = 0,\\
&\gamma_{197}w_{u,t,s,1} + \gamma_{201}w_{u,t,s,2} + \gamma_{94}w_{u,t,s,3} + a_4w_{u,t,s,4} + \gamma_{98}w_{u,t,s,5} + a_5w_{u,t,s,6}\\
&\quad + \gamma_{207}w_{u,t,s,7} + \gamma_{132}w_{u,t,s,8} + a_6w_{u,t,s,9} + \gamma_{139}w_{u,t,s,10} + \gamma_{150}w_{u,t,s,11}\\
&\quad + \gamma_{153}w_{u,t,s,12} + \gamma_{165}w_{u,t,s,13} + \gamma_{159}w_{u,t,s,14} + \gamma_{163}w_{u,t,s,15}\\
&\quad + \gamma_{173}w_{u,t,s,16} + \gamma_{176}w_{u,t,s,17} + \gamma_{179}w_{u,t,s,18} + \gamma_{190}w_{u,t,s,19}\\
&\quad + \gamma_{\{163,165,176,179,184,190,230\}}w_{u,t,s,20} + \gamma_{\{163,179,184\}}w_{u,t,s,21} = 0,
\end{aligned}$$

where

$$a_1 = \begin{cases}\gamma_{\{108,218\}}, & s = 3,\\ \gamma_{108}, & s \geqslant 4,\end{cases}\quad a_2 = \begin{cases}\gamma_{\{111,219\}}, & s = 3,\\ \gamma_{111}, & s \geqslant 4,\end{cases}$$

$$a_3 = \begin{cases}\gamma_{\{133,220\}}, & s = 3,\\ \gamma_{133}, & s \geqslant 4,\end{cases}$$

$$a_4 = \begin{cases}\gamma_{\{167,185,186,191,202,209,210,211,212,214,218,231\}}, & s = 3,\\ \gamma_{202}, & s \geqslant 4,\end{cases}$$

$$a_5 = \begin{cases}\gamma_{\{166,167,180,185,192,203,210,213,215,219\}}, & s = 3,\\ \gamma_{203}, & s \geqslant 4,\end{cases}$$

$$a_6 = \begin{cases}\gamma_{\{180,191,193,208,211,216,217,220\}}, & s = 3,\\ \gamma_{208}, & s \geqslant 4.\end{cases}$$

These equalities imply

$$\text{(8.9.8.3)}\qquad \begin{cases}a_i = i,\ i = 1,2,3,4,5,6,\ \gamma_j = 0,\ j = 91,\dots,98,\\ 109,110,112,113,129,\dots,132,134,\dots,141,142,\\ 149,\dots,165,168,\dots,179,181,\dots,184,187,\dots,190,\\ 194,\dots,201,204,\dots,207,227,\dots,230.\end{cases}$$

With the aid of (8.9.8.2) and (8.9.8.3), the homomorphism h sends (8.9.8.1) to

$$\begin{aligned}
&a_7w_{u,t,s,1} + a_8w_{u,t,s,2} + \gamma_{209}w_{u,t,s,3} + \gamma_{166}w_{u,t,s,4} + \gamma_{210}w_{u,t,s,5}\\
&\quad + \gamma_{167}w_{u,t,s,6} + a_9w_{u,t,s,7} + \gamma_{211}w_{u,t,s,8} + \gamma_{180}w_{u,t,s,9}\\
&\quad + \gamma_{212}w_{u,t,s,10} + \gamma_{213}w_{u,t,s,11} + \gamma_{185}w_{u,t,s,12} + \gamma_{214}w_{u,t,s,13}\\
&\quad + \gamma_{215}w_{u,t,s,14} + \gamma_{186}w_{u,t,s,15} + \gamma_{216}w_{u,t,s,16} + \gamma_{217}w_{u,t,s,17}\\
&\quad + \gamma_{191}w_{u,t,s,18} + \gamma_{193}w_{u,t,s,19} + \gamma_{192}w_{u,t,s,20} + \gamma_{231}w_{u,t,s,21} = 0,
\end{aligned}$$

where

$$a_7 = \begin{cases} \gamma_{202}, & s = 3, \\ \gamma_{218}, & s \geqslant 4, \end{cases} \quad a_8 = \begin{cases} \gamma_{203}, & s = 3, \\ \gamma_{219}, & s \geqslant 4, \end{cases} \quad a_9 = \begin{cases} \gamma_{208}, & s = 3, \\ \gamma_{220}, & s \geqslant 4. \end{cases}$$

From the above equalities, it implies

$$\begin{cases} a_7 = a_8 = a_9 = 0,\ \gamma_j = 0,\ j = 166, 167, 180, \\ 185, 186, 191, 192, 193, 209, \ldots, 217, 231. \end{cases} \tag{8.9.8.4}$$

Combining (8.9.8.2), (8.9.8.3) and (8.9.8.4), we get $\gamma_j = 0$ for all j. The proposition is proved. □

9. Final Remark

1. It is well known that the τ-sequence associated with an admissible monomial in P_3 is decreasing. However, this is not true in P_4. The admissible monomials in P_4 with non-decreasing τ-sequences are:

$$(1,2,2,2), (3,1,2,2), (1,3,2,2), (1,2,3,2), (1,2,2,3),$$
$$(1,3,6,6), (3,1,6,6), (3,5,2,6), (3,5,6,2).$$

2. Consider the $s\times 4$-matrices $A_{s,i}$, $1 \leqslant i \leqslant \mu(s)$, where $\mu(1) = 4, \mu(2) = 10, \mu(3) = 14, \mu(s) = 15$ for $s \geqslant 4$, which are determined as follows:

For $s \geqslant 1$,

$$A_{s,1} = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad A_{s,2} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 1 & 1 \end{pmatrix}$$

$$A_{s,3} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & 0 & 1 \end{pmatrix} \quad A_{s,4} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & 1 & 0 \end{pmatrix}.$$

For $s \geqslant 2$,

$$A_{s,5} = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad A_{s,6} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad A_{s,7} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 1 \end{pmatrix}$$

$$A_{s,8} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 1 & 1 \end{pmatrix} \quad A_{s,9} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 1 & 1 \end{pmatrix} \quad A_{s,10} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & 0 & 1 \end{pmatrix}.$$

For $s \geqslant 3$,

$$A_{s,11} = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad A_{s,12} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 1 \end{pmatrix}$$

$$A_{s,13} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad A_{s,14} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 1 & 1 \end{pmatrix}.$$

For $s \geqslant 4$,

$$A_{s,15} = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 1 \end{pmatrix}.$$

3. Consider the $t \times 4$-matrices $B_{t,j}$, $1 \leqslant j \leqslant \rho(t)$, where $\rho(1) = 6, \rho(2) = 20, \rho(t) = 35$ for $t \geqslant 3$, which are determined as follows:

For $t \geqslant 1$,

$$B_{t,1} = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,2} = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 0 & 1 \end{pmatrix} \quad B_{t,3} = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$$B_{t,4} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 0 & 1 \end{pmatrix} \quad B_{t,5} = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 1 & 0 \end{pmatrix} \quad B_{t,6} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & 0 & 0 \end{pmatrix}.$$

For $t \geqslant 2$,

$$B_{t,7} = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,8} = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,9} = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 0 & 1 \end{pmatrix}$$

$$B_{t,10} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,11} = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,12} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 0 & 1 \end{pmatrix}$$

$$B_{t,13} = \begin{pmatrix} 1 & 0 & 1 & 0\\ 0 & 1 & 1 & 0\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 1 & 1 & 0 \end{pmatrix} \quad B_{t,14} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 1 & 0 & 1 \end{pmatrix} \quad B_{t,15} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 0 & 1 & 1 & 0\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$$B_{t,16} = \begin{pmatrix} 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 1 & 0 & 0 & 1 \end{pmatrix} \quad B_{t,17} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 1 & 0 & 0 & 1 \end{pmatrix} \quad B_{t,18} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 1 & 0\\ \vdots & \vdots & \vdots & \vdots\\ 1 & 0 & 1 & 0 \end{pmatrix}.$$

$$B_{t,19} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,20} = \begin{pmatrix} 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 1 & 0 & 1 \end{pmatrix}.$$

For $t \geqslant 3$,

$$B_{t,21} = \begin{pmatrix} 0 & 1 & 1 & 0\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,22} = \begin{pmatrix} 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 1\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,23} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 1 & 0 & 1 \end{pmatrix}$$

$$B_{t,24} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 1 & 0\\ 0 & 1 & 1 & 0\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 1 & 1 & 0 \end{pmatrix} \quad B_{t,25} = \begin{pmatrix} 1 & 0 & 0 & 1\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,26} = \begin{pmatrix} 1 & 0 & 1 & 0\\ 0 & 1 & 1 & 0\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix}$$

$$B_{t,27} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 0 & 1 & 1 & 0\\ 0 & 1 & 0 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 1 & 0 & 1 \end{pmatrix} \quad B_{t,28} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 1 & 0\\ 1 & 0 & 0 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 1 & 0 & 0 & 1 \end{pmatrix} \quad B_{t,29} = \begin{pmatrix} 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix}$$

$$B_{t,30} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 0 & 1 & 0 & 1\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,31} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 0 & 1 & 1 & 0\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,32} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 0 & 1\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix}$$

$$B_{t,33} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 1 & 0\\ 0 & 0 & 1 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 0 & 1 & 1 \end{pmatrix} \quad B_{t,34} = \begin{pmatrix} 1 & 1 & 0 & 0\\ 1 & 0 & 1 & 0\\ 0 & 1 & 0 & 1\\ \vdots & \vdots & \vdots & \vdots\\ 0 & 1 & 0 & 1 \end{pmatrix}.$$

For $t = 3$,

$$B_{3,35} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

For $t \geqslant 4$,

$$B_{t,35} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

4. Consider the $u \times 4$-matrices $C_{u,r}$, $1 \leqslant r \leqslant \mu(u)$, which are determined as follows:

For $u \geqslant 1$,

$$C_{u,1} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad C_{u,2} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$C_{u,3} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 0 & 0 \end{pmatrix} \quad C_{u,4} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

For $u \geqslant 2$,

$$C_{u,5} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad C_{u,6} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad C_{u,7} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$C_{u,8} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad C_{u,9} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad C_{u,10} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

For $u \geqslant 3$,

$$C_{u,11} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad C_{u,12} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$C_{u,13} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad C_{u,14} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

For $u \geqslant 4$,

$$C_{u,15} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

5. Let $A = (a_{ij}), B = (b_{ij})$ be $p \times k, q \times k$ matrices, respectively. Following Kameko [8], we define the $(p+q) \times k$-matrix $A * B = (c_{ij})$ by

$$c_{ij} = \begin{cases} a_{ij}, & \text{if } i \leqslant p, \\ b_{i-p,j}, & \text{if } i > p. \end{cases}$$

Suppose x is an admissible monomial in P_4 with $\tau(x)$ decreasing and $\tau_1(x) < 4$.

5.1. If $\deg(x) = 2^{s+1} - 3$ then the matrix associated with x is of the form $C_{1,j}$ or $A_{s-1,i} * C_{1,j}$ for some i, j.

5.2. If $\deg(x) = 2^{s+1} - 2$ then the matrix associated with x is of the form $B_{s,j}$ for some j.

5.3. If $\deg(x) = 2^{s+1} - 1$ then the matrix associated with x is of the form $C_{s+1,r}$ for some r or $A_{1,i} * B_{s-1,j}$ for some i, j.

5.4. If $\deg(x) = 2^{s+2} + 2^{s+1} - 3$ then the matrix associated with x is of the form $A_{s,i} * C_{2,r}$ for some i, r, or $A_{s+1,i}$ for some i, j.

5.5. If $\deg(x) = 2^{s+t+1} + 2^{s+1} - 3$ for $t \geqslant 2$, then the matrix associated with x is of the form $A_{s,i} * C_{t+1,r}$ for some i, r, or $A_{s+1,i} * B_{t-1,j}$ for some i, j.

5.6. If $\deg(x) = 2^{s+t} + 2^s - 2$ then the matrix associated with x is of the form $B_{s,j} * C_{t,r}$ for some j, r.

5.7. If $\deg(x) = 2^{s+t+u} + 2^{s+t} + 2^s - 3$ then the matrix associated with x is of the form $A_{s,i} * B_{t,j} * C_{u,r}$ for some i, j, r.

References

[1] J. M. Boardman, *Modular representations on the homology of power of real projective space*, Algebraic topology: Oaxtepec 1991, M. C. Tangora (ed.) Contemp. Math. **146** (1993), 49-70. MR1224907 (95a:55041)

[2] R. R. Bruner, L. M. Ha and N. H. V. Hưng, *On behavior of the algebraic transfer*, Trans. Amer. Math. Soc. **357** (2005), 473-487. MR2095619 (2005k:55010)

[3] D. P. Carlisle and R. M. W. Wood, *The boundedness conjecture for the action of the Steenrod algebra on polynomials*, Adams Memorial Symposium on Algebraic Topology 2, N. Ray and G. Walker (ed.) London Math. Soc. Lecture Notes Series **176** (1990), 203-216. MR1232207 (95f:55015)

[4] L. M. Hà, *Sub-Hopf algebras of the Steenrod algebra and the Singer transfer*, in: Proceedings of the International School and Conference in Algebraic Topology, Hà Nội 2004, Geom. Topol. Monogr., Geom. Topol. Publ., Coventry, vol. 11, 2007, 81-105, MR2402802.

[5] N. H. V. Hưng, *The cohomology of the Steenrod algebra and representations of the general linear groups*, Trans. Amer. Math. Soc. **357** (2005), 4065-4089. MR2159700 (2006f:55017)

[6] N. H. V. Hưng and T. N. Nam, *The hit problem for the Dickson algebra*, Trans. Amer. Math. Soc. **353** (2001), 5029-5040. MR1852092 (2002f:55041)

[7] N. H. V. Hưng and T. N. Nam, *The hit problem for the modular invariants of linear groups*, Journal of Algebra **246** (2001), 367-384. MR2872626 (2002j:13008)

[8] M. Kameko, *Products of projective spaces as Steenrod modules*, Thesis, Johns Hopkins University, 1990.

[9] T. N. Nam, *$\mathcal{A}$-générateurs génériquess pour l'algèbre polynomiale*, Adv. Math. **186** (2004), 334-362. MR2073910 (2005e:55029)

[10] F. P. Peterson, *Generators of $H^*(\mathbb{R}P^\infty \times \mathbb{R}P^\infty)$ as a module over the Steenrod algebra*, Abstracts Amer. Math. Soc. No. **833** April 1987.

[11] S. Priddy, *On characterizing summands in the classifying space of a group, I*, Amer. Jour. Math. **112** (1990), 737-748. MR1073007 (91i:55020)

[12] J. H. Silverman, *Hit polynomials and the canonical antimonomorphism of the Steenrod algebra*, Proc. Amer. Math. Soc. **123** (1995), 627-637. MR1254854 (95c:55023)

[13] W. M. Singer, *The transfer in homological algebra*, Math. Zeit. **202** (1989), 493-523. MR1022818 (90i:55035)

[14] W. M. Singer, *On the action of the Steenrod squares on polynomial algebras*, Proc. Amer. Math. Soc. **111** (1991), 577-583. MR1045150 (91f:55007)

[15] N. E. Steenrod, *Cohomology operations*, Lectures by N. E. Steenrod written and revised by D. B. A. Epstein, Annals of Mathematics No. 50, Princeton University Press, Princeton N.J (1962). MR0145525 (26 #3056)

[16] R. M. W. Wood, *Steenrod squares of polynomials and the Peterson conjecture*, Math. Proc. Cambriges Phil. Soc. **105** (1989), 307-309. MR0974986 (90a:55030)

Department of Mathematics, Quy Nhơn University,
170 An Dương Vương, Quy Nhơn, Bình Định, Việt Nam.

E-mail: nguyensum@qnu.edu.vn